

Name of journal: World Journal of Gastroenterology



ESPS Manuscript NO: 15053

Columns: ORIGINAL ARTICLE

Prospective Study

3.0T 1H magnetic resonance spectroscopy in the assessment the changes of steatosis in patients with chronic hepatitis C

Qian Zhang, Hui-Mao Zhang, Wen-Qian Qi, Yong-Gui Zhang, Ping Zhao, Jian Jiao, Jiang-Bin Wang, Chun-Yu Zhang

Match Overview			 
1	CrossCheck 206 words Frank Fischbach. "Assessment of in vivo 1H magnetic resonance spectroscopy in the liver: a review : Assessm ...	4%	
2	CrossCheck 138 words H.A. Marsman. "Noninvasive quantification of hepatic steatosis in rats using 3.0 T 1H-magnetic resonance spec ...	3%	
3	CrossCheck 128 words A. K. P. Lim. "31P MR Spectroscopy in Assessment of Response to Antiviral Therapy for Hepatitis C Virus Rela ...	3%	
4	CrossCheck 72 words Jesper Lundbom. "Long-TE 1H MRS suggests that liver fat is more saturated than subcutaneous and visceral fat"	2%	
5	Internet 63 words crawled on 09-Feb-2010 www.ajronline.org	1%	
6	Internet 50 words crawled on 29-Sep-2014 www.ncbi.nlm.nih.gov	1%	

[网页](#) [图片](#) [新闻](#) [视频](#) [更多](#) [搜索工具](#)

找到约 8,540 条结果 (用时 0.51 秒)

[\[PDF\] Hydrogen 1 \(1H\) magnetic resonance spectroscopy \(MR...](#)www.academicjournals.org/journal/AJB/.../14E34C149266 - [翻译此页](#)

2014年12月31日 - (MRS) of 3.0T in assessment of steatosis to antiviral therapy for ...
Ninety chronic hepatitis C patients undergoing antiviral therapy with interferon and
..... the magnetic resonance peak caused by these changes could be ...

[Publications - Julie A. Fitzpatrick - Imperial College London](#)www.imperial.ac.uk/people/.../publications.html?... - [翻译此页](#)

55 results - We compared in vivo hepatic 31P magnetic resonance spectroscopy
(31P ... agent to assess the severity of hepatitis C virus (HCV)-related liver disease. ...
Twelve healthy volunteers and 15 patients with chronic liver disease were studied. ...
resonance at 3.0 T relative to 1.5 T. There was no significant change in ...

[Publications - Julie A. Fitzpatrick - Imperial College London](#)www.imperial.ac.uk/people/julie.fitzpatrick/publications.html - [翻译此页](#)

55 results - Magnetic Resonance Imaging (MRI) and Spectroscopy (MRS) have ... for
the non-invasive assessment of hepatic fibrosis in chronic hepatitis C, ... (PME)
resonance at 3.0 T relative to 1.5 T. There was no significant change in ... Proton
magnetic resonance spectroscopy (1H MRS) is increasingly used for steatosis ...

[Publications - Julie A. Fitzpatrick - Imperial College London](#)www.imperial.ac.uk/.../publications.html?... - [翻译此页](#)

11 results - We compared in vivo hepatic 31P magnetic resonance spectroscopy
(31P MRS) and ... resonance at 3.0 T relative to 1.5 T. There was no significant
change in nuclear ... A total of 67 patients with biopsy-proven chronic hepatitis C
underwent all four ... Hepatic steatosis was assessed by ultrasound and 1H MRS.

[网页](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 8,540 条结果 (用时 0.51 秒)

[PDF] Hydrogen 1 (1H) magnetic resonance spectroscopy (MRS ...www.academicjournals.org/journal/AJB/.../14E34C149266 - 翻译此页

2014年12月31日 - (MRS) of 3.0T in assessment of steatosis to antiviral therapy for ...

Ninety chronic hepatitis C patients undergoing antiviral therapy with interferon and the magnetic resonance peak caused by these changes could be ...

Publications - Julie A. Fitzpatrick - Imperial College Londonwww.imperial.ac.uk/people/julie.fitzpatrick/publications.html ▾ 翻译此页

55 results - Magnetic Resonance Imaging (MRI) and Spectroscopy (MRS) have ... for the non-invasive assessment of hepatic fibrosis in chronic hepatitis C, ... (PME) resonance at 3.0 T relative to 1.5 T. There was no significant change in ... Proton magnetic resonance spectroscopy (1H MRS) is increasingly used for steatosis ...

Publications - Julie A. Fitzpatrick - Imperial College Londonwww.imperial.ac.uk/people/.../publications.html?... ▾ 翻译此页

55 results - We compared in vivo hepatic 31P magnetic resonance spectroscopy (31P ... agent to assess the severity of hepatitis C virus (HCV)-related liver disease. ... Twelve healthy volunteers and 15 patients with chronic liver disease were studied. ... resonance at 3.0 T relative to 1.5 T. There was no significant change in ...

Advanced MRI Methods for Assessment of Chronic Liver ...[www.ncbi.nlm.nih.gov/.../PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov/.../PubMed Central (PMC)) ▾ 翻译此页

作者：B Taouli - 2009 - 被引用次数：100 - 相关文章

HCV infection accounts for approximately 40% of all chronic liver disease, ... Why some patients with steatosis develop inflammatory and fibrotic changes,

Quantification by 1H localized magnetic resonance spectroscopy and ... Comparison of fat quantification methods: a phantom study at 3.0T. J Magn Reson Imaging.

Quantitative Assessment of Liver Fat with Magnetic ...[www.ncbi.nlm.nih.gov/.../PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov/.../PubMed Central (PMC)) ▾ 翻译此页

作者：SB Reeder - 2011 - 被引用次数：141 - 相关文章