

Name of journal World Journal of Gastroenterology

ESPS Manuscript NO: 16497

Column: EDITORIAL

**Somatostatin analogs for gastric carcinoids: For many, but not all**

Sara Masarova, Alessandra Zilli, Dario Coste

# Abstract

Gastric carcinoids (GCs) are classified as type I, related to hypergastrinemia due to chronic atrophic gastritis (CAG), type II, associated with Zollinger-Ellison syndrome in multiple endocrine neoplasia type 1 (MEN-1), and type III, which is normogastrinemic.

## Match Overview

100% 100%

1	Internet - Citations Internet - Citations www.sagepub.com	2%
2	Internet - Citations Internet - Citations www.sagepub.com	2%
3	Internet - Citations Internet - Citations www.sagepub.com	1%
4	Internet - Citations Internet - Citations www.sagepub.com	1%
5	Internet - Citations Internet - Citations www.sagepub.com	1%



Somatostatin analogs for gastric carcinoids: For many, but not all



网页

图片

新闻

视频

购物

更多 ▾

搜索工具

找到约 328,000 条结果 (用时 0.68 秒)

## Google 学术: Somatostatin analogs for gastric carcinoids: For many, but not all

Carcinoid tumour - Caplin - 被引用次数: 352

Carcinoid tumors of the gut: Our experience over three ... - Marshall - 被引用次数: 93

Pancreatic endocrine tumors: recent advances - Jensen - 被引用次数: 179

## A Review of Carcinoid Cancer | The Carcinoid Cancer ...

www.carcinoid.org/content/review-carcinoid-cancer ▾ 翻译此页

This drug is derived from the naturally occurring hormone somatostatin. ... Large studies of many individuals indicate that the occurrence of insignificant small ... Gastric carcinoids; Carcinoid tumors originating in the stomach are sometimes very special. ... Not all functioning carcinoid tumors produce the same large variety of ...

## Gastric Carcinoid Tumors

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: R Wardlaw - 2008 - 被引用次数: 6 - 相关文章

Gastric carcinoid tumors are unique in that three types have been described based upon ... multiple endocrine neoplasia (MEN) type 1, pernicious anemia, somatostatin, ... Results of all other laboratory tests were essentially within normal limits, .... to develop carcinoid tumors, but this has not been seen in humans.16 Thus, ...

## Carcinoid Tumours - Patient.co.uk

www.patient.co.uk > Professional Reference ▾ 翻译此页

Two thirds of carcinoid tumours occur somewhere in the gastrointestinal tract. ... About 70% of all carcinoids are located in the major bronchi and the remainder ... Most tumours are clinically silent, but they may cause pain, weight loss or present as ... of

找到约 323,000 条结果 (用时 0.62 秒)

## Google 学术 : Somatostatin analogs for gastric carcinoids: For many, but not all

Carcinoid tumour - Caplin - 被引用次数 : 353

Carcinoid tumors of the gut: Our experience over three ... - Marshall - 被引用次数 : 93

Pancreatic endocrine tumors: recent advances - Jensen - 被引用次数 : 179

### A Review of Carcinoid Cancer | The Carcinoid Cancer ...

[www.carcinoid.org/content/review-carcinoid-cancer](http://www.carcinoid.org/content/review-carcinoid-cancer) ▾ 翻译此页

This drug is derived from the naturally occurring hormone **somatostatin**. ... Large studies of **many** individuals indicate that the occurrence of insignificant small ... **Gastric carcinoids**; Carcinoid tumors originating in the stomach are sometimes very special. ... **Not all** functioning carcinoid tumors produce the same large variety of ...

### Gastric Carcinoid Tumors

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > PubMed Central (PMC) ▾ 翻译此页

作者 : R Wardlaw - 2008 - 被引用次数 : 6 - 相关文章

**Gastric carcinoid** tumors are unique in that three types have been described based upon ... **multiple** endocrine neoplasia (MEN) type 1, pernicious anemia, **somatostatin**, ... Results of **all** other laboratory tests were essentially within normal limits, .... to develop carcinoid tumors, **but** this has **not** been seen in humans.<sup>16</sup> Thus, ...

### Carcinoid Tumours - Patient.co.uk

[www.patient.co.uk](http://www.patient.co.uk) > Professional Reference ▾ 翻译此页

Two thirds of **carcinoid** tumours occur somewhere in the **gastrointestinal** tract. ...

However, a small **but** significant proportion are malignant and difficult to manage. ...

About 70% of **all carcinoids** are located in the major bronchi and the remainder in .... of **carcinoid** syndrome do **not** resolve with a **somatostatin analogue** alone.