

Match Overview

Name of Journal: World Journal of Gastroenterology

ISSN Manuscript NO: 14905

Column: ORIGINAL ARTICLE

Basic Study

Activated rat hepatic stellate cells influence Th1/Th2 profile *in vitro*

Zuo Zhu Xing, Liu Ye Huang, Cheng Kong Wu, Hong Yoo, Hong Ma, Li Dong
Jin

Abstract

AIM: To investigate the effects of activated rat hepatic stellate cells (HSCs) on rat Th1/Th2 profile *in vitro*.

METHODS: Growth and survival of activated HSCs and CD4⁺ T lymphocytes cultured alone or together was assessed after 24 or 48 h. CD4⁺ T lymphocytes were then cultured with or without activated HSCs for 24 h or



Activated rat hepatic stellate cells influence Th1/Th2 profile in vitro



网页

图片

视频

购物

新闻

更多 ▾

搜索工具

找到约 42,500 条结果 (用时 0.49 秒)

Google 学术: Activated rat hepatic stellate cells influence Th1/Th2 profile in vitro

Fibrotic disease and the TH1/TH2 paradigm - Wynn - 被引用次数: 790

Identification of novel molecules and pathogenic ... - Shackel - 被引用次数: 120

The regulation of inflammation by galectin - 3 - Henderson - 被引用次数: 145

Hepatic Stellate Cell in Fibrosis: Stellate Cell Activation

www.medscape.com/viewarticle/436461_3 ▾ 翻译此页

Phenotypic features of hepatic stellate cell activation during liver injury and resolution. ...

Another means by which Kupffer cells can influence stellate cells is through ... Th1 cytokines inhibit the development of Th2 cells and Th2 cytokines inhibit ... examined in mice with several different lymphocyte profiles, including T-cell ...

Hepatocyte - Stellate Cell Crosstalk in the Liver Engenders ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: C Coulouarn - 2012 - 被引用次数: 57 - 相关文章

2012年3月14日 - Activated hepatic stellate cells (HSC) play a critical role in fibrogenesis ... by tumor cells but is also influenced by the microenvironment (1-3). In vitro angiogenesis ... LX2 were significantly enriched in a rat model of hepatic fibrosis and were and switch the expression of cytokines toward a Th2 profile.

Effects of Liver-Derived Dendritic Cell Progenitors on Th1 ...

www.jimmunol.org/content/164/3/1346.full - 翻译此页

作者: A Khanna - 2000 - 被引用次数: 180 - 相关文章

2000年2月1日 - Notably, these liver DCp cocultured with allogeneic T cells stained strongly for IL-10. ... 37), which blocks the B7-CD28 costimulatory pathway of T cell

[网页](#) [图片](#) [视频](#) [购物](#) [新闻](#) [更多 ▾](#) [搜索工具](#)

找到约 41,500 条结果 (用时 0.58 秒)

Google 学术 : Activated rat hepatic stellate cells influence Th1/Th2 profile in vitro

Fibrotic disease and the TH1/TH2 paradigm - Wynn - 被引用次数 : 791

Identification of novel molecules and pathogenic ... - Shackel - 被引用次数 : 120

The regulation of inflammation by galectin-3 - Henderson - 被引用次数 : 147

Hepatic Stellate Cell in Fibrosis: Stellate Cell Activation

www.medscape.com/viewarticle/436461_3 ▾ 翻译此页

Phenotypic features of **hepatic stellate cell activation** during **liver** injury and resolution. ... Another means by which Kupffer **cells** can **influence stellate cells** is through ... **Th1** cytokines inhibit the development of **Th2 cells** and **Th2** cytokines inhibit ... examined in mice with several different lymphocyte **profiles**, including T-cell ...

Hepatocyte - Stellate Cell Crosstalk in the Liver - Cancer ...

cancerres.aacrjournals.org/.../0008-5472.CAN-11-3317.full.pdf ▾ 翻译此页

作者 : C Coulouarn - 2012 - 被引用次数 : 58 - 相关文章

tumorigenesis suggests that tumor-stroma crosstalk may likely **influence** the phenotype of tumor **cells** ... **Activation** of **hepatic stellate cells** (HSC) is a key feature of **liver** fibrosis and LX2 cultured alone (LX2-CM) failed to induce in **vitro** angiogenesis (Fig. 4C). and switch the expression of cytokines toward a **Th2 profile**.

Hepatocyte–Stellate Cell Cross-Talk in the Liver Engenders ...

cancerres.aacrjournals.org/.../0008-5472.CAN-11-3317.full.pdf ▾ 翻译此页

作者 : C Coulouarn - 2012 - 被引用次数 : 58 - 相关文章

2012年5月15日 - **Activated hepatic stellate cells** (HSC) play a critical role in fibrogenesis associated ... only determined by tumor **cells** but also **influenced** by the In **vitro** angiogenesis **TH1** and **TH2** cytokines has been reported in the microenvi-