

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 16339

Columns: ORIGINAL ARTICLE

Retrospective Study

Living donor liver transplantation with body-weight more or less than 10 kilograms

Yang SC *et al.*, Pediatric LDLT based on body weight

Sheng-Chun Yang, Chia-Jung Huang, Chao-Long Chen, Chih-Hsien Wang,
Shao-Chun Wu, Tsung-Hsiao Shih, Sin-Ei Juang, Ying-En Lee, Bruno Jawan,
Yu-Feng Cheng, Kwok-Wai Cheng

**Sheng-Chun Yang, Chia-Jung Huang, Chih-Hsien Wang, Shao-Chun Wu,
Tsung-Hsiao Shih, Sin-Ei Juang, Ying-En Lee, Bruno Jawan, Kwok-Wai**

Match Overview

- | Match Number | Source | Words | Similarity |
|--------------|----------|---|------------|
| 1 | Internet | 100 words
crawled on 18-Mar-2015
www.ncbi.nlm.nih.gov | 2% |
| 2 | Internet | 20 words
crawled on 05-Feb-2015
etd.lib.nsysu.edu.tw | <1% |

学术搜索

找到约 4,140 条结果 (用时0.23秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

英语网页

☒ 包括专利☒ 包含引用☐ 创建快讯

Application of reduced-size liver transplants as split grafts, auxiliary orthotopic grafts, and living related segmental transplants.

CE Broelsch, JC Emond, PF Whittington... - Annals of ..., 1990 - ncbi.nlm.nih.gov

... Full size 46/57 (81%) 24% Reduced 18/23 (78%) 17% Split 14/21 (67%) 35% Live 5/5 (100% ... Living

Related Transplantation: Donor and Recipient Outcomes ... The first donor experienced two intraoperative compli- cations, a retractor injury to the spleen and a 5-mm lac- eration in ...

被引用次数: 547 相关文章 所有 6 个版本 引用 保存

[HTML] Living donor liver transplantation for children in Brazil weighing less than 10 kilograms

JS Neto, E Carone, V Pugliese, A Salzedas... - ... transplantation, 2007 - Wiley Online Library

... transplantation from live donors. Lancet 1989; 2: 497. ... S. Living-donor liver transplantation in children. Pediatr Transplant 2002; 6: 43-46. ... Techniques for and outcome of liver transplantation in neonates and infants weighing up to 5 kilograms. J Pediatr Surg 2002; 37: 159-164. ...

被引用次数: 24 相关文章 所有 4 个版本 引用 保存

Noninvasive evaluation of graft steatosis in living donor liver transplantation

M Iwasaki, Y Takada, M Hayashi, S Minamiguchi... - ..., 2004 - journals.lww.com

... for definitive preoperative diagnosis of hepatic steatosis, it is not a universally safe procedure and should not be routinely applied to all living donor candidates ... In a recent study on living liver donors, Mary et al ... 8) reported that BMI was a reliable predictor of hepatic steatosis with ...

被引用次数: 99 相关文章 所有 6 个版本 引用 保存

Body surface area and body weight predict total liver volume in Western adults

JN Vauthey, EK Abdalla, DA Doherty... - ... transplantation, 2002 - Wiley Online Library

... Adam R, Castaing D, Bismuth H. Transplantation of small donor liver in adult recipients. ... Impact of graft size mismatching on graft prognosis in liver transplantation from living donors. ...

Measurement of liver volume and hepatic func- tional reserve as a guide to decision-making ...

被引用次数: 258 相关文章 所有 7 个版本 引用 保存

[HTML] Graft and patient survival after adult live donor liver transplantation compared to a matched cohort who received a deceased donor transplantation

PJ Thuluvath, HY Yoo - Liver transplantation, 2004 - Wiley Online Library

... graft (b) survival by Kaplan-Meier analysis of patients with HCV cirrhosis who had live donor liver transplantation (LDLT, n ... 157 adult-to-adult living donor liver transplantation. ... Germin BI, Marcos A. One hundred consecutive hepatic biopsies in the workup of living donors for right ...

被引用次数: 110 相关文章 所有 7 个版本 引用 保存

[HTML] Vascular reconstruction and complications in living donor liver transplantation in infants wiley.com 中的 [HTML]

nih.gov 中的 [PDF]

wiley.com 中的 [HTML]

researchgate.net 中的 [PDF]

wiley.com 中的 [PDF]

wiley.com 中的 [HTML]

[网页](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 15,100 条结果 (用时 0.60 秒)

Google 学术 : Living donor liver transplantation with body-weight more or less than 10 kilograms

... grafts, and living related segmental transplants. - Broelsch - 被引用次数 : 547

Living donor liver transplantation for children in Brazil ... - Neto - 被引用次数 : 24

... of graft steatosis in living donor liver transplantation - Iwasaki - 被引用次数 : 99

PDF(142K) - Wiley Online Library

onlinelibrary.wiley.com/doi/10.1002/lt.21206/pdf ▾ [翻译此页](#)

作者 : JS Neto - 2007 - 被引用次数 : 24 - [相关文章](#)

in Brazil Weighing Less Than 10 Kilograms. Joao Seda Neto ... Living donor liver transplantation (LDLT) is the only option for timely 4.8-10 kg). The body weight distribution of these unit was 5 days (range, 1-77 days); most patients (85%).

Successful living donor liver transplant in a very small child

www.ncbi.nlm.nih.gov ▸ ... ▸ [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者 : V Kumar - 2010 - [相关文章](#)

It may be either less than or more than 10 kg body weight as the surgical anastomosis in <10 kg child is usually considered difficult. Small infants continued to ...

[PDF] The outcome of primary liver transplantation from

<https://www.rug.nl/research/portal/files/.../Chapter%208> ▾ [翻译此页](#)

primary liver transplantation from deceased donors in our unit. The aim of ... The outcome of liver transplantation in children with body weight ≤ 10 kg | 137 regel ... not age reflects more accurately the size of these children. The aim of results of LRLT in children weighing less than 10 kg did not show a negative impact on.