

No items found.

Search details

Increased[All Fields] AND ("serum"
[MeSH Terms] OR "serum"[All Fields])
AND soluble[All Fields] AND
("scavenger receptors, class e"[MeSH
Terms] OR "class e scavenger

Search

See more...

Recent Activity

Turn Off Clear

Increased serum soluble lectin-like oxidized
low-density lipoprot... (0) PubMed

PubMed Help - PubMed Help

A General Introduction to the E-utilities -
Entrez Programming Utilities Help

Increased risk of second malignancy in
pancreatic intraductal pap... (0) PubMed

16795993[uid] (1) PubMed

See more...

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 16888

Columns: ORIGINAL ARTICLE

Case Control Study

Increased serum soluble lectin-like oxidized low-density lipoprotein receptor-1 levels in patients with biopsy-proven nonalcoholic fatty liver disease

Oguzhan Ozturk, Yasar Colak, Ebubekir Senates, Yusuf Yilmaz, Celal Ulasoglu, Levent Doganay, Seyma Ozkanli, Yasemin Musteri Oltulu, Ender Coskunpinar, Ilyas Tuncer

Match Overview

1	CrossCheck 111 words Colak, Yasar, Ebubekir Senates, Ender Coskunpinar, Yasemin Musteri Oltulu, Ebru Zemheri, Oguzhan Ozturk, Levent Doganay, Seyma Ozkanli, Ilyas Tuncer	3%
2	CrossCheck 104 words Yasar Colak. "Association of Serum Lipoprotein-Associated Phospholipase A2 Level with Nonalcoholic Fatty Liver Disease"	2%
3	CrossCheck 74 words Yasar Colak. "Assessment of endothelial function in patients with nonalcoholic fatty liver disease", Endocrine, 06/2012	2%
4	Internet 46 words crawled on 09-Nov-2012 www.uscnk.us	1%
5	CrossCheck 33 words Ebubekir Senates. "Serum Levels of Hepcidin in Patients with Biopsy-Proven Nonalcoholic Fatty Liver Disease", World Journal of Gastroenterology, 2012	1%
6	CrossCheck 29 words Yasar Colak. "Serum concentrations of human insulin-like growth factor-1 and levels of insulin-like growth factor receptor-1 in patients with nonalcoholic fatty liver disease"	1%

[网页](#)[图片](#)[新闻](#)[视频](#)[更多](#)[搜索工具](#)

找到约 6,630 条结果 (用时 1.00 秒)

[Pro-atherosclerotic markers and cardiovascular risk factors ...](#)[www.ncbi.nlm.nih.gov](#) > ... > Literature > PubMed Central (PMC) - [翻译此页](#)作者: MR Alvares-da-Silva - 2014 - [相关文章](#)

2014年7月14日 - Core tip: Liver transplant (LT) patients a have high risk of long-term ... biopsy-proven nonalcoholic steatohepatitis (NASH) patients and healthy ... The serum cytokine levels (TNF- α , IL-8, IFN γ , and IL-10) were then CRP can induce endothelial lectin-like oxidized low-density lipoprotein (ox-LDL) receptor-1, ...

[Involvement of the TAGE-RAGE system in non-alcoholic ...](#)[www.ncbi.nlm.nih.gov](#) > ... > Literature > PubMed Central (PMC) - [翻译此页](#)作者: M Takeuchi - 2014 - [相关文章](#)

2014年12月27日 - Non-alcoholic fatty liver disease (NAFLD) is a major cause of liver disease around the world. ... TAGE synthesis is increased in NASH, and NASH patients were found to concentrations of 43 patients with a combination of biopsy-proven Horiuchi S. Lectin-like oxidized low density lipoprotein receptor-1 ...

[\[PDF\] 13 march 2015, friday poster no presenter title - APASL ...](#)[www.apasl2015.org/.../Poster%20Presentations_13_03_2015.p...](#) - [翻译此页](#)

2015年3月13日 - TGF- β 1/IL-23 PATHWAY IS UP-REGULATED IN PATIENTS WITH ... THE EVALUATION OF NON-ALCOHOLIC FATTY LIVER DISEASE ... SERUM FERRITIN LEVELS IN PREDICTING HISTOLOGICAL INCREASED SERUM SOLUBLE LECTIN-LIKE OXIDIZED LOW-DENSITY LIPOPROTEIN ...

[\[DOC\] Name of journal: World Journal of Hepatology ESPS ...](#)[www.wjgnet.com/esps/DownloadFile.aspx?Type=M...](#) - [翻译此页](#)作者: M Takeuchi - [相关文章](#)

2014年10月22日 - Non-alcoholic fatty liver disease (NAFLD) is a major cause of liver disease ... TAGE synthesis is increased in NASH, and NASH patients were

[网页](#) [购物](#) [图片](#) [新闻](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 6,710 条结果 (用时 0.57 秒)

[\[PDF\] 13 march 2015, friday poster no presenter title - apasl](#)www.apasl2015.org/.../Poster%20Presentations_13_03_2015.p... [翻译此页](#)

2015年3月13日 - TGF- β 1/IL-23 PATHWAY IS UP-REGULATED IN PATIENTS WITH ... THE EVALUATION OF NON-ALCOHOLIC FATTY LIVER DISEASE ... SERUM FERRITIN LEVELS IN PREDICTING HISTOLOGICAL INCREASED SERUM SOLUBLE LECTIN-LIKE OXIDIZED LOW-DENSITY LIPOPROTEIN ...

[\[PDF\] Role of Scavenger Receptors in the Pathophysiology of ...](#)www.researchgate.net/...receptors...liver_diseases/.../54ae720c0... [翻译此页](#)

作者 : C Armengol - 2013 - 被引用次数 : 4 - 相关文章

Ruti, camí de les escoles s/n, Edifici de Recerca, Planta 1, 08916 Badalona, ...

biomarkers of advanced liver disease, including cirrhosis and cancer. ... low-density lipoprotein; LOX1: lectin-like oxidized LDL receptor; LPS: lipopolysaccharide; LTA: lipoteichoic acid; MΦ: ... non-alcoholic fatty liver disease (NAFLD), which.

[Pro-atherosclerotic markers and cardiovascular risk factors ...](#)[www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC)) - [翻译此页](#)

作者 : MR Alvares-da-Silva - 2014 - 相关文章

2014年7月14日 - Core tip: Liver transplant (LT) patients a have high risk of long-term ... biopsy-proven nonalcoholic steatohepatitis (NASH) patients and healthy ... The serum cytokine levels (TNF- α , IL-8, IFN γ , and IL-10) were then ... CRP can induce endothelial lectin-like oxidized low-density lipoprotein (ox-LDL) receptor-1, ...

[Involvement of the TAGE-RAGE system in non-alcoholic ...](#)[www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC)) - [翻译此页](#)

作者 : M Takeuchi - 2014 - 相关文章

2014年12月27日 - Non-alcoholic fatty liver disease (NAFLD) is a major cause of liver