

Toll like receptor-4 activation upregulates cystathionine β synthetase expressic

学术搜索

获得 4 条结果 (用时0.23秒)

我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

创建快讯

您是不是要找: Toll like receptor-4 activation upregulates cystathionine β **synthase** expression through an NF- κ B-dependent mechanism in a rat model of chronic visceral pain

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Nuclear factor- κ B—a pivotal transcription factor in **chronic** inflammatory diseases

FH Epstein, [PJ Barnes](#), M Karin - New England Journal of ..., 1997 - Mass Medical Soc

... and between the receptor and NF- κ B.28,38-40 Thus, glucocorticoids **activate** glucocorticoid receptors ... NF- κ B is **activated** by many of the factors (viral infection, oxidants, and ... Effects of glucocorticoids on transcription factor **activation** in human peripheral blood mononuclear cells.

被引用次数: 4373 相关文章 所有 8 个版本 引用 保存

[academia.edu](#) 中的 [PDF]

Anti-inflammatory and cytoprotective actions of hydrogen sulfide: translation to therapeutics

[JL Wallace](#), RW Blackler, [MV Chan](#)... - ... & redox signaling, 2014 - online.liebertpub.com

... biopsies from patients with ulcerative colitis, allyl disulfide significantly inhibited NF- κ B **activation** and production ... H₂S has also been shown to **activate** the stress response pathway, up ... of a protein (Keap1) that tonically suppresses nuclear factor (erythroid-derived 2)-like 2 (Nrf2 ...

被引用次数: 12 相关文章 所有 3 个版本 引用 保存

Inflammation, atherosclerosis, and coronary artery disease

[GK Hansson](#) - New England Journal of Medicine, 2005 - Mass Medical Soc

... 1 helper T (Th1) response activates macrophages, initiates an inflammatory response **similar** to delayed hypersensitivity ... **Expression of toll-like** receptors in human atherosclerotic lesions: a possible pathway for ... CD1d-dependent **activation** of NKT cells aggravates atherosclerosis ...

被引用次数: 5885 相关文章 所有 22 个版本 引用 保存

[vtcardsfellows.com](#) 中的 [PDF]

Survival rate= eZ/(1+ eZ) Z= j10. 977+ 0.322 x SBP j0. 019 x HR

C HEMORRHAGIC - journals.lww.com

引用 保存

创建快讯

您是不是要找: Toll like receptor-4 activation upregulates cystathionine β **synthase** expression through an NF- κ B-dependent mechanism in a rat model of chronic visceral

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 17058

Column: ORIGINAL ARTICLE

Basic study

TLR4 upregulates CBS expression through NF- κ B activation in rats with chronic visceral hypersensitivity

Bo Yuan, Wei-Hong Tang, Li-Juan Lu, Yuan Zhou, Hong-Yan Zhu, You-Lang Zhou, Hong-Hong Zhang, Chuang-Ying Hu, Guang-Yin Xu

Abstract

AIM: To investigate the roles of TLR4 and NF- κ B on CBS expression and visceral hypersensitivity in rats.

Match Overview

1	CrossCheck: 137 words Chen R., J. Tian, F. Jiang, X. Zhou, C. Hu, X. Song, C. H. Yao, X. Jiang and G. H. Xu: Thymocyte release inflammation	2%
2	Internet: 116 words Internet: http://www.ncbi.nlm.nih.gov/pmc/articles/PMC2811111/	2%
3	CrossCheck: 35 words Hu, Shuang, Shengqi Su, Yuhua Mao, Yanyang Dai, Li Lu, Zhu, Xiaohong Zhou, Ying Gao and Guang-Yin Xu: TLR4	1%
4	CrossCheck: 17 words Teplovskaya, Ekaterina, Anna C. Pashina, Alexander V. Gerasimov, et al.: Lipopolysaccharide (LPS)-induced NF-κB activation in macrophages	1%
5	CrossCheck: 37 words Zhang H, Wu J, Wu F, Li Z, Zhou Y, Wu Y, Wu J, Wang H, Li H, Li Y, Zhang Y, Li M, Huang Y, Wang H, Li Y, Wu Y, Zhang H	1%
6	CrossCheck: 15 words Zhang, Chunhui, Yanyang Hu, Yanyang Hu, Yanyang Hu, Yanyang Hu, Yanyang Hu, Yanyang Hu, Yanyang Hu, Yanyang Hu, Yanyang Hu	1%
7	CrossCheck: 15 words Tang, Shuang, Shengqi Su, Yuhua Mao, Yanyang Dai, Li Lu, Zhu, Xiaohong Zhou, Ying Gao and Guang-Yin Xu: TLR4	1%
8	CrossCheck: 40 words Tang, Shuang, Shengqi Su, Yuhua Mao, Yanyang Dai, Li Lu, Zhu, Xiaohong Zhou, Ying Gao and Guang-Yin Xu: TLR4	1%



TLR4 upregulates CBS expression through NF- κ B activation in rats with



网页

图片

新闻

视频

更多 ▾

搜索工具

找到约 5,440 条结果 (用时 0.66 秒)

小提示: 仅限搜索简体中文结果。您可以在设置中指定搜索语言

Upregulation of cystathionine β -synthetase by toll-like ...

cpfd.cnki.com.cn/Article/CPFDTOTAL-OGSC201410001336.htm

作者: 徐广银 - 2014

Methods:Visceral hypersensitivity was induced by colonic injection of ... NCI significantly upregulated expression of TLR4,NF- κ B and CBS protein ... Conclusion:Our results suggest that upregulation of CBS expression via activation of toll-like ... on c-Fos expression in the spinal cord and brain of rats with chronic visceral ...

TRPV1 and SP: key elements for sepsis outcome?

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: JV Bodkin - 2013 - 被引用次数: 5 - 相关文章

2013年11月14日 - Determination of their reactivity depends solely on their expression of surface receptors. (H2S) and PPT proteins act together to upregulate NF- κ B and TLR4 in a This resulted in reduced ERK and NF- κ B activation, associated with Cervero F: Deficits in visceral pain and hyperalgesia of mice with a ...

ibuprofen upregulates expressions - WorldWideScience

worldwidescience.org/topicpages/i/ibuprofen+upregulates+expressions.html

Neutrophil recruitment and activation are principal events in inflammation. Interleukin-6 upregulates paraoxonase 1 gene expression via an AKT/NF- κ B ... chronic visceral hypersensitivity in conditions such as post-infectious irritable bowel B, respectively, impaired the TLR4-manipulated VSMC foam cell formation in ...

[PDF] Mikrobiota, Probiotika und Wirt Microbiota, Probiotics and...

www.dahm.org/wissenschaftlichethemenforschung/.../m_242 - 翻译此页

About 6,780 results

Any time

Past hour

Past 24 hours

Past week

Past month

Past year

All results

Verbatim

[TRPV1 and SP: key elements for sepsis outcome?](#)www.ncbi.nlm.nih.gov/pmc/articles/PMC3838676/

Nov 14, 2013 ... Passage of cations **through** this channel, particularly calcium, forms the basis of ... TRPV1 has also shown varying levels of **expression** in a diverse array of sulphide (H₂S) and PPT proteins act together to **upregulate NF-κB** and **TLR4** This resulted in reduced ERK and **NF-κB activation**, associated with ...

[Upregulation of cystathionine β-synthetase by toll-like receptor 4 ...](#)cpfd.cnki.com.cn/Article/CPFDTOTAL-OGSC201410001336.htm

Methods: **Visceral hypersensitivity** was induced by colonic injection of 0.5% acetic ... NCI significantly **upregulated expression** of **TLR4**, **NF-κB** and **CBS** protein levels in ... of **CBS expression** via **activation** of toll-like receptor 4 and **NF-κB** signaling ... on c-Fos **expression** in the spinal cord and brain of **rats with chronic** visceral ...

[Prrxl1 - paired related homeobox protein-like 1 - iHOP](#)www.ihop-net.org/UniPub/iHOP/bng/325731.html ▼

The DRG and sciatic nerve of adult **rats expressed** the hexokinase I isoform only. ... In addition, **NF-kappaB** decoy, but not p38 MAPK inhibitor, SB203580 reduced by AT(1) receptor **activation** by Ang II **through** an autocrine mechanism. ... root ganglion (DRG) neurons in a **rat** model of IBS-like **visceral hypersensitivity**.

[Full Article - Wiley Online Library](#)onlinelibrary.wiley.com/doi/10.1111/1751-2980.12194/full

Oct 20, 2014 ... Enteral nutrition **through** Trelumina placed by endoscopy in severe acute pancreatitis Objective Recent studies have implied that ectopic **activation** of the Wnt SST significantly inhibited highly **expression** of **TLR4**, **NF-κB** and IFN- γ while the **visceral hypersensitivity** induced by **chronic** stress in **rats**.

[\[PDF\] 2008 - Deutsche Gesellschaft für Hygiene und Mikrobiologie](#)www.dghm.org/wissenschaftlichethemenforschung/.../m_242 ▼

VSL#3 attenuates inflammatory gene **expression** in the cecum of IL10KO ... Role of **NF-κB activation** in intestinal epithelial cells in intestinal immune ... intestinal inflammation **through**