



Wiley-Blackwell



Rank	Article Title	Journal	Year	CrossCheck Score
1	Effects of presenting hyperphagic rats with automatic appetitive or satiation cues	Journal of Experimental Psychology: Animal Behavior Processes	1980	5%
2	Interoception: The sense of the state of the body	Philosophical Transactions of the Royal Society B: Biological Sciences	2011	2%
3	How Taster, Taster DNA, Tasteless, Taster's Brain, Taster's Gut, Taster's Personality, Taster's Body, Taster's Mind	Journal of Experimental Psychology: Animal Behavior Processes	1980	2%
4	How Taster, Taster DNA, Tasteless, Taster's Brain, Taster's Gut, Taster's Personality, Taster's Body, Taster's Mind	Journal of Experimental Psychology: Animal Behavior Processes	1980	1%
5	How Taster, Taster DNA, Tasteless, Taster's Brain, Taster's Gut, Taster's Personality, Taster's Body, Taster's Mind	Journal of Experimental Psychology: Animal Behavior Processes	1980	1%
6	How Taster, Taster DNA, Tasteless, Taster's Brain, Taster's Gut, Taster's Personality, Taster's Body, Taster's Mind	Journal of Experimental Psychology: Animal Behavior Processes	1980	1%
7	How Taster, Taster DNA, Tasteless, Taster's Brain, Taster's Gut, Taster's Personality, Taster's Body, Taster's Mind	Journal of Experimental Psychology: Animal Behavior Processes	1980	1%
8	How Taster, Taster DNA, Tasteless, Taster's Brain, Taster's Gut, Taster's Personality, Taster's Body, Taster's Mind	Journal of Experimental Psychology: Animal Behavior Processes	1980	<1%



Female Spontaneously Diabetic Torii fatty rats develop Nonalcoholic steatosis



网页

图片

新闻

视频

更多

搜索工具

找到约 6,340 条结果 (用时 0.94 秒)

Development of novel rat model for high-fat and high ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: K Kitamori - 2012 - 被引用次数: 11 - 相关文章

2011年8月19日 - Non-alcoholic fatty liver disease (NAFLD) includes a progressive liver disorder ... and the fibrosis results in hepatic cirrhosis or, finally, in tumor formation. ... The stroke-prone spontaneously hypertensive rat, SHRSP5/Dmcr, ... responses (about 600–900 mg/dl for females and 300–600 mg/dl for ... Diabetes.

Obesity, Inflammation and Liver Cancer

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: B Sun - 2012 - 被引用次数: 115 - 相关文章

2011年11月25日 - NASH is characterized by fatty liver inflammation and is believed to cause ... syndrome, type 2 diabetes, non-alcoholic fatty liver disease (NAFLD), and the ... mice spontaneously develop obesity even on normal chow [77–79]. ... in induction of pre-neoplastic liver lesions in DEN-administered rats [141].

The Role of Metformin in the Management of NAFLD

www.hindawi.com/journals/jdr/2012/716404/ ▾ 翻译此页

作者: A Mazza - 2011 - 被引用次数: 71 - 相关文章

2011年10月4日 - Nonalcoholic fatty liver disease (NAFLD) is the most common liver ... NAFLD is characterized by liver damage and functional impairment similar to those observed in ... risk factor even in patients with a normal body mass index (BMI). Diet ... a potential role of genetic factors for the development of NAFLD.

Related articles

www.hitachi-aloka.co.jp/english/products/.../lct200_03.html ▾ 翻译此页

Persistent hepatitis C virus infections and hepatopathological manifestations in ...



网页 图片 购物 新闻 视频 更多 ▾ 搜索工具

找到约 11,400 条结果 (用时 1.22 秒)

The Role of Metformin in the Management of NAFLD

www.hindawi.com/journals/jdr/2012/716404/ ▾ 翻译此页

作者：A Mazza - 2011 - 被引用次数：71 - 相关文章

2011年10月4日 - **Nonalcoholic fatty liver** disease (NAFLD) is the most common **liver** ...
NAFLD is characterized by **liver** damage and functional impairment **similar** to those
observed in ... risk factor even in patients with a normal body **mass** index (BMI). Diet ...
a potential role of genetic factors for the **development** of NAFLD.

Development of novel rat model for high-fat and high ...

www.ncbi.nlm.nih.gov ▸ ... ▸ PubMed Central (PMC) ▾ 翻译此页

作者：K Kitamori - 2012 - 被引用次数：11 - 相关文章

2011年8月19日 - **Non-alcoholic fatty liver** disease (NAFLD) includes a progressive **liver**
disorder ... and the fibrosis results in **hepatic** cirrhosis or, finally, in **tumor** formation. ...
The stroke-prone **spontaneously** hypertensive **rat**, SHRSP5/Dmcr, ... responses (about
600–900 mg/dl for **females** and 300–600 mg/dl for **Diabetes**.

Obesity, Inflammation and Liver Cancer

www.ncbi.nlm.nih.gov ▸ ... ▸ PubMed Central (PMC) ▾ 翻译此页

作者：B Sun - 2012 - 被引用次数：118 - 相关文章

2011年11月25日 - NASH is characterized by **fatty** liver inflammation and is believed to
cause ... syndrome, type 2 **diabetes**, **non-alcoholic fatty** liver disease (NAFLD), and
the mice **spontaneously develop** obesity even on normal chow [77–79]. in
induction of pre-neoplastic **liver lesions** in DEN-administered **rats** [141].

Related articles

www.hitachi-aloka.co.jp/english/products/.../lct200_03.html ▾ 翻译此页

Persistent **hepatitis** C virus infections and hepatopathological manifestations in ...
Matsumoto A. Fish oil prevents excessive **hepatic** lipid accumulation without ...
Asakura H, Yoneyama H. A murine model for **non-alcoholic steatohepatitis** M.
Pathophysiological analysis of **female Spontaneously Diabetic Torii fatty rats**.