

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 17183

Columns: ORIGINAL ARTICLE

Retrospective Cohort Study

Diagnostic value of drain amylase for detecting intrathoracic leakage after esophagectomy

Berkelmans GHK *et al.* Amylase detection of leakage after esophagectomy

Gijs HK Berkelmans, Ewout A Kouwenhoven, Boudewijn JJ Smeets, Teus J Weijs, Luis C Silva Corten, Marc J van Det, Grard AP Nieuwenhuijzen, Misha DP Luyer

Gijs HK Berkelmans, Boudewijn JJ Smeets, Teus J Weijs, Grard AP Nieuwenhuijzen, Misha DP Luyer, Department of Surgery, Catharina Hospital, Michelangelolaan 2, 5623 EJ Eindhoven, The Netherlands

Match Overview

1 Internet 114 words
Crawled on 04-May-2015
www.wjgnet.com

2%

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

英语网页

☒ 包括专利☒ 包含引用☐ 创建快讯

Systematic review of the definition and measurement of anastomotic leak after gastrointestinal researchgate.net 中的 [PDF] surgery

J Bruce, ZH Krukowski, G Al-Khairi... - British Journal of ..., 2001 - Wiley Online Library
... sensitiv-ity, speci@city, inter- and intra-rater reliability, practicality and acceptability of the de@nition, measure or diagnostic test. ... Reference Description Drain fluid volume Amylase value
Timing Bottger et al.41 Pancreatic fistula D Drain fluid amylase level > 2000 units/l Daily ...
被引用次数: 363 相关文章 所有 7 个版本 引用 保存

Prospective trial comparing contrast swallow, computed tomography and endoscopy to identify anastomotic leak following oesophagogastric surgery

BA Hogan, D Winter, D Broe, P Broe, MJ Lee - Surgical endoscopy, 2008 - Springer
... There were 29 males and 9 females, with a median age of 61 (range 28-75) years. ... of the surgical anastomosis provided by the referring surgeon is invaluable for accurate diagnosis by the ... A, Busch C, Bause H, Izbickei JR (1996) Gastric to- nometry and drain amylase analysis in ...
被引用次数: 37 相关文章 所有 9 个版本 引用 保存

Chylous ascites after pancreaticoduodenectomy: introduction of a grading system

NA van der Gaag, AC Verhaar, EB Haverkort... - Journal of the American ..., 2008 - Elsevier
... of the treating surgeon, in one of these patients also after a (diagnostic) paracentesis ... Female gender and a pathologic diagnosis of (focal) chronic pancreatitis were independently associated with ... In these circumstances, to diagnose CA, based on TG level and appearance (milky ...
被引用次数: 36 相关文章 所有 5 个版本 引用 保存

Earlier prediction of anastomotic insufficiency after thoracic esophagectomy by intramucosal pH

T Tarui, A Murata, Y Watanabe, SP Kim... - Critical care ..., 1999 - journals.lww.com
... viability of the gastric tube, especially when combined with the rectal pH values. ... The actual pH value is calculated theoretically by a modified Henderson-Hasselbalch equation ... Early detection of impaired perfusion enables caregivers to provide appropriate treatment to improve ...
被引用次数: 22 相关文章 所有 4 个版本 引用 保存

Does continuous mucosal partial carbon dioxide pressure measurement predict leakage of researchgate.net 中的 [PDF] intrathoracic esophagogastronomy?

W Schröder, D Stippel, M Lacher, CA Gutschow... - The Annals of thoracic ..., 2002 - Elsevier
... hours revealed a sensitivity of 0.6, a specificity of 0.88, and a positive predictive value of 0.36 ... advantage of continuous pCO₂ i measurement—that it serves as an early diagnostic indicator of ... At the time of diagnosis, endoscopy revealed a small leak but a well-perfused mucosa of ...



of drain amylase for detecting intrathoracic leakage after esophagectomy



网页

图片

购物

视频

新闻

更多 ▾

搜索工具

找到约 4,720 条结果（用时 0.85 秒）

Google 学术 : Diagnostic value of drain amylase for detecting intrathoracic leakage after esophagectomy

... definition and measurement of anastomotic leak after ... - Bruce - 被引用次数 : 368

... and endoscopy to identify anastomotic leak following ... - Hogan - 被引用次数 : 39

... ascites after pancreaticoduodenectomy: introduction of ... - van der Gaag - 被引用次数 : 35

The value of amylase level measurements in drain ... - nvge

<https://www.nvge.nl/ewiseabstract2?func...type...> ▾ 翻译此页

The value of amylase level measurements in drain fluid for early diagnosis of intrathoracic leakage after esophagectomy. ... Several diagnostic modalities are available to detect anastomotic leakage such as CT-scan and endoscopy. However ...

Relationship of drain amylase and anastomotic leak after ...

meetinglibrary.asco.org/content/106169-133 ▾ 翻译此页

Relationship of drain amylase and anastomotic leak after esophagectomy. ... Elevated drain amylase was defined as any level > 400 IU/L. Chi-square and ... Stapled intra-thoracic anastomoses were performed in 47, 3 had a cervical anastomoses. ... prediction for the detection of anastomotic leak after esophagectomy.

缺少字词 : diagnostic

Gijs Berkelmans | LinkedIn

<https://www.linkedin.com/pub/gijs-berkelmans/a6/32a/4b1> - 翻译此页

Eindhoven Area, Netherlands - Arts-onderzoeker heelkunde bij Catharina Hospital Eindhoven

The value of amylase level measurements in drain fluid for early diagnosis of intrathoracic leakage after esophagectomy. March 2014 – Present. Retrospective ...

Intrathoracic Leaks Following Esophagectomy Are No ...