



Comparison of two types of colectomy in treating slow transit constipation



网页

图片

新闻

视频

更多 +

搜索工具

找到约 3,640 条结果 (用时 0.80 秒)

Google 学术: Comparison of two types of colectomy in treating slow transit constipation with or without melanosis coli

... of young women: 'idiopathic slow transit constipation' - Preston - 被引用次数: 379

... types of subtotal colectomy for slow-transit constipation ... - Feng - 被引用次数: 32

Myths and misconceptions about chronic constipation - Müller-Lissner - 被引用次数: 371

Surgical Outcomes after Total Colectomy with Ileorectal ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: G Sohn - 2011 - 被引用次数: 25 - 相关文章

2011年8月31日 - Keywords: Colonic inertia, Colectomy, Treatment outcome.

Postoperative complication ... physiological studies: 1) the normal transit constipation group without abnormality in ... This study analyzed 37 patients with slow transit constipation who In addition, melanosis coli was shown in three patients. Fig. 2.

[PDF] Surgical Outcomes after Total Colectomy with Ileorectal ...

www.researchgate.net/...colectomy...slow_transit_constipation/ ... ▾ 翻译此页

作者: G Sohn - 2011 - 被引用次数: 25 - 相关文章

ity to treat medically intractable slow transit constipation. 2. 10-20. 3. More than 20. 4. Assistance: type of assistance. Without assistance. 0 ... In addition, melanosis coli was shown in three [10] compared the number of ganglion cells and.

Chronic Constipation: An Evidence-Based Review - The ...


www.jabfm.org/content/24/4/436.full ▾ 翻译此页

作者: L Leung - 2011 - 被引用次数: 94 - 相关文章

Chronic constipation is either caused by slow colonic transit or pelvic floor ... into 2



colectomy in treating slow transit constipation with or without melanosis coli



网页 图片 新闻 视频 购物 更多 搜索工具

找到约 1,300 条结果 (用时 0.78 秒)

Google 学术 : Comparison of two types of colectomy in treating slow transit constipation with or without melanosis coli

... of young women:'idiopathic slow transit constipation' - Preston - 被引用次数 : 378

... types of subtotal colectomy for slow-transit constipation ... - Feng - 被引用次数 : 32

Myths and misconceptions about chronic constipation - Müller-Lissner - 被引用次数 : 367

Gastrointestinal neuromuscular pathology in chronic ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) > 翻译此页

作者 : CH Knowles - 2011 - 被引用次数 : 30 - 相关文章

Some patients with chronic constipation may undergo colectomy yielding tissue ... Are tissues studied from slow transit constipation representative of the ... with silver nitrate revealing two types of neurons: argyrophilic and argyrophobic. ... has failed to identify a persistent abnormality other than melanosis coli [10,21,39] with ...

Treatment of Constipation in Older Adults - American Family ...

www.aafp.org > ... > aafp > Vol. 72/No. 11(December 1, 2005) > 翻译此页

2005年12月1日 - Functional constipation is diagnosed when no secondary causes can be ... Two or more of the following for at least 12 weeks in the preceding 12 months: ... Slow transit constipation is characterized by prolonged delay in the passage of stool pigmentation of the colonic mucosa known as melanosis coli.

[PDF] Chronic Constipation

www.motilitysociety.org/.../lembo_camilleri_NEJM_constipatio... > 翻译此页

作者 : NT Constipation - 2003 - 被引用次数 : 691 - 相关文章

uation and treatment of patients whose constipation is chronic and severe and does not resolve with the ... normal-transit constipation, slow-transit constipation, and disorders of defecatory or ... Two or more of the following for at least 12 weeks (not necessarily consecu- Melanosis coli does not lead to the development of.