



Thyroid Dysfunction in Chinese HCV patients: the Role of TPOAb and CXC



学术搜索

获得 2 条结果 (用时0.06秒)

我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

[The presence of thyroid peroxidase antibody of IgG2 subclass is a risk factor for thyroid dysfunction in chronic hepatitis C patients](#)

C Shao, N Huo, L Zhao, Y Gao, X Fan, Y Zheng... - European Journal of ..., 2013 - EFES

... of thyroglobulin antibody (TGA) in patients with (TD) or without (WTD) thyroid dysfunction. ... IgG2 subclass was significantly higher in hypothyroidism (51.5%) and subclinical hypothyroidism (33.3%) than in ... and differed in effector function from IgG1 and IgG4 thyroid antibodies. ...

被引用次数: 5 相关文章 所有 7 个版本 引用 保存

[eje-online.org 中的 \[HTML\]](#)

[A prospective, randomized trial of intravenous glucocorticoids therapy with different protocols for patients with Graves' ophthalmopathy](#)

W Zhu, L Ye, L Shen, Q Jiao, F Huang... - The Journal of ..., 2014 - press.endocrine.org

... 2, Thyroid dysfunction. 3, Proptosis. 4, Optic nerve involvement. ... Graves' hyperthyroidism, 33 (84.62), 32 (91.43), .27. Primary hypothyroidism, 2 (5.13), 0, .49. ... Visual acuity, 0.91 ± 0.27, 0.96 ± 0.38, .48. Abbreviations: BMI, body mass index; TPOAb, thyroid peroxidase autoantibodies ...

被引用次数: 7 相关文章 所有 4 个版本 引用 保存

[endocrine.org 中的 \[HTML\]](#)

创建快讯

[关于 Google 学术搜索](#)

[隐私权](#)

[条款](#)

[提供反馈](#)

## Match Overview

1	Internet 209 words crawled on 22-Oct-2014 <a href="http://www.eje-online.org">www.eje-online.org</a>	3%
2	CrossCheck 180 words J-F. Huang. "The role of thyroid autoantibodies in the development of thyroid dysfunction in Taiwanese chronic l ...	3%
3	Internet 92 words crawled on 10-Nov-2009 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	1%
4	Internet 65 words crawled on 15-Jul-2010 <a href="http://www.peoriaendocrine.com">www.peoriaendocrine.com</a>	1%
5	CrossCheck 54 words M. Rotondi. "Serum CXCL10 levels and occurrence of thyroid dysfunction in patients treated with interferon- the ...	1%
6	CrossCheck 39 words Sadat, U., S. R. Walsh, A. G. Norden, J. H. Gillard, and J. R. Boyle. "Does Oral N-Acetylcysteine Reduce Contrast-I	1%
7	Publications 36 words Barut, Sener, Gunal, Ozgur, Erkorkmaz, Unal and Yildiz, Feyza. "Thyroid dysfunction in Turkish patients with chro	1%
8	CrossCheck 10 words C Carella. "The addition of ribavirin to interferon-alpha therapy in patients with hepatitis C virus-related chronic he	<1%

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 18162

Manuscript Type: Retrospective Study

Thyroid Dysfunction in Chinese HCV Patients: Prevalence and Correlation of TPOAb and CXCL10

Ren-Wen Zhang, Cui-Ping Shao, Na Huo, Min-Ran Li, Hong-Li Xi, Min Yu,

Xiao-Yuan Xu

### Abstract

**AIM:** To investigate the relationship among pretreatment serum CXC chemokine ligand 10 (CXCL10), thyroid peroxidase antibody (TPOAb) levels and thyroid dysfunction (TD) in Chinese hepatitis C virus (HCV) patients.





Thyroid dysfunction in Chinese hepatitis C virus patients: Prevalence



网页

图片

新闻

视频

购物

更多

搜索工具

找到约 1,670 条结果 (用时 0.81 秒)

## Google 学术: Thyroid dysfunction in Chinese hepatitis C virus patients: Prevalence and correlation of TPOAb and CXCL10

... for thyroid dysfunction in chronic hepatitis C patients - Shao - 被引用次数: 5

## Possible factors affecting thyroid dysfunction in hepatitis C ...

www.ncbi.nlm.nih.gov › ... › Literature › PubMed Central (PMC) - 翻译此页

作者: XR MAO - 2014 - 被引用次数: 5 - 相关文章

2014年5月14日 - In China, a country with a high incidence of HCV infection, there are ~40 ... the correlation between TD and HCV-infected patients without interferon ...

Hypothyroidism is more common in patients with chronic HCV infection than ... The prevalence of TD in HCV-infected patients has been frequently reported; ...

## The presence of thyroid peroxidase antibody of IgG2 ...

www.eje-online.org/content/168/5/717.full.pdf - 翻译此页

作者: C Shao - 2013 - 被引用次数: 5 - 相关文章

Objective: To investigate the prevalence of thyroid dysfunction (TD) and IgG subclasses of ... and twelve untreated hepatitis C virus-infected patients without a history of ... factors were statistically analyzed to determine the correlation between TD and ... The positive percentage of TPOAb IgG2 subclass in the TD group was ...

## The presence of thyroid peroxidase antibody of IgG2 ...

www.eje-online.org/content/168/5/717.full - 翻译此页

作者: C Shao - 2013 - 被引用次数: 5 - 相关文章



atitis C virus patients: Prevalence and correlation of TPOAb and CXCL10



[网页](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 2,050 条结果 ( 用时 0.66 秒 )

## Google 学术 : Thyroid dysfunction in Chinese hepatitis C virus patients: Prevalence and correlation of TPOAb and CXCL10

... for thyroid dysfunction in chronic hepatitis C patients - Shao - 被引用次数 : 5

### Possible factors affecting thyroid dysfunction in hepatitis C ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者 : XR MAO - 2014 - 被引用次数 : 5 - [相关文章](#)

2014年5月14日 - In **China**, a country with a high incidence of **HCV** infection, there are ~40 million individuals infected with **HCV**, with a **prevalence** of ~3.2% (2). ... the **correlation** between TD and **HCV**-infected **patients** without interferon ...

**Hypothyroidism** is more common in **patients** with chronic **HCV** infection than in normal ...

### The presence of thyroid peroxidase antibody of IgG2 ...

[www.eje-online.org/content/168/5/717.full.pdf](http://www.eje-online.org/content/168/5/717.full.pdf) - [翻译此页](#)

作者 : C Shao - 2013 - 被引用次数 : 5 - [相关文章](#)

Objective: To investigate the **prevalence** of **thyroid dysfunction** (TD) and IgG subclasses of ... and twelve untreated **hepatitis C virus**-infected **patients** without a history of ... factors were statistically analyzed to determine the **correlation** between TD and ... The positive percentage of **TPOAb** IgG2 subclass in the TD group was ...

### The presence of thyroid peroxidase antibody of IgG2 ...

[www.eje-online.org/content/168/5/717.full](http://www.eje-online.org/content/168/5/717.full) - [翻译此页](#)

作者 : C Shao - 2013 - 被引用次数 : 5 - [相关文章](#)

2013年5月1日 - Objective To investigate the **prevalence** of **thyroid dysfunction** (TD) and IgG ... analyzed to determine the **correlation** between TD and this **patient**