



Cutaneous Metastasis as Initial Presentation of a High Grade Non-functi



网页

图片

新闻

视频

购物

更多 ▾

搜索工具

找到约 960,000 条结果 (用时 0.76 秒)

Google 学术: Cutaneous Metastasis as Initial Presentation of a High Grade Non-functioning Pancreatic Neuroendocrine Tumor

... : well-differentiated pancreatic non-functioning tumors - Falconi - 被引用次数: 135

... biology and histopathology of neuroendocrine tumours - Klöppel - 被引用次数: 158

... pancreatic nonfunctioning tumors/carcinoma - Jean - 被引用次数: 165

Pancreatic neuroendocrine tumors: biology, diagnosis, and ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: C Ro - 2013 - 被引用次数: 5 - 相关文章

The clinical presentation, diagnosis, and treatment of specific types of PNETs and familial Functioning pancreatic neuroendocrine tumor (PNET) syndromes ... Surgical removal of primary, non-metastatic PNET is the only clinical cure, and surgical ... Chemotherapy is reserved for intermediate- and high-grade PNETs.

Metastatic pancreatic neuroendocrine tumors and poorly ...

www.uptodate.com/.../metastatic-pancreatic-neuroendocrine-tu... ▾ 翻译此页

2015年1月8日 - (See 'High-grade well- or moderately-well differentiated tumors' below and ... in the treatment of functional and nonfunctional neuroendocrine tumors. In situ radiotherapy with ¹¹¹In-pentetreotide: initial observations and future directions. neuroendocrine tumors: Presentation, prognosis, imaging, and ...

BMC Cancer | Full text | Multiple scalp metastases from ...

www.biomedcentral.com/1471-2407/14/305 ▾ 翻译此页

作者: S Wang - 2014 - 相关文章

2014年5月1日 - Cutaneous metastases of colonic neuroendocrine carcinomas ... therefore, cutaneous metastases as initial distant metastases can be ... presentation

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 18489

Columns: CASE REPORT

Cutaneous metastasis as initial presentation of a non-functioning pancreatic neuroendocrine tumor

Woo Young Shin, Keon Young Lee, Seung Ik Ahn, Shin-Young Park, Keun-Myoung Park

Abstract

Non-functioning pancreatic neuroendocrine tumors (NF-PNETs) are rare tumors and they account for 2% of all pancreatic malignancy. About 60% of NF-PNETs present distant metastases and usually hepatic metastases. However, cutaneous metastases are very rare. Herein, we reported our experience. A 60-year-old man visited our outpatient clinic with a mass on his left hip. An abdominal computerized tomography scan

Match Overview

1	CrossCheck 26 words Jan Franko. "Non-functional Neuroendocrine Carcinoma of the Pancreas: Incidence, Tumor Biology, and Outcomes ...	1%
2	CrossCheck 25 words Modlin, I.M.. "Gastroenteropancreatic neuroendocrine tumors", Lancet Oncology, 200801	1%
3	Internet 22 words crawled on 06-Mar-2014 www.carcinoid.org	1%
4	CrossCheck 19 words Lee F Starker. "Molecular genetics of gastroenteropancreatic neuroendocrine tumors", Current Opinion in Oncology, 0	1%
5	Internet 19 words crawled on 08-Dec-2010 www.wjso.com	1%
6	CrossCheck 14 words Jeon, Young Bae, Sangchul Yum, and Dongho Choi. "Transfusion free radical antegrade modular pancreaticosplenect	1%



Cutaneous metastasis as initial presentation of a non-functioning pancreatic



网页

图片

新闻

视频

购物

更多

搜索工具

找到约 303,000 条结果 (用时 0.68 秒)

Google 学术: Cutaneous metastasis as initial presentation of a non-functioning pancreatic neuroendocrine tumor

Endocrine tumours of the pancreas - Öberg - 被引用次数: 372

Neuroendocrine Pancreatic Tumors Clinical Findings ... - Eriksson - 被引用次数: 115

... of gastroenteropancreatic neuroendocrine tumours (... - Öberg - 被引用次数: 160

Endocrine tumors of the pancreas - Surgical Treatment ...

www.ncbi.nlm.nih.gov › NCBI › Literature › Bookshelf ▾ 翻译此页

作者: RG Holzheimer - 2001 - 相关文章

Endocrine tumors of the pancreas (ETP) or islet cell tumors are rare lesions, the ... both for their dramatic and sometimes puzzling initial clinical presentation, and ... ETP are classified as functioning or non functioning according to the type and ... after resection of an apparently benign tumor, by the outcome of metastases.

Islet Cell Tumors of the Pancreas / Endocrine ... - Pathology

pathology.jhu.edu/pc/treatmentendocrine.php?area=tr ▾ 翻译此页

First, islet cell tumors / endocrine neoplasms grow much slower than does pancreatic cancer. ... more islet cell tumors / pancreatic endocrine neoplasms are non-functional. The liver is carefully assessed for evidence of metastatic disease. may be suggested by the clinical presentation and biopsy of the skin lesions, but ...

Pancreatic Endocrine Tumours. Learn about PETs | Patient



initial presentation of a non-functioning pancreatic neuroendocrine tumor



网页

图片

新闻

视频

购物

更多 ▾

搜索工具

找到约 1,730,000 条结果 (用时 0.49 秒)

Google 学术 : Cutaneous metastasis as initial presentation of a non-functioning pancreatic neuroendocrine tumor

Endocrine tumours of the pancreas - Öberg - 被引用次数 : 372

Neuroendocrine Pancreatic Tumors Clinical Findings ... - Eriksson - 被引用次数 : 115

... of gastroenteropancreatic neuroendocrine tumours (... - Öberg - 被引用次数 : 162

Endocrine tumors of the pancreas - Surgical Treatment ...

www.ncbi.nlm.nih.gov > NCBI > Literature > Bookshelf ▾ 翻译此页

作者 : RG Holzheimer - 2001 - 相关文章

Endocrine tumors of the pancreas (ETP) or islet cell tumors are rare lesions, the ... both for their dramatic and sometimes puzzling initial clinical presentation, and ... ETP are classified as functioning or non functioning according to the type and ... after resection of an apparently benign tumor, by the outcome of metastases.

Pancreatic neuroendocrine tumors: biology, diagnosis, and ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者 : C Ro - 2013 - 被引用次数 : 5 - 相关文章

The clinical presentation, diagnosis, and treatment of specific types of PNETs and familial Functioning pancreatic neuroendocrine tumor (PNET) syndromes ... Surgical removal of primary, non-metastatic PNET is the only clinical cure, and The initial treatment of VIPoma entails replacing large fluid losses, correcting ...

Pancreatic Endocrine Tumours. Learn about PETs | Patient

patient.info > Professional Reference ▾ 翻译此页

Epidemiology; Presentation; Investigations; Management; Prognosis; References ...

Liver metastases are the most common secondaries beyond spread to regional ... type