

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 17699

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Computed tomography and magnetic resonance imaging features of
lipid-rich neuroendocrine tumors of the pancreas

Fukukura Y *et al.* CT and MRI of Lipid-rich PanNETs

Yoshihiko Fukukura, Toshikazu Shindo, Michiyo Higashi, Koji Takumi,
Tomokazu Umanodan, Tomohide Yoneyama, Takashi Yoshiura

Yoshihiko Fukukura, Toshikazu Shindo, Koji Takumi, Tomokazu
Umanodan, Tomohide Yoneyama, Takashi Yoshiura, Department of
Radiology, Kagoshima University Graduate School of Medical and Dental
Sciences, Kagoshima City 890-8544, Japan

Match Overview

1	Internet 98 words crawled on 01-Jul-2015 www.wjgnet.com	1%
2	CrossCheck 76 words Kengo Yoshimitsu. "Papillary renal carcinoma: Diagnostic approach by chemical shift gradient-echo and echo-planar ..."	1%
3	CrossCheck 28 words Yoshihiko Fukukura. "Comparison of ferucarbotran-enhanced fluid-attenuated inversion-recovery echo-planar, T2-weighted ..."	<1%
4	CrossCheck 27 words Koji Takumi. "Epithelial-myoepithelial carcinoma of the parotid gland: correlation of dynamic magnetic resonance imaging ..."	<1%
5	CrossCheck 22 words Souya Nunobe. "Clear Cell Endocrine Tumor of the Pancreas Which is Not Associated with von Hippel-Lindau Disease"	<1%

学术搜索

找到约 181 条结果 (用时0.11秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

英语网页

☒ 包括专利☒ 包含引用☐ 创建快讯

Localization of pancreatic endocrine tumors by endoscopic ultrasonography

T Rösch, CJ Lightdale, JF Botet... - ... England Journal of ..., 1992 - Mass Medical Soc
... ampullary carcinoma by endoscopic ultrasonography: comparison with conventional sonography,
computed tomography, and angiography ... 2011) Magnetic resonance imaging versus endoscopic ...
2011) EUS is still superior to multidetector computerized tomography for detection ...

被引用次数: 641 相关文章 所有 6 个版本 引用 保存

Adrenal radiology: distinguishing benign from malignant adrenal masses.

NR Dunnick, M Korobkin... - AJR. American journal of ..., 1996 - Am Roentgen Ray Soc
... The adrenal glands may also be imaged by MR. ... Distinguishing Benign from
Malignant Causes CT, MR imaging, and adrenal scintigra- phy may each be used
to distinguish a malig- nant adrenal mass from a benign lesion. ...

被引用次数: 130 相关文章 所有 4 个版本 引用 保存

The optimal imaging of adrenal tumours: a comparison of different methods

I Ilias, A Sahdev, RH Reznek... - ... -Related Cancer, 2007 - Soc Endocrinology
... SPECT) scintigraphy with various radionuclide tracers, and positron emission tomographic (PET)
scintigraphy ... (A) Whole-body single-photon emission computed tomography with [123 I ... image
obtained with [18 F]-fluorodopamine (FDA) positron emission tomography (PET) of ...

被引用次数: 156 相关文章 所有 3 个版本 引用 保存

Clear cell "sugar" tumor of the pancreas: a novel member of the family of lesions characterized by the presence of perivascular epithelioid cells

G Zamboni, M Pea, G Martignoni... - The American journal ..., 1996 - journals.lww.com
... Workup included ultrasonography and computed tomography (CT) abdominal scan, both of which
revealed a solid ... An extensive metastatic workup, including bone scan, chest, abdominal and
pelvic CT scans was ... The case reported here shows unusual and interesting features. ...

被引用次数: 292 相关文章 所有 7 个版本 引用 保存

Imaging of neuroendocrine tumors

EP Tamm, EE Kim, CS Ng - Hematology/oncology clinics of North America, 2007 - Elsevier
... some pheochromocytomas have been reported to have similar noncontrast and washout
characteristics to adenomas ... The imaging features of extra-adrenal tumors are similar to those
of tumors of ... than CT for the detection of extra-adrenal paraganglioma, but CT has comparable ...

被引用次数: 59 相关文章 所有 6 个版本 引用 保存

Functional imaging of endocrine tumors: role of positron emission tomography

K Pasak, C Eisenhofer, DS Goldstein, Endocrine Reviews, 2004, mass.endocrine.org

endocrinology-journals.org 中的 [HTML]

endocrine.org 中的 [HTML]



ance imaging features of lipid-rich neuroendocrine tumors of the pancreas



网页 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 253,000 条结果 (用时 0.91 秒)

Google 学术 : Computed tomography and magnetic resonance imaging features of lipid-rich neuroendocrine tumors of the pancreas

Localization of **pancreatic** endocrine tumors by ... - Rösch - 被引用次数 : 642

... : distinguishing benign from malignant adrenal masses. - Dunnick - 被引用次数 : 130

... optimal **imaging** of adrenal tumours: a comparison of ... - Ilias - 被引用次数 : 155

Imaging and Pathology of Pancreatic Neoplasms: A Pictorial ...

<https://books.google.com/books?isbn=8847056780> - 翻译此页

Mirko D'Onofrio, Paola Capelli, Paolo Pederzoli - 2014 - Medical

Cancer 92:1984–1991 Fryer E, Serra S, Chetty R (2012) **Lipid-rich** ("clear cell") ...

Basturk O, Klimstra DS et al (2006) **Lipid-rich** variant of **pancreatic endocrine neoplasms**. ... **computed tomography**, and **magnetic resonance imaging features**.

Multiple Neuroendocrine Tumors of the Pancreas in von ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者 : IA Lubensky - 1998 - 被引用次数 : 197 - 相关文章

Although **pancreatic neuroendocrine tumors** (NETs) in von Hippel-Lindau (VHL) disease have ... and the clear cells additionally revealed abundant intracytoplasmic **lipid**. ... morphology are important histological **features** of VHL-associated NETs. ... during screening by **computed tomography** or **magnetic resonance imaging**.

Imaging in Cushing's syndrome - SciELO

www.scielo.br/scielo.php?pid=s0004...script=sci_arttext ▾ 翻译此页

作者 : A Sahdev - 2007 - 被引用次数 : 29 - 相关文章