

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 16282

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

Risk factors for adverse in-hospital outcomes in acute colonic diverticular hemorrhage

Nagata N *et al.* Inhospital outcomes of bleeding diverticula

Naoyoshi Nagata, Ryota Niikura, Tomonori Aoki, Shiori Moriyasu, Toshiyuki Sakurai, Takuro Shimbo, Katsunori Sekine, Hidetaka Okubo, Kazuhiro Watanabe, Chizu Yokoi, Junichi Akiyama, Mikio Yanase, Masashi Mizokami, Kazuma Fujimoto, Naomi Uemura

Naoyoshi Nagata, Ryota Niikura, Tomonori Aoki, Shiori Moriyasu, Katsunori Sekine, Hidetaka Okubo, Kazuhiro Watanabe, Toshiyuki Sakurai,

Match Overview

1	Internet 125 words crawled on 01-Jul-2015 www.wjgnet.com	2%
2	CrossCheck 92 words Nagata, Naoyoshi, Ryota Niikura, Tomonori Aoki, Takuro Shimbo, Yoshihiro Kishida, Katsunori Sekine, Shohel Tanak	2%
3	Internet 56 words crawled on 28-May-2014 www.ncbi.nlm.nih.gov	1%
4	Internet 28 words crawled on 16-Jun-2015 www.mdpi.com	<1%
5	Internet 25 words crawled on 23-Apr-2015 www.researchgate.net	<1%
6	Internet 20 words crawled on 20-Jan-2010 www.nature.com	<1%

学术搜索

找到约 1,320 条结果 (用时0.08秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

英语网页

☒ 包括专利☒ 包含引用

创建快讯

Risk factors for mortality in lower intestinal bleeding

LL Strate, JZ Ayanian, G Kotler, S Syngal - Clinical Gastroenterology and ..., 2008 - Elsevier

... Factors associated with significantly increased in-hospital mortality were advanced age, intestinal ischemia ... colorectal polyps, and anal fissures were at a decreased risk of in ... 3%) died; ongoing or recurrent bleeding was the only independent predictor of adverse outcome. ...

被引用次数: 70 相关文章 所有 10 个版本 引用 保存

nih.gov 中的 [HTML]

Early predictors of severe lower gastrointestinal bleeding and adverse outcomes: a prospective study

FS Velayos, A Williamson, KH Sousa, E Lung... - Clinical ..., 2004 - Elsevier

... Severe LGIB is a significant risk factor for global adverse outcomes. ... Clinical factors in the prediction of further haemorrhage or mortality in acute upper gastrointestinal haemorrhage. Br J Surg, 73 (1986), pp. ... Risk assessment after acute upper gastrointestinal haemorrhage. ...

被引用次数: 79 相关文章 所有 6 个版本 引用 保存

researchgate.net 中的 [PDF]

[引用] Lower GI bleeding: epidemiology and diagnosis

LL Strate - Gastroenterology Clinics of North America, 2005 - Elsevier

被引用次数: 139 相关文章 所有 8 个版本 引用 保存

upenn.edu 中的 [PDF]

Management of the adult patient with acute lower gastrointestinal bleeding

G Zuccaro - The American journal of gastroenterology, 1998 - Elsevier

... The treating physician must consider several factors when ... Among the characteristics of the group at high risk for adverse out- comes were: a greater prevalence of comorbid illness (renal, hepatic, pulmonary, hematological, neurological, or cardi- ac), a lower serum albumin ...

被引用次数: 239 相关文章 所有 16 个版本 引用 保存

acggi.net 中的 [HTML]

Incidence of perforated diverticulitis and risk factors for death in a UK population

CR Morris, IM Harvey, WSL Stebbings... - British Journal of ..., 2008 - Wiley Online Library

... Of the in-hospital factors, only an ASA grade above II was independently associated with death. ... diverticulitis may help in identifying a population at high risk, and risk factor modification or ... Anti-inflammatory drugs, analgesics and the risk of perforated colonic diverticular disease. ...

被引用次数: 82 相关文章 所有 5 个版本 引用 保存

nature.com 中的 [HTML]

Diagnosis and management of lower gastrointestinal bleeding

J Barnert, H Messmann - Nature Reviews Gastroenterology and ..., 2009 - nature.com

... Using these criteria, patients were stratified into three risk groups: low (no risk factor), moderate (1-3 risk factors), and high (>3 risk factors). ... There are several risk scores to stratify patients with acute LGIB to appropriate levels of care and intervention

[网页](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 133,000 条结果 (用时 0.57 秒)

Google 学术 : Risk factors for adverse in-hospital outcomes in acute colonic diverticular hemorrhage

Risk factors for mortality in lower intestinal bleeding - Strate - 被引用次数 : 70

... bleeding and adverse outcomes: a prospective study - Velayos - 被引用次数 : 79

Lower GI bleeding: epidemiology and diagnosis - Strate - 被引用次数 : 139

Gastrointestinal bleeding in the elderly - Nature

www.nature.com > Home > Archive > Vol 5 No 2 > Review ▾ [翻译此页](#)

作者 : PS Yachimski - 2008 - 被引用次数 : 40 - [相关文章](#)

Gastrointestinal bleeding can be classified as acute (presenting as hematemesis, ... not been shown to correlate with adverse outcomes, elderly patients who are taking ... Most people who have colonic diverticulosis are asymptomatic and have an ... Additional risk factors include small-vessel vascular disease (including ...

Risk Factors for Mortality in Lower Intestinal Bleeding

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : LL Strate - 2008 - 被引用次数 : 70 - [相关文章](#)

2008年6月16日 - We sought to identify characteristics associated with in-hospital mortality in a ... limited the ability to study rare outcomes of LIB, including in-hospital mortality. ... adverse outcome measures including death and bleeding severity. ... included: diverticulosis or diverticulitis of the colon with hemorrhage (ICD-9 ...

Management of the Adult Patient with Acute Lower ...

gi.org > Guideline ▾ [翻译此页](#)

作者 : G Zuccaro Jr - 1998 - 被引用次数 : 239 - [相关文章](#)

The source of acute gastrointestinal bleeding is not always apparent from initial history