



Are faecal markers good indicators of mucosal healing in inflammatory t



网页

图片

购物

新闻

视频

更多

搜索工具

找到约 120,000 条结果 (用时 0.64 秒)

Google 学术: Are faecal markers good indicators of mucosal healing in inflammatory bowel disease?

... of mucosal healing in patients with inflammatory bowel ... - Røseth - 被引用次数: 182

Mucosal healing predicts sustained clinical remission ... - Baert - 被引用次数: 359

... as a biological marker in inflammatory bowel disease - Gisbert - 被引用次数: 147

Looking beyond symptom relief: evolution of mucosal ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) - 翻译此页

作者: M Iacucci - 2011 - 被引用次数: 12 - 相关文章

We review the natural history of inflammatory bowel disease, the markers of disease control Faecal calprotectin as a marker showed high diagnostic accuracy in Crohn's Currently the best indicator of this appears to be mucosal healing.

The role and utility of faecal markers in inflammatory bowel ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者: FS Lehmann - 2015 - 被引用次数: 1 - 相关文章

Faecal markers such as calprotectin or lactoferrin have emerged as new In modern IBD treatment, mucosal healing (MH) rather than clinical symptom control is all patients with endoscopic and histological signs of pouchitis had elevated ... a good correlation with the traditional laboratory ELISA tests [Wassell et al.

The Role and Utility of Fecal Markers in IBD - Medscape

www.medscape.com/viewarticle/837153 - 翻译此页

How could fecal markers assist clinicians in better understanding the pathophysiology of IBD and how to best treat the disease?



19441-Review

BY GUDULA JAM BOON

Quotes Excluded
Bibliography Excluded2%
SIMILAR

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 19441

Manuscript Type: SYSTEMATIC REVIEWS

Are faecal markers good indicators of mucosal healing in inflammatory bowel disease?

Gudula JAM Boon, Andrew S Day, Chris J Mulder, Richard B Gearry

Abstract

The inflammatory bowel diseases (IBD), including Crohn's disease and ulcerative colitis, are chronic diseases characterised by inflammation of the gastrointestinal

Match Overview

1	CrossCheck 42 words Scafoli, Eleonora, Michele Scagliarini, Carla Cardamone, Elisa Liverani, Giampaolo Ugolini, Davide Festi, Franco Bazz	1%
2	CrossCheck 23 words Guidi, Luisa, Manuela Marzo, Gianluca Andrisani, Carla Felice, Daniela Pugliese, Giammarco Mucci, Olga Nardone, It	<1%
3	Internet 22 words crawled on 04-Mar-2012 otago.ourarchive.ac.nz	<1%
4	CrossCheck 12 words Sipponen, Taina, and Kaija-Leena Kolho. "Fecal calprotectin in diagnosis and clinical assessment of inflammatory b...	<1%
5	CrossCheck 10 words Dhaliwal, A., Z. Zeino, C. Tomkins, M. Cheung, C. Nwokolo, S. Smith, C. Harmston, and R. P. Arasaradnam. "Utility of fa	<1%




[学术搜索](#)

找到约 9,670 条结果 (用时0.14秒)

[我的著作引用情况](#)
[文章](#)
[我的图书馆](#)
[时间不限](#)
[2015以来](#)
[2014以来](#)
[2011以来](#)
[自定义范围...](#)
[按相关性排序](#)
[按日期排序](#)
[搜索所有网页](#)
[中文网页](#)
[简体中文网页](#)
☒ [包括专利](#)
☐ [包含引用](#)

小提示： 只搜索中文(简体)结果，可在 [学术搜索设置](#) 指定搜索语言

Mucosal healing predicts sustained clinical remission in patients with early-stage Crohn's disease hhr.be 中的 [PDF]

..., Belgian Inflammatory Bowel Disease Research Group - Gastroenterology, 2010 - Elsevier

For the best product experience, we recommend you upgrade to a newer version of IE ... **Mucosal healing** was a predictive **marker** irrespective of the CDAL ... **Stool markers**, including calprotectin and lactoferrin, have been suggested as reliable noninvasive substitutes for endoscopy ...

被引用次数: 400 [相关文章](#) [所有 18 个版本](#) [引用](#) [保存](#)

Questions and answers on the role of faecal calprotectin as a biological marker in inflammatory bowel disease

JP Gisbert, AG McNicholl - Digestive and Liver Disease, 2009 - Elsevier

... Biological **marker**; Calprotectin; Crohn's disease; Fecal marker; Inflammatory bowel disease; Ulcerative colitis. ... using the, inexpensive and easy, ELISA technique [2]. Five grams of **faeces** is enough ... As it has been stated by Konikoff and Denson [3], **faecal** calprotectin should not ...

被引用次数: 161 [相关文章](#) [所有 4 个版本](#) [引用](#) [保存](#)

Fecal calprotectin, lactoferrin, and endoscopic disease activity in monitoring anti - TNF - alpha therapy for Crohn's disease researchgate.net 中的 [PDF]

T Sipponen, E Savilahti, P Kärkkäinen... - Inflammatory bowel ..., 2008 - Wiley Online Library

... The concentration of calprotectin and lactoferrin in stool declined in treatment responders significantly from the ... **Faecal** calprotectin: a **marker** of inflammation throughout the intestinal tract. ... **Fecal** lactoferrin is a sensitive and specific **marker** of disease activity in children and young ...

被引用次数: 143 [相关文章](#) [所有 7 个版本](#) [引用](#) [保存](#)

Faecal calprotectin and lactoferrin are reliable surrogate markers of endoscopic response during lab-tech.no 中的 [PDF]



[网页](#) [图片](#) [新闻](#) [视频](#) [更多 ▾](#) [搜索工具](#)

找到约 175,000 条结果 (用时 0.49 秒)

Google 学术 : Are faecal markers good indicators of mucosal healing in inflammatory bowel disease?

Mucosal healing predicts sustained clinical remission ... - Baert - 被引用次数 : 401
... as a biological marker in inflammatory bowel disease - Gisbert - 被引用次数 : 161
... monitoring anti-TNF-alpha therapy for Crohn's disease - Sipponen - 被引用次数 : 144

Looking beyond symptom relief: evolution of mucosal ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : M Iacucci - 2011 - 被引用次数 : 14 - [相关文章](#)

We review the natural history of inflammatory bowel disease, the markers of disease ...
disease activity score worldwide in clinical trials of Crohn's disease [Best et al. ... At 10
weeks, only 36% of patients with mucosal healing were in remission as ... A
combination of stool markers, CRP and activity index may increase the ...

The role and utility of faecal markers in inflammatory bowel ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : FS Lehmann - 2015 - 被引用次数 : 7 - [相关文章](#)

Faecal markers such as calprotectin or lactoferrin have emerged as new In modern
IBD treatment, mucosal healing (MH) rather than clinical symptom control is all
patients with endoscopic and histological signs of pouchitis had elevated ... a good
correlation with the traditional laboratory ELISA tests [Wassell et al.

Inflammatory bowel disease activity assessed by fecal ...

www.biomedcentral.com/1756-0500/2/221 ▾ [翻译此页](#)

作者 : A Vieira - 2009 - 被引用次数 : 63 - [相关文章](#)