



Evaluation of a multiplex PCR assay for detection of cytomegalovirus in :



网页

图片

新闻

购物

视频

更多

搜索工具

找到约 9,060 条结果 (用时 0.73 秒)

Google 学术: Evaluation of a multiplex PCR assay for detection of cytomegalovirus in stool samples from patients with ulcerative colitis

... to detect cytomegalovirus (CMV)-DNA in stool samples ... - Herfarth - 被引用次数: 22

... subspecies paratuberculosis and Crohn's disease: a ... - Feller - 被引用次数: 332

Detection of Mycobacterium avium subspecies ... - Hulten - 被引用次数: 131

(CMV)-DNA in Stool Samples of Patients with Inflammatory ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: HH Herfarth - 2010 - 被引用次数: 22 - 相关文章

2010年2月18日 - The prevalence of CMV IgG in patients with ulcerative colitis (UC) was recently ... for the qualitative assay but as a multiplex in the quantitative assay. ... The detection of systemic CMV DNA by evaluation of plasma by PCR was ...

BMC Infectious Diseases | Full text | Persistent digestive ...

www.biomedcentral.com/1471-2334/13/37 ▾ 翻译此页

作者: SL Becker - 2013 - 被引用次数: 18 - 相关文章

2013年1月24日 - Hence, multiplex PCR assays have become the test of choice with excellent ... On fresh stool samples, direct microscopy is performed by mixing a small ... In particular immunosuppressed patients are at risk for cytomegalovirus (CMV) ... Detection of CMV-DNA by molecular methods alone is not sufficient.

Evaluation of a new automated homogeneous PCR assay ...

www.jove.com/.../evaluation-new-automated-homogeneous-pc... ▾ 翻译此页

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 20331

Manuscript Type: ORIGINAL ARTICLE

Prospective study

Evaluation of a multiplex PCR assay for detection of cytomegalovirus in stool samples from patients with ulcerative colitis

Saifun Nahar, Atsushi Iraha, Akira Hokama, Ayako Uehara, Gretchen Parrott, Tetsuya Ohira, Masatoshi Kaida, Tetsu Kinjo, Takeshi Kinjo, Tetsuo Hirata, Nagisa Kinjo, Jiro Fujita

Abstract

AIM: To evaluate a multiplex PCR assay for the detection of bacterial and viral enteropathogens in stool samples from patients with ulcerative colitis (UC).

Match Overview

1	CrossCheck 75 words Hans H. Herfarth. "Evaluation of a Non-invasive Method to Detect Cytomegalovirus (CMV)-DNA in Stool Sample ..."	1%
2	Internet 57 words crawled on 09-Oct-2012 bio.alt.ua	1%
3	CrossCheck 50 words "Posters", Clinical Microbiology and Infection, 05/2011	1%
4	Internet 37 words crawled on 08-Jun-2015 www.science.gov	1%
5	CrossCheck 29 words COUPLAND, L. J., I. McELARNEY, E. MEADER, K. COWLEY, L. ALCOCK, J. NAUNTON, and J. GRAY. "Simultaneous"	<1%
6	CrossCheck 28 words Onori, Manuela, Luana Coltella, Livia Mancinelli, Marta Argentieri, Donato Menichella, Alberto Villani, Annalisa Gra	<1%
7	Internet 24 words crawled on 06-Jan-2015 sgh.org.sa	<1%
8	Internet 23 words crawled on 27-Jan-2015 doaj.org	<1%



Evaluation of a multiplex PCR assay for detection of cytomegalovirus ir



网页

新闻

图片

视频

更多

搜索工具

找到约 7,680 条结果 (用时 0.71 秒)

(CMV)-DNA in Stool Samples of Patients with Inflammatory ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: HH Herfarth - 2010 - 被引用次数: 25 - 相关文章

2010年2月18日 - **Evaluation** of a Non-invasive Method to **Detect Cytomegalovirus** ... The prevalence of CMV IgG in **patients with ulcerative colitis** (UC) was ... Qualitative and/or quantitative **stool** CMV DNA testing by **PCR** is a ... The internal control was detected separately for the qualitative **assay** but as a **multiplex** in the ...

Evaluation of a Non-invasive Method to Detect ... - Academic

65.54.113.239/.../**evaluation-of-a-non-invasive-method-to-dete**... ▾ 翻译此页

Background A severe flare of **colitis** in **patients** with IBD treated with ... chain reaction (PCR) to **detect** CMV DNA in **stool samples** of IBD **patients**. ... **Multiplex**, Quantitative, Real-Time **PCR Assay** for Cytomegalovirus and Human DNA (Citations: 7) ... Cytomegalovirus disappearance after treatment for refractory **ulcerative** ...

Findings From a Randomized Controlled Trial of Fecal ...

www.sciencedirect.com/science/article/pii/S0016508515004485 - 翻译此页

作者: NG Rossen - 2015 - 被引用次数: 15 - 相关文章

The general hypothesis on the cause of **ulcerative colitis** (UC) is that there is an ... proof-of-concept phase 2 trial, the efficacy of FMT was **evaluated** in **patients** with ... The maximum **sample** size of 80 would allow **detection** of a treatment delta of ... Fecal **samples** were collected at baseline before **bowel** lavage and 6 and 12 ...

BMC Infectious Diseases | Full text | Persistent digestive ...

www.biomedcentral.com/1471-2334/13/37 ▾ 翻译此页

作者: SL Becker - 2013 - 被引用次数: 27 - 相关文章

2013年1月24日 - We performed a literature **review** to identify and define the bacterial,

[网页](#)[新闻](#)[图片](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 7,380 条结果 (用时 0.62 秒)

(CMV)-DNA in Stool Samples of Patients with Inflammatory ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者：HH Herfarth - 2010 - 被引用次数：25 - 相关文章

2010年2月18日 - [Evaluation](#) of a Non-invasive Method to [Detect Cytomegalovirus](#) ...

The prevalence of CMV IgG in [patients with ulcerative colitis](#) (UC) was ... Qualitative and/or quantitative [stool](#) CMV DNA testing by [PCR](#) is a ... The internal control was detected separately for the qualitative [assay](#) but as a [multiplex](#) in the ...

Evaluation of a Non-invasive Method to Detect ... - Academic

65.54.113.239/.../evaluation-of-a-non-invasive-method-to-dete... ▾ [翻译此页](#)

Background A severe flare of [colitis](#) in [patients](#) with IBD treated with ... chain reaction

(PCR) to [detect CMV](#) DNA in [stool samples](#) of IBD [patients](#). ... [Multiplex](#), Quantitative,

Real-Time [PCR Assay](#) for Cytomegalovirus and Human DNA (Citations: 7) ...

Cytomegalovirus disappearance after treatment for refractory [ulcerative](#) ...

Findings From a Randomized Controlled Trial of Fecal ...

www.sciencedirect.com/science/article/pii/S0016508515004485 - [翻译此页](#)

作者：NG Rossen - 2015 - 被引用次数：15 - 相关文章

The general hypothesis on the cause of [ulcerative colitis](#) (UC) is that there is an ...

proof-of-concept phase 2 trial, the efficacy of FMT was [evaluated](#) in [patients](#) with ...

The maximum [sample](#) size of 80 would allow [detection](#) of a treatment delta of ... Fecal [samples](#) were collected at baseline before [bowel](#) lavage and 6 and 12 ...

American Journal of Gastroenterology - Temporal Bacterial ...

www.nature.com > ... > [Archive](#) > [Original Contributions](#) ▾ [翻译此页](#)

作者：S Angelberger - 2013 - 被引用次数：96 - 相关文章

2013年9月24日 - Five [patients](#) with moderately to severely active [ulcerative colitis](#)

The ethics committee of the Medical University of Vienna approved the retrospective