


学术搜索

找到约 25 条结果 (用时0.07秒)

 我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...


按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用 创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

Stool color card screening for biliary atresia

JJ Tseng, MS Lai, MC Lin, YC Fu - Pediatrics, 2011 - Am Acad Pediatrics

... The median ages of patients receiving the Kasai operation advanced a little after the stool color card ... if we only include the patients who had a Kasai operation, there was a reduction in the ... 26 Reducing the age at which patients receive the Kasai operation is still the leading goal ...

被引用次数: 21 相关文章 所有 5 个版本 引用 保存

Early diagnosis of neonatal cholestatic jaundice Test at 2 weeks

El Benchimol, CM Walsh, SC Ling - Canadian Family Physician, 2009 - cfp.ca

... atresia is directly affected by the speed with which health care providers arrive at the diagnosis, ... and 99.9%, respectively.42 The average age at the time of portoenterostomy was reduced to 53 ... SerinetMOWildhaberBEBrouéPLachauxASarlesJJacqueminElmpact of age at Kasai ...

被引用次数: 31 相关文章 所有 10 个版本 引用 保存

Peculiar features and tailored management of adult cancers occurring in pediatric age

A Ferrari, M Casanova, M Massimino, I Sultan - 2010 - informahealthcare.com

... View larger version(47K), Figure 2. Frequency of colorectal cancer across ages from the ... 2 trial investigated chemotherapy intensification in HCC, adding carboplatin to PLADO and reducing the interval ... found to significantly delay the onset of acute toxicity and reduce its severity ...

被引用次数: 11 相关文章 所有 4 个版本 引用 保存

Surgeon volume and operative mortality in the United States

JD Birkmeyer, TA Stukel, AE Siewers... - ... England Journal of ..., 2003 - Mass Medical Soc

... used Medicare data, our study was restricted to patients 65 years of age or older. ... However, to reduce any potential bias against low-volume surgeons, we excluded physicians who ... diagnoses and the association between distance to an Alzheimer clinic and Alzheimer diagnosis. ...

aappublications.org 中的 [HTML]

cfp.ca 中的 [PDF]

uro-basel.ch 中的 [PDF]

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 21555

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Reduction of the ages at diagnosis and operation of biliary atresia in Taiwan: A
15-year population-based cohort study

Lin JS *et al.* Reducing ages at diagnosis and operation of BA

Jen-Shyang Lin, Solomon Chih-Cheng Chen, Chin-Li Lu, Hung-Chang Lee,
Chun-Yan Yeung, Wai-Tao Chan

Abstract

AIM: To describe the ages at diagnosis and operation of biliary atresia (BA) and its
incidence over a 15-year period in Taiwan.

Match Overview

1	CrossCheck 38 words Lin, C.C.. "Effects of surgeon and hospital volume on 5 ... ear survival rates following oral cancer resections: The e	1%
2	Internet 36 words crawled on 30-Mar-2012 www.basca.ch	1%
3	CrossCheck 34 words Mao-Meng Tiao. "Epidemiological features of biliary at ... sia in Taiwan, a national study 1996?2003", Journal of G	1%
4	Internet 30 words crawled on 24-Nov-2013 www.recentmedicalfindings.com	1%
5	Internet 18 words crawled on 24-Aug-2015 www.biomedcentral.com	<1%
6	CrossCheck 15 words Wu, Jia-Feng, Pei-Chi Kao, Huey-Ling Chan, Hong-Shia e Lai, Hong-Yuan Hsu, Mei-Hwei Chang, and Yen-Hsua	<1%
7	Internet 15 words crawled on 11-May-2014 www.nimonline.nl	<1%

学术搜索

找到约 31 条结果 (用时0.09秒)

 我的著作引用情况

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...


按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Stool color card screening for biliary atresia

JJ Tseng, MS Lai, MC Lin, YC Fu - Pediatrics, 2011 - Am Acad Pediatrics

... The median ages of patients receiving the Kasai operation advanced a little after the stool color card ... if we only include the patients who had a Kasai operation, there was a reduction in the ... 26
Reducing the age at which patients receive the Kasai operation is still the leading goal ...

被引用次数: 24 相关文章 所有 5 个版本 引用 保存

aappublications.org 中的 [HTML]

Early diagnosis of neonatal cholestatic jaundice Test at 2 weeks

El Benchimol, CM Walsh, SC Ling - Canadian Family Physician, 2009 - cfp.ca

... atresia is directly affected by the speed with which health care providers arrive at the diagnosis. ... and 99.9%, respectively.42 The average age at the time of portoenterostomy was reduced to 53 ... SerinetMOWildhaberBEBrouéPLachauxASarlesJJacqueminElmpact of age at Kasai ...

被引用次数: 33 相关文章 所有 10 个版本 引用 保存

cfp.ca 中的 [PDF]

Peculiar features and tailored management of adult cancers occurring in pediatric age

A Ferrari, M Casanova, M Massimino, I Sultan - 2010 - informahealthcare.com

... View larger version(47K), Figure 2. Frequency of colorectal cancer across ages from the ... 2 trial investigated chemotherapy intensification in HCC, adding carboplatin to PLADO and reducing the interval ... found to significantly delay the onset of acute toxicity and reduce its severity ...

被引用次数: 11 相关文章 所有 4 个版本 引用 保存

alsendibad.net 中的 [PDF]

[引用] EXTRA HEPATIC BILIARY ATRESIA

RL Meyers, EG Pearson - Surgical Care of Major Newborn ..., 2012 - World Scientific

相关文章 所有 3 个版本 引用 保存

The Causal Effect of Early Surgical Intervention on Biliary Atresia Patients Clinical Outcome Using Economic and Education Models

E Hoshino - 2015 - academiccommons.columbia.edu

... Observational studies suggested a number of possible benefits of treatment with ursodeoxycholic acid, ranging from enhanced weight gain to reduced ... nationwide stool color card was associated with a decrease in the median time to diagnosis ... early age. Tien et al.

columbia.edu 中的 [PDF]

[网页](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 6,400 条结果 (用时 0.63 秒)

Stool Color Card Screening for Biliary Atresia | Articles ...

pediatrics.aappublications.org/content/128/5/e1209 - 翻译此页

作者：JJ Tseng - 2011 - 被引用次数：24 - 相关文章

In this **study**, we evaluated the effectiveness of stool color card screening by using ...

Only early **diagnosis** of **biliary atresia** can lead to early **operation**. ... Although the NHI of **Taiwan** was initiated in 1995, the database for that **year** was incomplete.

Reducing the **age** at which patients receive the Kasai **operation** is still the ...

Stool Color Card Screening for Biliary Atresia ... - Pediatrics

pediatrics.aappublications.org/content/128/5/e1209.full.pdf ▾ 翻译此页

作者：JJ Tseng - 2011 - 被引用次数：24 - 相关文章

charts of all inpatients who were **diagnosed** with **biliary atresia** from 1996 to 2008

were collected from **Taiwan's** National Health Insurance ... **biliary atresia**, stool color

card screening, Kasai **operation** ... This was a retrospective **cohort study**. Although

than 60 days of **age**.12,15 In a 2-**year** ob- a **population-based** study.

Screening and Outcomes in Biliary Atresia: Summary of a ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者：RJ Sokol - 2007 - 被引用次数：131 - 相关文章

Earlier **diagnosis** (<30-45 days of life) is associated with improved outcomes

following ... **Biliary atresia** (BA) is the most common pediatric cause of cirrhosis,

The resolution of jaundice after the HPE **operation** was 39% in the second **cohort** of patients. A full-scale **population-based** screening observational **study** using ...

Early diagnosis of neonatal cholestatic jaundice

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者：El Benchimol - 2009 - 被引用次数：33 - 相关文章

Approximately 60% to 70% of patients with **biliary atresia** will develop cirrhosis and ...