



Spontaneous aortic pseudoaneurysm rupture into the sigmoid colon via the



Spontaneous aortic pseudoaneurysm rupture into the sigmoid colon via the mesocolon as a vascular complication of Behçet's disease

学术搜索

获得 10 条结果 (用时0.10秒)

我的著作引用情况



文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☐ 包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Beyond decreased bowel enhancement: acute abnormalities of the mesenteric and portal vasculature

CK Sandstrom, CR Ingraham, EJ Monroe... - Abdominal ..., 2015 - Springer

... Isolated visceral arterial dissection, most often of the SMA, is a rare condition not associated with aortic dissection. ... SAM may be under recognized and may underlie an apparently spontaneous SMA dissection [29]. Fig. 12 ... Reflux of contrast into abdominal aorta.

引用 保存

[图书] Emergency Radiology: Imaging of Acute Pathologies

A Singh - 2013 - books.google.com

... Intramural hematoma is characterized by lack of direct communication of the arterial lumen with the ... (a) Contrast-enhanced CT demonstrates asymmetric aortic wall thickening, consistent with intramural hematoma (arrowheads) in the descending thoracic aorta. ...

相关文章 所有 6 个版本 引用 保存 更多

Diverticulum, Oesophagus

O Ekberg - Encyclopedia of Diagnostic Imaging, 2008 - Springer

... Injury of the endplate cartilage causes an aseptic reaction of the subchondral bone, with increase of ... become serious alternatives to surgical procedures and other treatment options. ... severity into four grades, from normal to severe osteoarthritis (6). In addition and among others ...

相关文章 引用 保存

Direct Imaging

CT Chan, SS Gambhir - Encyclopedia of Diagnostic Imaging, 2008 - Springer

... Injury of the endplate cartilage causes an aseptic reaction of the subchondral bone, with increase of ... become serious alternatives to surgical procedures and other treatment options. ... severity into

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 21603

Manuscript Type: CASE REPORT

Spontaneous aortic pseudoaneurysm rupture into the sigmoid colon in Behçet's disease patient

Lee SL *et al.* Behçet's disease vascular complication

Su Lim Lee, Young Mi Ku, Yoodong Won

Abstract

Behçet's disease (BD) is a multisystem autoimmune disorder characterized by recurrent orogenital ulcers, uveitis, and skin lesions. The vascular manifestations include thrombophlebitis, stenosis, occlusion, and pseudoaneurysm. BD infrequently precipitates aortic pseudoaneurysm

Match Overview

1	CrossCheck 19 words Murat Ugurlucan. "Complication of Behcet's Disease: Spontaneous Aortic Pseudoaneurysm", <i>Journal of Cardiac Surg</i>	1%
2	CrossCheck 18 words Hwang, Yunsup, Su Lim Lee, Eun Deok Chang, Chang Hyek Ahn, and Young Mi Ku. "CT and MR Imaging Findings o	1%
3	CrossCheck 15 words Jang, Suyoung, Myung Ho Jeong, Hyung Ki Jung, Kyung Jin Lee, Kye Hun Kim, Young Joon Hong, Ju Han Kim, and Yo	1%

小提示: 只搜索中文(简体)结果, 可在 [学术搜索设置](#) 指定搜索语言

Behçet's disease

T Sakane, M Takeno, N Suzuki... - New England Journal of ... , 1999 - Mass Medical Soc

... The lesions are painful and usually resolve **spontaneously**, leaving a deeply pigmented area, but they sometimes ulcerate. ... The **rupture** of such aneurysms may be fatal. ... Pulmonary **arterial** aneurysms in **Behçet's** syndrome: a report of 24 cases. ...

被引用次数: 1439 [相关文章](#) [所有 8 个版本](#) [引用](#) [保存](#)

[HTML] Aortitis: imaging spectrum of the infectious and inflammatory conditions of the aorta

CS Restrepo, D Ocazonez, R Suri, D Vargas - Radiographics, 2011 - pubs.rsna.org

... coronary ostia, this may lead to myocardial ischemia (2). Multiple **aortic** aneurysms as well as **spontaneous rupture** in **patients** ... images show a saccular **pseudoaneurysm** (arrow) that arises from the lateral wall of the **aorta**, resulting in extrinsic compression of the **aortic** lumen. ...

被引用次数: 53 [相关文章](#) [所有 6 个版本](#) [引用](#) [保存](#)

Abdominal aortic aneurysm

N Sakalihasan, R Limet, OD Defawe - The Lancet, 2005 - Elsevier

... and colleagues 5 proposed the definition of an abdominal **aortic** aneurysm as an **aorta** with an ... This dilatation affects the three layers of the vascular tunic; otherwise, the dilatation is called a **pseudoaneurysm**. ... 25 Thus, most abdominal **aortic** aneurysms are called non-specific. ...

被引用次数: 603 [相关文章](#) [所有 12 个版本](#) [引用](#) [保存](#)

Visceral and Renal Artery Aneurysms: A Pictorial Essay on Endovascular Therapy 1

JL Noshier, J Chung, LS Brevetti, AM Graham... - Radiographics, 2006 - pubs.rsna.org

... guidance, a 20- or 22-gauge needle is introduced **into** the **pseudoaneurysm** and thrombin ... of all VAAs (1). CAAs are often degenerative, reportedly being associated with **aortic** aneurysms in ... aneurysm embolization of the peripheral vascular bed or narrowing of the **arterial** lumen ...

被引用次数: 123 [相关文章](#) [所有 4 个版本](#) [引用](#) [保存](#)

Endoluminal stent-grafts for infrarenal abdominal aortic aneurysms

rsna.org 中的 [\[HTML\]](#)

ulg.ac.be 中的 [\[PDF\]](#)

rsna.org 中的 [\[HTML\]](#)

[网页](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 28,700 条结果 (用时 0.62 秒)

Related Articles - PubMed Central Canada

pubmedcentralcanada.ca/pmcc/solr/mlt;...acc... ▾ [翻译此页](#)

Results 1 - 25 - **Rupture** of abdominal **aortic aneurysm** into sigmoid colon: A case ...

Primary aorto-colic fistula is a **spontaneous rupture** of abdominal **aortic aneurysm** into the in **patients** with the multisystemic inflammatory disorder **Behçet's disease**. An aorto-atrial fistula secondary to **Behçet's disease** without a sinus of ...

Cases reported • Colonic Diseases - LookForDiagnosis.com

lookfordiagnosis.com/.../cases.php?...Colonic+Diseases...aneury... ▾ [翻译此页](#)

Lower gastrointestinal bleeding in the **patient** who has an **aortic aneurysm** ... Two cases of **spontaneous rupture** of colic artery aneurysms are presented, ... hepatic artery **aneurysm**; right hepatic artery **aneurysm** with erosion **into** the ... A 16-year-old boy had incomplete **Behcet's disease** with common carotid artery **aneurysm** ...

RadioGraphics: Aortitis: Imaging Spectrum of the Infectious ...

pubs.rsna.org/doi/full/10.1148/rg.312105069 ▾ [翻译此页](#)

作者 : CS Restrepo - 2011 - [被引用次数 : 53](#) - [相关文章](#)

Takayasu arteritis is commonly divided **into** two clinical stages, Multiple **aortic** aneurysms as well as **spontaneous rupture** in **patients** with ... **Aortic pseudoaneurysm** secondary to **Behçet disease** in a 51-year-old woman. ... **aortic-sigmoid colon** fistula with bleeding, and secondary bacterial infection (eg, with Salmonella),

Aortocolic fistula, a lethal cause of lower gastrointestinal ...

www.ncbi.nlm.nih.gov/pubmed/1086196 ▾ [翻译此页](#)

作者 : SE Wilson - 1976 - [被引用次数 : 14](#) - [相关文章](#)

Dis Colon Rectum. ... Aortocolic fistula occurs with **spontaneous rupture** of **aortic** and iliac aneurysms **into the sigmoid colon**, or due to involvement of ... Lower gastrointestinal bleeding in the **patient** who has an **aortic aneurysm** and ... **Aortic**