



Predictive Value of Fatty liver index for non-alcoholic fatty Liver disease: A pop



学术搜索

找到约 9,730 条结果 (用时0.10秒)

我的著作引用情况



文章

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

我的图书馆

[HTML] The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population

G Bedogni, S Bellentani, L Miglioli... - BMC ..., 2006 - biomedcentral.com

... from a practical viewpoint, there is no need to measure skinfolds for predicting FL ... if it may be noticed that the selection of glucose as potential predictor increased after ... FE, Borsboom GJ, Eijkemans MJ, Vergouwe Y, Habbema JD: Internal validation of predictive models: efficiency ...

被引用次数: 324 相关文章 所有 15 个版本 引用 保存 更多

biomedcentral.com 中的 [HTML]

时间不限

2015以来

2014以来

2011以来

自定义范围...

[PDF] Prevalence of non-alcoholic fatty liver disease: population based study

D Amarapurkar, P Kamani, N Patel, P Gupte... - Ann ..., 2007 - medigraphic.com

... Variables Values p value Significance ... Ultrasonography can not differentiate NAFLD and NASH. 11 Positive predictive value of ultrasound is only 34%. 15 We could ... study on diabetics we have demonstrated more than 90% sensitivity of ultrasound in predicting NAFLD when ...

被引用次数: 145 相关文章 所有 12 个版本 引用 保存

medigraphic.com 中的 [PDF]

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

The epidemiology of nonalcoholic fatty liver disease in adults

JM Clark - Journal of clinical gastroenterology, 2006 - journals.lww.com

... 14 However, in defining the epidemiology of NAFLD, it is worth noting that the prevalence ... Imaging data for hepatic fat and biopsies of large population-based samples are not available ... of middle-aged Japanese workers found a prevalence of ultrasound detected fatty liver of 22 ...

被引用次数: 660 相关文章 所有 3 个版本 引用 保存

☒ 包括专利

☐ 包含引用

[HTML] Nonalcoholic fatty liver disease: from steatosis to cirrhosis

GC Farrell, CZ Larter - Hepatology, 2006 - Wiley Online Library

wiley.com 中的 [HTML]



22549-Review

BY NIMA MOTAMED

Quotes Excluded
Bibliography Excluded10%
SIMILAR

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 22549

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Fatty liver index value *vs* waist circumference for predicting of non-alcoholic fatty liver disease

Nima Motamed, Masoudreza Sohrabi, Hossein Ajdarkosh, Gholamreza Hemmasi, Mansooreh Maadi, Reza Pirzad, Khadijeh Abedi, Sivil Aghapour, Farhad Zamani

Abstract

AIM: To determine the discriminatory performance of fatty liver index (FLI) for nonalcoholic fatty liver disease (NAFLD).

Match Overview

1	Internet 160 words crawled on 10-Nov-2015 bib.irb.hr	4%
2	Internet 133 words crawled on 18-Oct-2015 onlinelibrary.wiley.com	3%
3	CrossCheck 30 words Wing-Kin Syn. "Nonalcoholic fatty liver disease in a dis ... ct general hospital: clinical presentation and risk factors"	1%
4	CrossCheck 28 words Giuseppe Labruna. "High Leptin/Adiponectin Ratio and Serum Triglycerides Are Associated With an "At-Risk" F ...	1%
5	Internet 16 words crawled on 08-May-2015 www.ncbi.nlm.nih.gov	<1%
6	CrossCheck 14 words Jeong, Dong Wook, Hye Won Lee, Young Hye Cho, Don g Won Yi, Sang Yeoup Lee, Seok Man Son, and Yang Ho	<1%



Fatty liver index value vs waist circumference for predicting of non-alcoholic fat ▾



学术搜索

找到约 5,710 条结果 (用时0.13秒)

我的著作引用情况



文章

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

Nonalcoholic fatty liver disease a feature of the metabolic syndrome

[G Marchesini](#), [M Brizi](#), [G Bianchi](#), [S Tomassetti](#)... - Diabetes, 2001 - Am Diabetes Assoc

... Only the small difference in glucose disposal between NAFLD patients and subjects with pure fatty livers might be ... to reduce the obesity-related insulin resistance, which might be partly responsible for liver fat deposition (7 ... Wanless IR, Lentz JS: Fatty liver hepatitis (steatohepatitis ...

被引用次数: 2041 相关文章 所有 6 个版本 引用 保存

diabetesjournals.org 中的 [HTML]

Prediction of non-alcoholic fatty liver disease and liver fat using metabolic and genetic factors

[A Kotronen](#), [M Peltonen](#), [A Hakkarainen](#)... - Gastroenterology, 2009 - Elsevier

... protons of water and protons of methylene (CH2) n-2 groups in the fatty acid chains ... covariance was used to adjust liver fat for age, gender, and body mass index (BMI). ... fS low-density lipoprotein (LDL) cholesterol concentrations were not associated with increased liver fat content ...

被引用次数: 212 相关文章 所有 13 个版本 引用 保存

ejso.com 中的 [HTML]

[HTML] The Fatty Liver Index: a simple and accurate predictor of hepatic steatosis in the general population

[G Bedogni](#), [S Bellentani](#), [L Miglioli](#)... - BMC ..., 2006 - biomedcentral.com

... Durnin JV, Womersley J: Total body fat, calculated from body density, and its ... markers (FibroTest-FibroSURE) for the prediction of liver fibrosis in patients with non-alcoholic fatty liver disease. ... J: The diagnostic value of biomarkers (SteatoTest) for the prediction of liver steatosis. ...

被引用次数: 341 相关文章 所有 15 个版本 引用 保存 更多

biomedcentral.com 中的 [HTML]

Nonalcoholic fatty liver disease: predictors of nonalcoholic steatohepatitis and liver fibrosis in the severely obese

[网页](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 156,000 条结果 (用时 0.61 秒)

Non-invasive diagnosis of non-alcoholic fatty liver disease ...

www.sciencedirect.com/science/article/pii/S0168827812008902 - 翻译此页

作者：MV Machado - 2013 - 被引用次数：88 - 相关文章

Non-alcoholic fatty liver disease (NAFLD) affects one in every three subjects in the ...
predictive value; PPV, positive **predictive value**; FLI, **fatty liver index**; US, through
a formula incorporating: body mass index (BMI), **waist circumference**, CAP
performed better than Steatotest, FLI or Hepatic Steatosis Index [51].

ZJU index: a novel model for predicting nonalcoholic fatty ...

www.nature.com > Scientific Reports - 翻译此页

2015年11月16日 - **Non-alcoholic fatty liver disease** (NAFLD) is an important health
issue worldwide. ... At a **value** of <32.0, the ZJU **index** could rule out NAFLD with a
sensitivity of ... Standing height, body weight and **waist circumference** were ...

Validation of the Fatty Liver Index for Nonalcoholic Fatty ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者：X Huang - 2015

2015年10月9日 - **Nonalcoholic fatty liver disease** (NAFLD), defined as hepatic ... The
fatty liver index (FLI) is an algorithm based on **waist circumference**, body mass
fatty liver disease, NPV = negative **predictive value**, OR = odds ratio, PBG ...

Diagnosis and Evaluation of Nonalcoholic Fatty Liver Disease

www.hindawi.com/journals/jdr/2012/145754/ - 翻译此页

作者：M Obika - 2011 - 被引用次数：73 - 相关文章

2011年8月31日 - **Nonalcoholic fatty liver disease** (NAFLD) is the most common cause
of Using a cutoff serum GGT **value** of 96.5, GGT **predicted** advanced fibrosis