

Biomarkers for detection of alcohol consumption in liver transplantation

全部 图片 视频 新闻 购物 地图 图书

找到约 396,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Biomarkers for detection of alcohol consumption in liver transplantation](#)

[Alcoholic liver disease](#) - O'Shea - 被引用次数: 729

[... testing detects alcohol consumption in alcoholic liver ...](#) - Erim - 被引用次数: 61

[... of liver fibrosis in patients with chronic alcoholic liver ...](#) - Naveau - 被引用次数: 145

[\[PDF\] Standard PDF - Wiley Online Library](#)

[onlinelibrary.wiley.com/doi/10.1002/lt.23596/pdf](#)

The accurate assessment of drinking by patients with alcoholic liver disease is important both before and ... tional alcohol biomarkers are liver function tests [eg, ...

[Full Article - Wiley Online Library](#)

[onlinelibrary.wiley.com/doi/10.1002/lt.23596/full](#)

17 Mar 2013 ... At least heavy drinking after liver transplantation has been linked to substantially ... Traditional alcohol biomarkers are liver function tests [eg, ...

[Hair Ethyl Glucuronide is Highly Sensitive and Specific for Detecting ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC3523385/](#)

25 Sep 2012 ... EtG in urine has been shown to identify recent drinking in liver disease patients ... as has been suggested in organ transplant patients (Haller et al., 2010). ... Use of other biomarkers to detect under-reporting of alcohol use.

[\[PDF\] Biomolecules and Biomarkers Used in Diagnosis of Alcohol ...](#)

[www.mdpi.com/2218-273X/5/3/1339/pdf](#)

29 Jun 2015 ... the successful treatment of alcohol misuse in transplantation of patients ... alcohol consumption and the incidence of hepatic injury was noted.

[Determination of Ethyl Glucuronide in Hair for Detection of Alcohol](#)

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 23877

Manuscript Type: REVIEW

Biomarkers for detection of alcohol consumption in liver transplantation

Katharina Stauer, Michel Yegles

Abstract

Alcoholic liver disease is an established, yet controversial indication for liver transplantation. Although an abstinence period of up to 6 months prior to transplantation is mandatory, alcohol relapse after transplantation is a common event. In case of recurrence of heavy drinking, graft survival is significantly impaired. Guidelines on detection and surveillance of alcohol consumption in this patient cohort are lacking. This review summarizes the challenge of patient selection, as well as the current knowledge on established and novel alcohol biomarkers with

Match Overview

1

Internet 177 words
crawled on 26-Apr-2015
emjreviews.com

2%

2

CrossCheck 67 words
Katharina Stauer, "Urinary ethyl glucuronide as a novel screening tool in patients pre- and post-liver transplantation ...

1%

Biomarkers for detection of alcohol consumption in liver transplantation

全部 图片 视频 新闻 购物 地图 图书

找到约 186,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Biomarkers for detection of alcohol consumption in liver transplantation](#)

[Alcoholic liver disease](#) - O'Shea - 被引用次数: 756

[... testing detects alcohol consumption in alcoholic liver ...](#) - Erim - 被引用次数: 62

[... of liver fibrosis in patients with chronic alcoholic liver ...](#) - Naveau - 被引用次数: 146

[\[PDF\] Standard PDF \(132.8 KB\) - Wiley Online Library](#)

onlinelibrary.wiley.com/doi/10.1002/lt.23596/pdf

The accurate assessment of drinking by patients with alcoholic liver disease is important both before and ... tional alcohol biomarkers are liver function tests [eg, ...

[Assessing the drinking status of liver transplant patients with ...](#)

onlinelibrary.wiley.com/doi/10.1002/lt.23596/full

17 Mar 2013 ... At least heavy drinking after liver transplantation has been linked to substantially ... Traditional alcohol biomarkers are liver function tests [eg, ...

[Biomolecules and Biomarkers Used in Diagnosis of Alcohol ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4598755/

29 Jun 2015 ... Background: The quantitative, measurable detection of drinking is ... during evaluation for liver transplantation, in legal cases, or during alcohol ...

[Hair Ethyl Glucuronide is Highly Sensitive and Specific for Detecting ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC3523385/

25 Sep 2012 ... EtG in urine has been shown to identify recent drinking in liver disease patients ... as has been suggested in organ transplant patients (Haller et al., 2010). ... Use of other biomarkers to detect under-reporting of alcohol use.

[Assessment of alcohol consumption in liver transplant candidates](#)

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 401,000 条结果 (用时 0.54 秒)

Google 学术 : Biomarkers for detection of alcohol consumption in liver transplantation

... testing detects alcohol consumption in alcoholic liver ... - Erim - 被引用次数 : 63

Alcoholic liver disease - O'Shea - 被引用次数 : 793

... of liver fibrosis in patients with chronic alcoholic liver ... - Naveau - 被引用次数 : 152

Assessment of alcohol consumption in liver transplant ...

www.ncbi.nlm.nih.gov/pubmed/24692331 ▾ 翻译此页

作者 : S Piano - 2014 - 被引用次数 : 10 - 相关文章

2014年5月26日 - The detection of alcohol consumption in liver transplant candidates (LTCs) ... and other indirect markers of alcohol consumption were evaluated ...

Biomolecules and Biomarkers Used in Diagnosis of Alcohol ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - 翻译此页

作者 : RM Nanau - 2015 - 被引用次数 : 2 - 相关文章

2015年6月29日 - Background: The quantitative, measurable detection of drinking is ... during evaluation for liver transplantation, in legal cases, or during alcohol ...

Novel Objective Biomarkers of Alcohol Use: Potential ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者 : RK Kalapatapu - 2009 - 被引用次数 : 13 - 相关文章

2009年1月1日 - Keywords: Alcohol biomarker, dual diagnosis, alcohol abuse, alcohol ... detoxification and abstinence clinics, liver transplant programs, and ...

Standard PDF (132.8 KB) - Wiley Online Library

onlinelibrary.wiley.com/doi/10.1002/lt.23596/pdf ▾ 翻译此页