

学术搜索

找到约 2,840 条结果 (用时0.06秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Tumor microsatellite-instability status as a predictor of benefit from fluorouracil-based **adjuvant chemotherapy for colon cancer** nih.gov 中的 [HTML]

CM Ribic, DJ Sargent, MJ Moore... - *New England ...*, 2003 - Mass Medical Soc
... that occur during the progression from adenoma to carcinoma to **metastatic** disease.13,14 ...
Molecular predictors of survival **after adjuvant chemotherapy for colon cancer**. ... Determination of microsatellite instability, p53 and K-RAS mutations in **hepatic metastases** from patients with ...
被引用次数: 1433 相关文章 所有 8 个版本 引用 保存

[HTML] Cryosurgical ablation and radiofrequency ablation for unresectable **hepatic malignant neoplasms: a proposed algorithm** jamanetwork.com 中的 [HTML]

AJ Bilchik, TF Wood, D Allegra... - ... of *Surgery*, 2000 - archotol.jamanetwork.com
Research from JAMA Surgery — Cryosurgical Ablation and Radiofrequency Ablation for Unresectable Hepatic Malignant Neoplasms — A Proposed Algorithm.
被引用次数: 301 相关文章 所有 10 个版本 引用 保存

Small **liver colorectal metastases treated with percutaneous radiofrequency ablation: local response rate and long-term survival with up to 10-year follow-up** rsna.org 中的 [HTML]

L Solbiati, M Ahmed, L Cova, T Ierace, M Brioschi... - *Radiology*, 2012 - pubs.rsna.org
... Of these, 29 **patients** were **lost** to follow-up, largely due to logistic and ... the time of primary tumor resection, 20 **patients** (20.2%) underwent resection of synchronous **liver metastases** (one to ... that were possible at the time of **treatment**, and they included **hepatic** ultrasonography (US ...
被引用次数: 57 相关文章 所有 4 个版本 引用 保存

Improved survival with ipilimumab in **patients with metastatic melanoma** nih.gov 中的 [HTML]

FS Hodi, SJ O'Day, DF McDermott... - *New England ...*, 2010 - Mass Medical Soc
... or office work)24; positive status for HLA-A*0201; normal hematologic, **hepatic**, and renal ... survival was independent of age, sex, baseline serum lactate dehydrogenase levels, **metastasis** stage of ... 70% of the **patients** had M1c disease (presence of visceral **metastases**) and more ...
被引用次数: 4791 相关文章 所有 25 个版本 引用 保存

The management of **colorectal liver metastases**: expanding the role of **hepatic** resection in the age of multimodal **therapy**

A Chiappa, M Makawuchi, NU Lyvic

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 23984

Manuscript Type: Review

Missing metastases as a model to challenge current therapeutic algorithms in colorectal liver metastases

Lucidi V *et al.* Adjuvant surgery in patients with missing colorectal liver metastases after chemotherapy

Valerio Lucidi, Alain Hendlisz, Jean Luc Van Laethem, Vincent Donckier

Valerio Lucidi, Department of Abdominal Surgery, Hôpital Erasme, Centre de Chirurgie Hépato-Biliaire de l'ULB, Université Libre de Bruxelles, 1070 Brussels, Belgium

Alain Hendlisz, Department of Medical Oncology, Institut Jules Bordet, Université Libre de Bruxelles, 1000 Brussels, Belgium

Match Overview

1 **Internet** 175 words
crawled on 25-Feb-2016
www.wjgnet.com 5%

学术搜索

找到约 10,800 条结果 (用时0.11秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

包括专利

包含引用

创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Rescue surgery for unresectable colorectal liver metastases downstaged by chemotherapy: a model to predict long-term survival researchgate.net 中的 [PDF]

R Adam, V Delvart, G Pascal, A Valeanu... - Annals of ..., 2004 - journals.lww.com
 ... by systemic chemotherapy and to use prognostic factors of outcome for a model predictive of ...
 1-5 However, the resectability rate of metastases at the time of diagnosis is low ... 5,6 Chemotherapy
 as a first-line treatment of metastatic colorectal cancer has greatly improved within the ...
 被引用次数: 1255 相关文章 所有 7 个版本 引用 保存

Tumor progression while on chemotherapy: a contraindication to liver resection for multiple colorectal metastases? researchgate.net 中的 [PDF]

R Adam, G Pascal, D Castaing, D Azoulay... - Annals of ..., 2004 - journals.lww.com
 ... bilobarity, number, size, resectability, serum CEA, serum CA 19-9, metastatic pedicular lymph ...
 or cryotherapy, major hepatectomy, anatomic, curative, number and size of metastases on the ...
 A multivariate analysis using a Cox model was performed for all factors emerging as ...
 被引用次数: 596 相关文章 所有 8 个版本 引用 保存

[HTML] Predicting hepatitis B virus-positive metastatic hepatocellular carcinomas using gene expression profiling and supervised machine learning nature.com 中的 [HTML]

QH Ye, LX Qin, M Forgues, P He, JW Kim, AC Peng... - Nature medicine, 2003 - nature.com
 ... levels equal to the median; red, transcript levels greater than the median; gray, missing data ... The
 authors reported that a metastasis and its primary tumor were as similar in their overall ... that the
 molecular program of a primary tumor may generally be retained in its metastases 23. ...
 被引用次数: 717 相关文章 所有 14 个版本 引用 保存

Irinotecan plus fluorouracil and leucovorin for metastatic colorectal cancer researchgate.net 中的 [PDF]

LB Saltz, JV Cox, C Blanke, LS Rosen... - ... England Journal of ..., 2000 - Mass Medical Soc
 ... primary tumor, the time from diagnosis of disease to the occurrence of metastasis, the number ...
 Prospective study of prognostic factors in patients with unresected hepatic metastases from
 colorectal cancer. ... 2015) Current and advancing treatments for metastatic colorectal cancer. ...
 被引用次数: 3018 相关文章 所有 8 个版本 引用 保存

Resection of colorectal liver metastases
 J Scheele, R Stang, A Altendorf-Hofmann... - World journal of surgery, 1995 - Springer
 ... of satellite metastases, primary tumor grade, the time of metastasis diag- nosis, diameter of the
 largest metastasis, anatomic versus ... Lost to follow-up 1 -- 2 ... deliberate palliative debulking (solid
 thin line; n = 35) or no resection (broken thin line; n = 1249) of hepatic metastases. ...

被引用次数: 1535 相关文章 所有 9 个版本 引用 保存



全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 174,000 条结果 (用时 0.60 秒)

Google 学术 : Missing metastases as a model to challenge current therapeutic algorithms in colorectal liver metastases

... surgery for unresectable colorectal liver metastases ... - Adam - 被引用次数 : 1258

... to liver resection for multiple colorectal metastases? - Adam - 被引用次数 : 598

Predicting hepatitis B virus-positive metastatic ... - Ye - 被引用次数 : 717

“Vanishing liver metastases”—A real challenge for liver ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者 : A Zendel - 2014 - 被引用次数 : 6 - 相关文章

2014年9月16日 - Expanded surgical intervention in colorectal liver metastasis (LM) and improved ... and even the chance for cure, and so it is the current paradigm of treatment (8-17). ... The logic basis behind the decision-making algorithm for DLM built on ... Preoperative imaging—Are the metastases missing indeed?

Surgical Options for Initially Unresectable Colorectal Liver ...

www.hindawi.com/journals/hpb/2012/454026/ ▾ 翻译此页

作者 : I Popescu - 2012 - 被引用次数 : 10 - 相关文章

2012年8月21日 - (2) Multiple bilobar liver metastases, the complete resection of which would not ... Cetuximab, and Panitumumab), the current survival rates for these cases are ... This therapeutic strategy is based on the observation that increasing the ... colorectal liver metastases downstaged by chemotherapy: a model to ...

缺少字词 : algorithms

Management of potentially resectable colorectal cancer liver ...

www.uptodate.com/.../management-of-potentially-resectable-co... ▾ 翻译此页

作者 : AP Venook - 被引用次数 : 5 - 相关文章

2015年11月4日 - Hepatic metastatic disease from colorectal cancer (CRC) is a ...