



Near-infrared fluorescence sentinel lymph node detection in gastric cancer



全部

图片

新闻

视频

更多 ▾

搜索工具

找到约 22,700 条结果 (用时 0.33 秒)

Google 学术: Near-infrared fluorescence sentinel lymph node detection in gastric cancer: A pilot study

... clinical trial in breast cancer sentinel lymph node ... - Trojan - 被引用次数: 369

... fluorescence imaging for sentinel lymph node detection ... - Crane - 被引用次数: 79

... detection of the sentinel lymph node in cervical cancer: ... - Crane - 被引用次数: 81

[PDF] Promotie L.M.A. Crane Binnenwerk.indd - Rijksuniversiteit...
www.rug.nl/research/portal/files/14459423/ThesisCrane.pdf

Intraoperative near-infrared fluorescence imaging for sentinel lymph node ... Optical imaging for lymph node surveillance in breast cancer patients: skin cancer and gastric cancer.10,11 A pilot study on fluorescence SLN detection in breast.

Intraoperative Multispectral Fluorescence Imaging for the ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: LMA Crane - 2011 - 被引用次数: 81 - 相关文章

2010年9月11日 - Real-time intraoperative near-infrared fluorescence (NIRF) imaging is a ... The purpose of this technical feasibility pilot study was to evaluate the ... Over the last decades, the sentinel lymph node (SLN) concept has proven green (ICG) and infrared ray detection in early gastric cancer: an accurate method ...



23710-Review

BY ANTONIO FACCIORUSSO

Quotes Excluded
Bibliography Excluded3%
SIMILAR

1

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 23710

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Lymphocyte-to-monocyte ratio predicts survival after radiofrequency ablation for colorectal liver metastases

Antonio Facciorusso, Valentina Del Prete, Nicola Crucinio, Gaetano Serviddio, Gianluigi Vendemiale, Nicola Muscatiello

Match Overview

1

CrossCheck 74 words

Facciorusso, Antonio, Valentina Del Prete, Matteo Antonino , Viviana Neve, Nicola Crucinio, Alfredo Di Leo, Brian I Carr

2%

2

CrossCheck 53 words

Facciorusso, Antonio, Valentina Del Prete, Matteo Antonino , Viviana Neve, Annabianca Amoroso, Nicola Crucinio, Alfr

1%



Near-infrared fluorescence sentinel lymph node detection in gastric cancer



全部

图片

新闻

视频

更多 ▾

搜索工具

找到约 22,700 条结果 (用时 0.33 秒)

Google 学术: Near-infrared fluorescence sentinel lymph node detection in gastric cancer: A pilot study

... clinical trial in breast cancer sentinel lymph node ... - Trojan - 被引用次数: 369

... fluorescence imaging for sentinel lymph node detection ... - Crane - 被引用次数: 79

... detection of the sentinel lymph node in cervical cancer: ... - Crane - 被引用次数: 81

[PDF] Promotie L.M.A. Crane Binnenwerk.indd - Rijksuniversiteit...
www.rug.nl/research/portal/files/14459423/ThesisCrane.pdf

Intraoperative near-infrared fluorescence imaging for sentinel lymph node ... Optical imaging for lymph node surveillance in breast cancer patients: skin cancer and gastric cancer.10,11 A pilot study on fluorescence SLN detection in breast.

Intraoperative Multispectral Fluorescence Imaging for the ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者: LMA Crane - 2011 - 被引用次数: 81 - 相关文章

2010年9月11日 - Real-time intraoperative near-infrared fluorescence (NIRF) imaging is a ... The purpose of this technical feasibility pilot study was to evaluate the ... Over the last decades, the sentinel lymph node (SLN) concept has proven green (ICG) and infrared ray detection in early gastric cancer: an accurate method ...

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 17,900 条结果 (用时 0.62 秒)

PubMed Result

www.ncbi.nlm.nih.gov/pubmed?LinkName=pubmed...uid... - [翻译此页](#)

Usefulness of the neutrophil-to-lymphocyte ratio in predicting short- and long-term ...
for colorectal liver metastasis after percutaneous radiofrequency ablation. ... ratio is
elevated in epithelial ovarian cancer and predicts survival after treatment. ... to the
Lymphocyte-To-Monocyte Ratio for Predicting the Long-Term Survival of ...

Elevated preoperative peripheral blood monocyte count ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者：SL Shen - 2014 - 被引用次数：4 - 相关文章

2014年10月3日 - Elevated preoperative peripheral blood monocyte count predicts poor
prognosis for ... However, even after hepatic resection, the long-term prognosis has ...
and platelet lymphocyte ratio (PLR) have prognostic value in cancer patients [5]. ...
factor of poor prognosis for patients with colorectal liver metastasis, ...

PubMed Result

www.ncbi.nlm.nih.gov/pubmed?db=pubmed&cmd... - [翻译此页](#)

Elevated neutrophil to lymphocyte ratio predicts poor prognosis in nasopharyngeal ...
for colorectal liver metastasis after percutaneous radiofrequency ablation.

Lymphocyte-to-monocyte ratio predicts survival of patients ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者：ZX Lin - 2015 - 被引用次数：1

2015年10月14日 - Lymphocyte-to-monocyte ratio predicts survival of patients with
hepatocellular ... anticancer therapy or intraoperative radiofrequency ablation. ... been
used to predict survival in patients with HCC after curative resection[3-6]. Patients
with LMR \leq 3.23 were also prone to have liver cirrhosis ($P = 0.003$), ...