



Suppository Naproxen Reduces Incidence and Severity of Post-E



网页

图片

新闻

视频

购物

更多

搜索工具

找到约 5,220 条结果 (用时 0.78 秒)

A Randomized Trial of Rectal Indomethacin to Prevent Post ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) > 翻译此页

作者: BJ Elmunzer - 2012 - 被引用次数: 220 - 相关文章

2012年4月12日 - The primary outcome was **post-ERCP pancreatitis**, which was defined as ... there is no conclusive evidence from **randomized, controlled trials** and because ... The **suppositories** were administered immediately after ERCP while the **reduced** the **incidence and severity of post-ERCP pancreatitis** in patients ...

缺少字词: **naproxen**

NSAIDs and Acute Pancreatitis: A Systematic Review

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) > 翻译此页

作者: R Pezzilli - 2010 - 被引用次数: 10 - 相关文章

2010年3月10日 - On the contrary, in a case-control study carried out in Sweden in 2002 [94], ... 95% CI: 4.2–5.9) and the lowest for **naproxen** (OR 1.1, 95% CI: 0.7–1.7). ... were **randomized controlled trials**, they had different modalities of **ERCP** may **reduce** the **incidence and severity** of acute **pancreatitis after ERCP**.

Indomethacin Reduces Post-ERCP Pancreatitis Risk

www.medscape.com/viewarticle/761933 > 翻译此页


2012年4月11日 - A single indomethacin **suppository** administered immediately after endoscopic ... because there is no conclusive evidence from **randomized controlled trials**. ... **reduced** the **incidence and severity of post-ERCP pancreatitis** in ...

缺少字词: **naproxen**

The Prevention of Post-ERCP Pancreatitis | Clinical ...

www.clinicalcorrelations.org/?p=7209 > 翻译此页

作者: SK Patel - 相关文章



ESPS Manuscript NO: 20893
Manuscript Type: ORIGINAL ARTICLE
Randomized Controlled Trial

Suppository naproxen reduces incidence and severity of post-endoscopic retrograde cholangiopancreatography pancreatitis: Randomized controlled trial

Mansour-Ghanaei F *et al.* Naproxen for post-ERCP pancreatitis

Fariborz Mansour-Ghanaei, Farahnaz Joukar, Zahra Taherzadeh, Homayoon Sokhanvar, Tolou Hasandokht

15
Abstract

AIM: To determine the efficacy of rectally administered naproxen for the prevention of post-endoscopic retrograde cholangiopancreatography (ERCP) pancreatitis (PEP).

METHODS: This double-blind randomized control trial conducted from January

Match Overview

1	CrossCheck 76 words Taiga Otsuka. "Low-dose rectal diclofenac for prevention of post-endoscopic retrograde cholangiopancreatogra ...	2%
2	Internet 69 words crawled on 28-Nov-2015 www.biomedcentral.com	1%
3	Internet 54 words crawled on 29-Jul-2010 www.wjgnet.com	1%
4	CrossCheck 49 words Rasoul Sotoudehmanesh. "Indomethacin May Reduce ... he Incidence and Severity of Acute Pancreatitis After ER	1%
5	CrossCheck 44 words Katsinelos, P.. "Intravenous N-acetylcysteine does not ... event post-ERCP pancreatitis", <i>Gastrointestinal Endosc</i>	1%
6	CrossCheck 32 words Solhjoo, Ehsan, Fariborz Mansour-Ghanaei, Roghaeyh Moulaei-Lahgorudi, and Farahnaz Joukar. "Comparison	1%
7	Internet 28 words crawled on 12-Feb-2011 www.mdpi.com	1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 2,560 条结果 (用时 0.69 秒)

Google 学术: Suppository naproxen reduces incidence and severity of post-endoscopic retrograde cholangiopancreatography pancreatitis: Randomized controlled trial

... NSAIDs in the prevention of post-ERCP pancreatitis - Elmunzer - 被引用次数: 210

[NSAIDs and acute pancreatitis: A Systematic Review](#) - Pezzilli - 被引用次数: 10

NSAIDs and Acute Pancreatitis: A Systematic Review

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: R Pezzilli - 2010 - 被引用次数: 10 - [相关文章](#)

2010年3月10日 - The Prophylactic Use of NSAIDs for Preventing Post-ERCP Acute ... were randomized controlled trials, they had different modalities of ERCP ... In the first controlled double-blind study, indomethacin, in the form of suppositories at a may reduce the incidence and severity of acute pancreatitis after ERCP.

A Randomized Trial of Rectal Indomethacin to Prevent Post ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: BJ Elmunzer - 2012 - 被引用次数: 254 - [相关文章](#)

2012年4月12日 - The primary outcome was post-ERCP pancreatitis, which was defined ... Moderate-to-severe pancreatitis developed in 13 patients (4.4%) in the ... pancreatitis, rectal indomethacin significantly reduced the incidence of the condition. ... no conclusive evidence from randomized, controlled trials and because ...

缺少字词: naproxen



[全部](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 2,250 条结果 (用时 0.70 秒)

Google 学术 : Suppository naproxen reduces incidence and severity of post-endoscopic retrograde cholangiopancreatography pancreatitis: Randomized controlled trial

... NSAIDs in the prevention of post-ERCP pancreatitis - Elmunzer - 被引用次数 : 211

NSAIDs and acute pancreatitis: A Systematic Review - Pezzilli - 被引用次数 : 10

NSAIDs and Acute Pancreatitis: A Systematic Review

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : R Pezzilli - 2010 - 被引用次数 : 10 - 相关文章

2010年3月10日 - The Prophylactic Use of NSAIDs for Preventing Post-ERCP Acute ... were randomized controlled trials, they had different modalities of ERCP ... In the first controlled double-blind study, indomethacin, in the form of suppositories at a may reduce the incidence and severity of acute pancreatitis after ERCP.

A Randomized Trial of Rectal Indomethacin to Prevent Post ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : BJ Elmunzer - 2012 - 被引用次数 : 255 - 相关文章

2012年4月12日 - The primary outcome was post-ERCP pancreatitis, which was defined ... Moderate-to-severe pancreatitis developed in 13 patients (4.4%) in the ... pancreatitis, rectal indomethacin significantly reduced the incidence of the condition. ... no conclusive evidence from randomized, controlled trials and because ...

缺少字词 : naproxen

The Prevention of Post-ERCP Pancreatitis | Clinical ...

www.clinicalcorrelations.org/?p=7209 ▾ [翻译此页](#)