

[学术搜索](#)

获得 7 条结果 (用时0.06秒)

[文章](#)
[我的图书馆](#)
[时间不限](#)
[2016以来](#)
[2015以来](#)
[2012以来](#)
[自定义范围...](#)
[按相关性排序](#)
[按日期排序](#)
[搜索所有网页](#)
[中文网页](#)
[简体中文网页](#)
☒ 包括专利

☒ 包含引用

 [创建快讯](#)

小提示： 只搜索 **中文(简体)** 结果，可在 [学术搜索设置](#) 指定搜索语言

Receptor activation of NADPH oxidases

A Petry, M Weitnauer, A Görlach - Antioxidants & redox ..., 2010 - online.liebertpub.com
 ... in vascular cells or macrophages in response to different stimuli including insulin, thrombin, urotensin-II ... NOX4 expression can also be induced by urotensin-II in pulmonary artery smooth muscle ... c-Abl was shown to phosphorylate NOX5 thus enhancing its calcium sensitivity (59).....
 被引用次数: 105 [相关文章](#) [所有 4 个版本](#) [引用](#) [保存](#)

Regulation of NADPH oxidase in vascular endothelium: the role of phospholipases, protein kinases, and cytoskeletal proteins

S Pendyala, PV Usatyuk, IA Gorshkova... - ... & redox signaling, 2009 - online.liebertpub.com
 ... man urotensin II (hU-II) activated NADPH oxidase, which was abrogated by p22phox or Nox4 ... Nox4, but not Nox2, appeared to be a major mediator of insulin-induced ROS ... the alveolar capillary endothelium and epithelium, resulting in arterial hypoxemia, resistant to oxygen ...
 被引用次数: 78 [相关文章](#) [所有 6 个版本](#) [引用](#) [保存](#)

Redox regulation of the coagulation cascade

A Görlach - Antioxidants & redox signaling, 2005 - online.liebertpub.com
 ... In vitro studies confirmed redox sensitivity of PAI-1 induction in response to HNE and AGE ... the hypoxia-inducible transcription factor HIF-1 (30, 32), oxidative stress and insulin enhanced PAI ... T, BelAiba RS, Bonello S, Pfeilschifter J, Hess J, and Gørlach A. Human urotensin II is a ...
 被引用次数: 32 [相关文章](#) [所有 3 个版本](#) [引用](#) [保存](#)

The hypoxia-inducible factor-2α is stabilized by oxidative stress involving NOX4

I Diebold, D Flügel, S Becht... - ... & redox signaling, 2010 - online.liebertpub.com
 ... these two transcription factors, suggesting differences regarding target gene specificity, sensitivity to oxygen ... T. Transcriptional regulation of plasminogen activator inhibitor-1 expression by insulin-like growth ... Bonello S, Pfeilschifter J, Hess J, and Gørlach A. Human urotensin II is ...
 被引用次数: 41 [相关文章](#) [所有 6 个版本](#) [引用](#) [保存](#)

[图书] Signaling Through NADPH Oxidases: Role in Vascular Remodeling Processes

T Djordjevic - 2005 - core.ac.uk
 ... involving HIF-1 121 4.3 NADPH oxidase-derived ROS mediate urotensin-II-induced signaling and ... pulmonary artery pressure and pulmonary vascular resistance which are primarily due to ... Vessel Human urotensin-II -log (EC) Endothelin-1 -log (EC) Relative potency ...
[相关文章](#) [引用](#) [保存](#)

[researchgate.net 中的 \[PDF\]](#)
[core.ac.uk 中的 \[PDF\]](#)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 25296

Manuscript Type: Original Article

Basic Study

Urotensin II-induced insulin resistance are mediated by NADPH oxidase-derived reactive oxygen species in HepG2 cells

Li YY *et al.* Urotensin II induced HepG2 cells insulin resistance

Ying-Ying Li, Zheng-Ming Shi, Xiao-Yong Yu, Ping Feng, Xue-Jiang Wang

Ying-Ying Li, Xiao-Yong Yu, Xue-Jiang Wang, Department of Physiology and Pathophysiology, School of Basic Medical Sciences, Capital Medical University, Beijing 100069, China

Ying-Ying Li, Xiao-Yong Yu, Xue-Jiang Wang, Beijing Key Laboratory for Cancer Invasion and Metastasis Research, Capital Medical University, Beijing

Match Overview

1	Internet 180 words crawled on 15-Mar-2016 www.wjgnet.com	5%
2	Internet 30 words crawled on 24-Nov-2015 www.stemcellres.com	1%

[学术搜索](#)

获得 7 条结果 (用时0.12秒)

[文章](#)
[我的图书馆](#)

 小提示: 只搜索[中文\(简体\)](#)结果, 可在 [学术搜索设置](#) 指定搜索语言

Receptor activation of NADPH oxidases

A Petry, M Weitnauer, A Görlach - Antioxidants & redox ..., 2010 - online.liebertpub.com
 ... in vascular cells or macrophages in response to different stimuli including insulin, thrombin, urotensin-II ... NOX4 expression can also be induced by urotensin-II in pulmonary artery smooth muscle ... c-Abl was shown to phosphorylate NOX5 thus enhancing its calcium sensitivity (59). ...
 被引用次数: 106 相关文章 所有 4 个版本 引用 保存

Regulation of NADPH oxidase in vascular endothelium: the role of phospholipases, protein kinases, and cytoskeletal proteins

S Pendyala, PV Usatyuk, IA Gorshkova... - ... & redox signaling, 2009 - online.liebertpub.com
 ... man urotensin II (hU-II) activated NADPH oxidase, which was abrogated by p22phox or Nox4 ... Nox4, but not Nox2, appeared to be a major mediator of insulin-induced ROS ... the alveolar capillary endothelium and epithelium, resulting in arterial hypoxemia, resistant to oxygen ...
 被引用次数: 80 相关文章 所有 6 个版本 引用 保存

Redox regulation of the coagulation cascade

A Görlach - Antioxidants & redox signaling, 2005 - online.liebertpub.com
 ... In vitro studies confirmed redox sensitivity of PAI-1 induction in response to HNE and AGE ... the hypoxia-inducible transcription factor HIF-1 (30, 32), oxidative stress and insulin enhanced PAI ... T, BelAiba RS, Bonello S, Pfeilschifter J, Hess J, and Gorlach A. Human urotensin II is a ...
 被引用次数: 34 相关文章 所有 3 个版本 引用 保存

The hypoxia-inducible factor-2α is stabilized by oxidative stress involving NOX4

I Diebold, D Flügel, S Becht... - ... & redox signaling, 2010 - online.liebertpub.com
 ... these two transcription factors, suggesting differences regarding target gene specificity, sensitivity to oxygen ... T. Transcriptional regulation of plasminogen activator inhibitor-1 expression by insulin-like growth ... Bonello S, Pfeilschifter J, Hess J, and Gorlach A. Human urotensin II is ...
 被引用次数: 43 相关文章 所有 6 个版本 引用 保存

[researchgate.net 中的 \[PDF\]](#)

[图书] Signaling Through NADPH Oxidases: Role in Vascular Remodeling Processes

T Djordjevic - 2005 - core.ac.uk
 ... involving HIF-1 121 4.3 NADPH oxidase-derived ROS mediate urotensin-II-induced signaling and ... pulmonary artery pressure and pulmonary vascular resistance which are primarily due to ... Vessel Human urotensin-II -log (EC) Endothelin-1 -log (EC) Relative potency ...
 相关文章 引用 保存 更多


[core.ac.uk 中的 \[PDF\]](#)

Signaling through NADPH oxidases

T Djordjevic - 2006 - edoc.ub.uni-muenchen.de
 ... involving HIF-1 121 4.3 NADPH oxidase-derived ROS mediate urotensin II-induced signaling

[uni-muenchen.de 中的 \[PDF\]](#)

- ☒ 包括专利
- ☒ 包含引用

 [创建快讯](#)

[全部](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 4,580 条结果 (用时 0.66 秒)

Google 学术 : Urotensin II-induced insulin resistance are mediated by NADPH oxidase-derived reactive oxygen species in HepG2 cells

Receptor activation of NADPH oxidases - Petry - 被引用次数 : 106

Regulation of NADPH oxidase in vascular endothelium ... - Pendyala - 被引用次数 : 80

Redox regulation of the coagulation cascade - Görlach - 被引用次数 : 33

The Effects of Palmitate on Hepatic Insulin Resistance Are M...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : D Gao - 2010 - 被引用次数 : 124 - [相关文章](#)

2010年7月20日 - NADPH oxidase 3 (NOX3)-derived reactive oxygen species (ROS) may ... We found that palmitate induced hepatic insulin resistance through ... 2, D and E, ROS levels were increased in HepG2 cells by exposure to palmitate.

缺少字词 : urotensin

NADPH Oxidase 4 is required for interleukin-1 β -mediated acti...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : R Ginnan - 2013 - 被引用次数 : 4 - [相关文章](#)

2012年9月26日 - Reactive oxygen species (ROS) are generated in the vascular wall To this end, we pre-incubated VSM cells with the H2O2 scavenger ... IL-1 β -induced JNK activation is ROS-dependent in rat aortic VSM The effects of palmitate on hepatic insulin resistance are mediated by NADPH Oxidase 3-derived ...

缺少字词 : hepg2

Qizhi Jiangtang Jiaonang Improves Insulin Signaling and Red...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - [翻译此页](#)