

学术搜索

获得 9 条结果 (用时0.11秒)

试用新功能!

小提示: 只搜索中文(简体)结果, 可在 [学术搜索设置](#) 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☒ 包含引用

[PDF] Ischemia-reperfusion injury and ischemic-type biliary lesions following liver transplantation

hindawi.com 中的 [PDF]

R Cursio, J Gugenheim - Journal of transplantation, 2012 - downloads.hindawi.com

... ITBLs were also termed as "ischemic-type biliary complications or ITBC" [3], "ischemic cholangitis" [34] ... use of extended-criteria donor (ECD) organs, including organs, which present risk factors mentioned ... did not show that chronic graft rejection was a risk factor for development ...

被引用次数: 40 相关文章 所有 11 个版本 引用

Intrathymic transplantation of allogeneic nonimmunogenic perinatal islet tissue does not induce donor-specific tolerance.

M BUNKE, B GANZEL, JONB KLEIN... - ..., 1993 - journals.lww.com

... (1). However, several factors must be ... Prolongation of islet allograft survival following in vitro culture (24°C) and a single injection of ALS. ... 1024, 2048 1024,1024 " Thymectomy of BCBA mice was at 4 weeks and bone marrow transplantation at 12 weeks after conditioning with ...

被引用次数: 7 相关文章 所有 4 个版本 引用

[HTML] Intravenous immunoglobulins in liver transplant patients: Perspectives of clinical immune modulation

nih.gov 中的 [HTML]

A Kornberg - World journal of hepatology, 2015 - ncbi.nlm.nih.gov

... accepted, which may, however, aggravate the patients' infectious and immunologic risk profiles. ... Ig against hepatitis B virus is current standard in post-LT antiviral ... of an escalating organ scarcity, decreasing posttransplant survival rates following liver transplantation have been ...

被引用次数: 1 相关文章 所有 8 个版本 引用

Rotterdam: Main port for organ transplantation research in the Netherlands

J Kwekkeboom, LJJ van der Laan, MGH Betjes... - Transplant ..., 2014 - Elsevier

... An important factor determining transplantation outcome is damage to the graft during procurement, cold ... Moreover we found that resident human liver MSC secrete factors that promote liver regeneration ... UGT1A9 were found to associate with MPA exposure and rejection risk [37] ...

相关文章 所有 4 个版本 引用

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 26479

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Cohort Study*

Risk factor for ischemic-type biliary lesion after **ABO-incompatible living donor liver transplantation**

**Jun Bae Bang, Bong-Wan Kim, Young Bae Kim, Hee-Jung Wang, Hyun Yeong Lee, Joohyun Sim, Taegyu Kim, Kyeong Lok Lee, Xu-Guang Hu, Wei Mao**

#### Abstract

**AIM:** To evaluate the risk factors for ischemic-type biliary lesion (ITBL) after ABO-incompatible (ABO-I) adult living donor liver transplantation (ALDLT).

**METHODS:** Among 141 ALDLTs performed in our hospital between 2008 and 2014, 27 (19%) were ABO-I ALDLT and 114 were ABO-identical/compatible ALDLT. In

#### Match Overview

1	<b>Internet</b> 56 words crawled on 23-Apr-2016 <a href="http://www.wjnet.com">www.wjnet.com</a>	1%
2	<b>CrossCheck</b> 41 words Resch, Thomas, Cornelia Fabritius, Susanne Ebner, Paul Ritschl, and Katja Kotsch. "The Role of Natural Killer ...	1%
3	<b>Internet</b> 31 words crawled on 17-Mar-2015 <a href="http://onlinelibrary.wiley.com">onlinelibrary.wiley.com</a>	1%
4	<b>CrossCheck</b> 15 words "Poster Panel [Liver Transplantation]", HPB, 2014.	<1%
5	<b>CrossCheck</b> 13 words "Abstracts .", Transplantation Journal, 2015.	<1%
6	<b>CrossCheck</b> 11 words Daisuke Morioka. "Six Consecutive Cases of Successful Adult ABO-Incompatible Living Donor Liver Transplants ...	<1%
7	<b>Internet</b> 11 words crawled on 27-May-2012 <a href="http://www.wjnet.com">www.wjnet.com</a>	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 11,200 条结果 (用时 0.66 秒)

## Google 学术: Risk factor for ischemic-type biliary lesion after ABO-incompatible living donor liver transplantation

... a risk factor for ischemic - type biliary lesions following ... - Moench - 被引用次数: 93

... reconstruction in living donor liver transplantation - Park - 被引用次数: 64

Biliary complications after liver transplantation from ... - Suárez - 被引用次数: 85

## Ischemia-Reperfusion Injury and Ischemic-Type Biliary Lesions ...

[www.hindawi.com/journals/jtrans/2012/164329/](http://www.hindawi.com/journals/jtrans/2012/164329/) ▾ [翻译此页](#)

作者: R Cursio - 2012 - 被引用次数: 43 - [相关文章](#)

2011年12月23日 - Ischemia-reperfusion (I-R) injury after liver transplantation (LT) induces .... injury (ABO incompatibility, rejection, autoimmune hepatic disease, CMV ... Risk factors involved in the development of ITBLs are old donor age [36, 37], ..... Europe and North America and in living-related liver transplantation (LRLT) ...

## [PDF] Incidence of and risk factors for ischemic-type biliary lesions following...

[www.custodiol.com/hansjb/wp.../09/HeidenhainIncidenceRiskFactors.pdf](http://www.custodiol.com/hansjb/wp.../09/HeidenhainIncidenceRiskFactors.pdf) ▾ [翻译此页](#)

作者: C Heidenhain - 2009 - 被引用次数: 115 - [相关文章](#)

Ischemic-type biliary lesions (ITBL) account for a major part of patients' mor- bidity and .... split-liver transplantation, living versus deceased donor and the use of ...

## The role of endoscopy in biliary complications after liver transplantation

[www.uptodate.com/.../the...biliary...after-liver-transplantation/.../7,12,17-23](http://www.uptodate.com/.../the...biliary...after-liver-transplantation/.../7,12,17-23) - [翻译此页](#)

Moreover, partial LT or living donor liver transplantation (LDLT) and donation after ... Especially



全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 12,500 条结果 (用时 0.66 秒)

## Google 学术 : Risk factor for ischemic-type biliary lesion after ABO-incompatible living donor liver transplantation

... a risk factor for ischemic-type biliary lesions following ... - Moench - 被引用次数 : 92

... reconstruction in living donor liver transplantation - Park - 被引用次数 : 64

Biliary complications after liver transplantation from ... - Suárez - 被引用次数 : 87

## Ischemia-Reperfusion Injury and Ischemic-Type Biliary Lesions ...

[www.hindawi.com/journals/jtrans/2012/164329/](http://www.hindawi.com/journals/jtrans/2012/164329/) ▾ 翻译此页

作者 : R Cursio - 2012 - 被引用次数 : 45 - 相关文章

2011年12月23日 - Ischemia-reperfusion (I-R) injury after liver transplantation (LT) induces .... injury (ABO incompatibility, rejection, autoimmune hepatic disease, CMV ... Risk factors involved in the development of ITBLs are old donor age [36, 37], .... Europe and North America and in living-related liver transplantation (LRLT) ...

## Aetiology and risk factors of ischaemic cholangiopathy after liver ...

[www.ncbi.nlm.nih.gov/NCBI/Literature/PubMed Central \(PMC\)](http://www.ncbi.nlm.nih.gov/NCBI/Literature/PubMedCentral(PMC)) ▾ 翻译此页

作者 : MM Mourad - 2014 - 被引用次数 : 14 - 相关文章

2014年5月28日 - Core tip: Biliary complications after liver transplantation represent a major challenge. ... or secondary to immunological causes (ABO-incompatibility, rejection, and ... can cause ischaemic-type biliary lesions and segmental strictures[48]. .... Donor age is a risk factor for the development of IC after LT[73,84].

## [PDF] Incidence of and risk factors for ischemic-type biliary lesions followin...

[www.custodiol.com/hansjb/wp.../09/HeidenhainIncidenceRiskFactors.pdf](http://www.custodiol.com/hansjb/wp.../09/HeidenhainIncidenceRiskFactors.pdf) ▾ 翻译此页

作者 : C Heidenhain - 2009 - 被引用次数 : 117 - 相关文章

Ischemic-type biliary lesions (ITBL) account for a major part of patients' mor- bidity and .... split-liver transplantation, living versus deceased donor and the use of ...