

Mitochondrial DNA from hepatocytes as a ligand for TLR9: Drivers of NASH?



全部 图片 视频 新闻 购物 地图 图书

找到约 16,700 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Google 学术: Mitochondrial DNA from hepatocytes as a ligand for TLR9: Drivers of NASH?](#)

[Apoptosis as a mechanism for liver disease ...](#) - Guicciardi - 被引用次数: 104

[... in the pathogenesis of NASH: beyond steatosis and ...](#) - Peverill - 被引用次数: 39

[Mechanisms and biomarkers of apoptosis in liver ...](#) - Chakraborty - 被引用次数: 51

[Hepatocyte mitochondrial DNA drives nonalcoholic steatohepatitis ...](#)

www.ncbi.nlm.nih.gov/pubmed/26808498

25 Jan 2016 ... Hepatocyte mitochondrial DNA drives nonalcoholic steatohepatitis by ... to identify NASH-associated ligands for TLR9, establish the cellular ...

[Wajahat Z Mehal - Publications - ResearchGate](#)

https://www.researchgate.net/profile/Wajahat_Mehal/publications

Article: Hepatocyte mitochondrial DNA drives nonalcoholic steatohepatitis by and Mice With NASH Have Increased Plasma Levels of TLR9 Ligand, and a ...

[\[PDF\] Evolving Concepts in the Pathogenesis of NASH: Beyond Steatosis ...](#)

www.mdpi.com/1422-0067/15/5/8591/pdf

14 May 2014 ... appears to be the central driver of cellular injury via oxidative stress. ... hepatocyte dysfunction leading to disease progression in NASH. Mitochondrial ROS, induced by SFAs, appear to contribute to JNK activation and cellular insulin tumour

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 26383

Manuscript Type: EDITORIAL

Mitochondrial DNA from hepatocytes as a ligand for TLR9: Drivers of nonalcoholic steatohepatitis?

Priya Handa, Akhila Vemulakonda, Kris V Kowdley, Misael Uribe, Nahum Méndez-Sánchez

1

Match Overview

1	CrossCheck 123 words Tilg, Herbert, Alexander R Moschen, and Gyongyi Szal"Interleukin-1 and inflammasomes in ALD/AAH and NA	4%
2	CrossCheck 44 words Hoque, Rafaz, Yoram Vodovotz, and Wajahat Mehal. "... erapeutic strategies in inflammasome mediated disease	2%
3	Publications 43 words "Researchers' Work from University of South Carolina F... cuses on Inflammation (Sparstolonin B attenuat", Health	2%
4	Internet 25 words crawled on 04-Mar-2016 www.wjgnet.com	1%
5	CrossCheck 21 words Arrese, Marco, Daniel Cabrera, Alexis M. Kalergis, and ... riel E. Feldstein. "Innate Immunity and Inflammation in N	1%
6	CrossCheck 20 words Friedman, Scott, Arun Sanyal, Zachary Goodman, Eric L efebvre, Mildred Gottwald, Laurent Fischer, and Vlad Rat	1%

Mitochondrial DNA from hepatocytes as a ligand for TLR9: Drivers of nonalco



全部 图片 视频 新闻 购物 地图 图书

找到约 19,700 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Hepatocyte mitochondrial DNA drives nonalcoholic steatohepatitis ...](#)

www.ncbi.nlm.nih.gov/pubmed/26808498

25 Jan 2016 ... Hepatocyte mitochondrial DNA drives nonalcoholic steatohepatitis by ... to identify NASH-associated ligands for TLR9, establish the cellular ...

[\[PDF\] Standard PDF - Wiley Online Library](#)

onlinelibrary.wiley.com/doi/10.1002/hep.22306/pdf

nonimmune cell populations, including hepatocytes, biliary epithelial cells, endothelial ... in the pathophysiology of nonalcoholic steatohepatitis ... spectively, driving the activation of mitogen-activated ... tolerance," whereas CpG -DNA(TLR9 ligand) pretreat- ... the mitochondrial membrane.58,130-132 Cleaved IPS-1 re-

[\[PDF\] Full-Text PDF - MDPI.com](#)

www.mdpi.com/1422-0067/15/5/8591/pdf

14 May 2014 ... appears to be the central driver of cellular injury via oxidative stress. ... Hepatocyte death and an ensuing inflammatory cascade likely ... Originally, the progression of NAFLD from simple steatosis to NASH was conceptualised as a ... Mitochondrial ROS, induced by SFAs, appear to contribute to JNK ...

[TLR9 activation in nonalcoholic steatohepatitis - YouTube](#)

全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 20,000 条结果 (用时 0.67 秒)

Hepatocyte mitochondrial DNA drives nonalcoholic steatohepatitis by ...

www.ncbi.nlm.nih.gov/pubmed/26808498 - 翻译此页

作者: I Garcia-Martinez - 2016

2016年1月25日 - **Hepatocyte mitochondrial DNA** drives **nonalcoholic steatohepatitis** by ... to identify NASH-associated **ligands for TLR9**, establish the cellular ...

Standard PDF - Wiley Online Library

onlinelibrary.wiley.com/doi/10.1002/hep.22306/pdf ▾ 翻译此页

作者: E Seki - 2008 - 被引用次数: 464 - 相关文章

nonimmune cell populations, including **hepatocytes**, biliary epithelial cells, endothelial in the pathophysiology of **nonalcoholic steatohepatitis** spectively, **driving** the activation of mitogen-activated tolerance," whereas CpG-DNA(**TLR9 ligand**) pretreat- the **mitochondrial** membrane.58,130-132 Cleaved IPS-1 re-

TLR9 activation in nonalcoholic steatohepatitis - YouTube



<https://www.youtube.com/watch?v=rIBlw8kGlnA>

2016年1月25日 - 上传者: Journal of Clinical Investigation

Nonalcoholic steatohepatitis (NASH) is the most common liver ... of this study identify **TLR9** activation via ...

缺少字词: hepatocytes ligand

Unbound MEDLINE | J Clin Invest journal articles from PubMed

www.unboundmedicine.com/medline/journal/J_Clin_Invest?start=120... ▾ 翻译此页

Hepatocyte mitochondrial DNA drives **nonalcoholic steatohepatitis** by activation of **TLR9**. [Journal Article]; J Clin Invest 2016 Mar 1; 126(3):859-64.

[PDF] Full-Text PDF - MDPI.com

www.mdpi.com/1422-0067/15/5/8591/pdf ▾ 翻译此页

作者: W Peverill - 2014 - 被引用次数: 46 - 相关文章

2014年5月14日 - appears to be the central **driver** of cellular injury via oxidative stress. ... **Hepatocyte**