

全部 新闻 图片 视频 购物 更多 ▾ 搜索工具

找到约 11,900 条结果 (用时 0.66 秒)

您是不是要找：**A total of 539 successive *phases* of pancreaticoduodenectomy: Analysis of the risk factors for postoperative pancreatic fistula**

Analysis of risk factors for postoperative pancreatic fistula ...

www.ncbi.nlm.nih.gov/pubmed/25516663 - 翻译此页

作者：QY Liu - 2014 - 被引用次数：7 - 相关文章

2014年12月14日 - Analysis of risk factors for postoperative pancreatic fistula following ... METHODS: The data from 196 consecutive patients who underwent pancreaticoduodenectomy, ... of the patients, and the incidence of CR-POPF was 32.7% (64/196). ... there were three postoperative deaths, giving a total mortality rate of ...

Risk factors and outcomes of postoperative pancreatic ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者：SJ Fu - 2015 - 被引用次数：2 - 相关文章

2015年3月26日 - PF was found in 65 (12.2%) cases, of whom 11 were classified into ISGPF grade A, ... Univariate and multivariate analysis showed that blood loss \geq 500 ml, ... Keywords: Pancreatic fistula, Pancreaticoduodenectomy, Risk factor, Outcome. Go to: A total of 303 (57.0%, 303/532) patients didn't develop any ...

Pancreatic fistula after pancreatectomy: Evolving definitions ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ 翻译此页

作者：SV Shrikhande - 2008 - 被引用次数：67 - 相关文章

2008年10月14日 - Keywords: Pancreatic fistula, Pancreaticoduodenectomy, Pancreatic ... that post-operative intra-abdominal collections after PD were PF. The rationale for total pancreatectomy is that it allows a more 2005;92:539-546. ... Distal pancreatectomy: risk factors for surgical failure in 302 consecutive cases.

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 26587

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Total of 539 successive cases of pancreaticoduodenectomy: Analysis of the risk factors for postoperative pancreatic fistula

Match Overview

- 1** **CrossCheck** 17 words **<1%**
Kim, Eun Young, Young Kyoung You, Dong Goo Kim, and Tae Ho Hong. "A simple pancreaticojejunostomy technique f
- 2** **CrossCheck** 15 words **<1%**
Su, An-Ping, Yi Zhang, Neng-Wen Ke, Hui-Min Lu, Bo-Le Tian, Wei-Ming Hu, and Zhao-Da Zhang. "Triple-layer duct-to-
- 3** **Internet** 14 words **<1%**
crawled on 15-Jul-2016
www.wjgnet.com

Total of 539 successive phases of pancreaticoduodenectomy: Analysis of the risl



全部 图片 视频 新闻 购物 地图 图书

找到约 256 条结果

时间不限

- 过去 1 小时内
- 过去 24 小时内
- 过去 1 周内
- 过去 1 个月内
- 过去 1 年内

所有结果

精确匹配

[Analysis of risk factors for postoperative pancreatic fistula following ...](#)

www.ncbi.nlm.nih.gov/pubmed/25516663

14 Dec 2014 ... Analysis of risk factors for postoperative pancreatic fistula following ... from 196 consecutive patients who underwent pancreaticoduodenectomy, ... there were three postoperative deaths, giving a total mortality rate of 1.5% ...

[Risk factors of postoperative complications of ... - NCBI](#)

www.ncbi.nlm.nih.gov/pubmed/25713915

Risk factors of postoperative complications of pancreatoduodenectomy: ... postoperative mortality and morbidity were selected for univariate analysis and ... pre-operative drainage, total serum bilirubin level, alanine aminotransferase, serum ... Pancreatic duct diameter was the independent risk factor of pancreatic fistula.

[Pancreatoduodenectomy: Risk Factors of Postoperative Pancreatic ...](#)

www.ncbi.nlm.nih.gov/pubmed/26158175

Pancreatoduodenectomy: Risk Factors of Postoperative Pancreatic Fistula ... present study is the analysis of risk factors of postoperative pancreatic fistula (POPF) and of clinical outcome after pancreatoduodenectomy (PD) in a retrospective multicentric review ... Forty % grade A, 13% grade B and 47% grade C total POPF.

[\[PDF\] Pancreatic fistula after pancreaticoduodenectomy: A comparison ...](#)

www.wignet.com/esns/DownloadFile.aspx?Type



全部 新闻 图片 购物 地图 更多 ▾ 搜索工具

找到约 5,160 条结果 (用时 0.58 秒)

Analysis of risk factors for postoperative pancreatic fistula following ...

www.ncbi.nlm.nih.gov/pubmed/25516663 ▾ 翻译此页

作者: QY Liu - 2014 - 被引用次数: 13 - 相关文章

2014年12月14日 - Analysis of risk factors for postoperative pancreatic fistula following ... METHODS: The data from 196 consecutive patients who underwent pancreaticoduodenectomy, ... of the patients, and the incidence of CR-POPF was 32.7% (64/196). ... there were three postoperative deaths, giving a total mortality rate of ...

Risk factors and outcomes of postoperative pancreatic fistula after ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: SJ Fu - 2015 - 被引用次数: 8 - 相关文章

2015年3月26日 - PF was found in 65 (12.2%) cases, of whom 11 were classified into ISGPF grade A, ... Univariate and multivariate analysis showed that blood loss \geq 500 ml, ... Keywords: Pancreatic fistula, Pancreaticoduodenectomy, Risk factor, Outcome. Go to: ... A total of 303 (57.0%, 303/532) patients didn't develop any ...

Pancreatic fistula after pancreatectomy: Evolving definitions ... - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: SV Shrikhande - 2008 - 被引用次数: 67 - 相关文章

2008年10月14日 - Keywords: Pancreatic fistula, Pancreaticoduodenectomy, Pancreatic ... study concluded that post-operative intra-abdominal collections after PD were PF. Total pancreatectomy should not be performed in most cases of 2005;92:539-546. ... risk factors for surgical failure in 302 consecutive cases.

Pancreatoduodenectomy: Risk Factors of Postoperative Pancreatic ...

www.ncbi.nlm.nih.gov/pubmed/26158175 - 翻译此页

作者: LG De Carlis - 2014 - 被引用次数: 1 - 相关文章

Pancreatoduodenectomy: Risk Factors of Postoperative Pancreatic Fistula. ... present study is the analysis of risk factors of postoperative pancreatic fistula (POPF) and of ... January 2003 to July 2013 143 patients underwent PD: 138 cases were included and 3 ... Forty % grade A, 13% grade B and