

[全部](#)
[图片](#)
[新闻](#)
[视频](#)
[购物](#)
[更多 ▾](#)
[搜索工具](#)

找到约 143,000 条结果 (用时 0.75 秒)

## Google 学术: Hepatocytes isolation from resected tissues with benign disease: results of a 5-year experience

Nonalcoholic fatty liver disease: a spectrum of clinical ... - [Matteoni](#) - 被引用次数: 2831

... of the liver: the need for complete resection - [Läuffer](#) - 被引用次数: 108

... with continuous clamping for major liver resection - [Petrowsky](#) - 被引用次数: 181

## Isolation of Primary Human Hepatocytes from Normal and Diseased ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者: [RH Bhogal](#) - 2011 - 被引用次数: 70 - [相关文章](#)

2011年3月29日 - Here we report our experience isolating human hepatocytes from organ ... donors [5]-[6] or from non-diseased tissue removed at surgical resection for ... liver, normal resected liver tissue and cirrhotic liver disease. ... For the purposes of the study, liver tissue obtained from resections for benign lesions was ...

## [PDF] A simple isolation and cryopreservation method for adult human hepatocytes

<https://www.researchgate.net/file.PostFileLoader.html?...AS%3A308588583...> ▾ [翻译此页](#)

Results: the viability and plating efficiency of hepatocytes after partial hepatectomy using two-step extracorporeal ... go to the healthy liver tissue; warm ischemic time did not ... was washed three times at 50×g for 5 minutes at 4°C to ... benign diseases (hepatic hemangioma and transplantation ... Mean donor age (years).

## Factors affecting hepatocyte isolation, engraftment, and replication in ...

[onlinelibrary.wiley.com/doi/10.1002/lt.22099/pdf](http://onlinelibrary.wiley.com/doi/10.1002/lt.22099/pdf) ▾ [翻译此页](#)

作者: [T Kawahara](#) - 2010 - 被引用次数: 31 - [相关文章](#)

Hepatocytes isolated from older donors (60 years) had lower viability and posttransplant engraftment (P: 0.01) .... Of these samples, 5 liver segment tissues were.

## Most Cited Hepatobiliary & Pancreatic Diseases International Articles ...

[www.journals.elsevier.com/hepatobiliary-and...diseases.../most-cited-articles](http://www.journals.elsevier.com/hepatobiliary-and...diseases.../most-cited-articles) ▾ [翻译此页](#)

RESULTS: Hepatocyte-like cells differentiated from MSCs are a promising cell ... Surgical treatment involves major resections of the liver, pancreas and bile duct, with ... found in 66.7% (24/36) of the primary sites of HCC, but not in benign lesions (4/6). ... The 5-year estimated overall survival rate was

**Name of Journal:** World Journal of Gastroenterology

**ESPS Manuscript NO:** 28716

**Manuscript Type:** ORIGINAL ARTICLE

*Basic Study*

**Hepatocytes isolation from resected benign tissues: Results of a 5-year experience**

Meng FY *et al.* Hepatocytes isolation from resected tissues

Fan-Ying Meng, Li Liu, Jun Liu, Chun-You Li, Jian-Ping Wang, Feng-Hui Yang, Zhi-Shui Chen, Ping Zhou

**Fan-Ying Meng, Jun Liu, Chun-You Li, Jian-Ping Wang, Feng-Hui Yang, Division of Hepato-Biliary-Pancreatic Surgery, Institute of Organ Transplantation, Shandong Provincial Hospital affiliated to Shandong University, Jinan 250021, Shandong Province, China**

**Li Liu, Zhi-Shui Chen, Ping Zhou, Institute of Organ Transplantation, Tongji**

Match Overview

1

Internet 107 words  
crawled on 16-Jun-2016  
[www.wjgnet.com](http://www.wjgnet.com)

3%

2

Internet 100 words  
crawled on 04-Dec-2015  
[office.wjgnet.com](http://office.wjgnet.com)

3%

3

Crossref 40 words  
[Rong, Zhong-Hou, Hong-Yuan Chen, Xin-Xing Wang, Zhi-Yi Wang, Guo-Zhe Xian, Bang-Zhen Ma, Cheng-Kun Qin, a](#)

1%



学术搜索

找到约 4,110 条结果 (用时0.07秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☐ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Biliary cystadenocarcinoma of the liver: the need for complete resection

JM L  uffer, HU Baer, CA Maurer, CH Stoupis... - European journal of ..., 1998 - Elsevier

... Endoscopic retrograde cholangiopancreatography (ERCP) with papillotomy and extraction of multiple calculi from ... by Kawarada and associates[28], which suggests that isolated biliary cystadenocarcinoma ... 3-year (5-year) survival rate (Kaplan-Meier) of all resected cases has ...

被引用次数: 109 相关文章 所有 8 个版本 引用 保存

Disseminated tumor cells in lymph nodes as a determinant for survival in surgically resected non-small-cell lung cancer

B Kubuschok, B Passlick, JR Izbicki... - Journal of clinical ..., 1999 - jco.ascopubs.org

... the cytoplasm of all epithelial cells except epidermal keratinocytes, gastric parietal cells, hepatocytes, thymic cortical ... T, Goya T, Tsuchiya R, et al; Prognosis and survival in resected lung cancer ... JR, Passlick B, et al; Frequency and prognostic significance of isolated tumor cells in ...

被引用次数: 182 相关文章 所有 5 个版本 引用 保存

[HTML] Over-expression of the mitogen-activated protein kinase (MAPK) kinase (MEK)-MAPK in hepatocellular carcinoma: its role in tumor progression and apoptosis

H Huynh, TTT Nguyen... - BMC ..., 2003 - bmcgastroenterol.biomedcentral, ...

... myc, c-fos and c-jun, rho, transforming growth factor- , hepatocyte growth factor ... Four weeks post-transfection, individual clones were isolated, expanded and assayed for MEK1 ... demonstration of phosphorylated MEK1/2 and MAPK in malignant and adjacent benign hepatocytes. ...

被引用次数: 257 相关文章 所有 17 个版本 引用 保存 更多

Simultaneous laparoscopic resection of primary colorectal cancer and associated liver metastases: a systematic review

RM Lupinacci, W Andraus, LBDP Haddad... - Techniques in ..., 2014 - Springer

... Colorectal surgery was considered if the primary tumor was completely resected. ... The choice of incision for specimen extraction showed a huge variability between centers (Table 4). Table ... of intestinal bacteria to the liver transection bed, or impaired hepatocyte regeneration [35 ...

被引用次数: 19 相关文章 所有 10 个版本 引用 保存

of macrophage colony-stimulating factor-1 receptor in peritumoral liver tissue

[HTML] biomedcentral.com

[全部](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 158,000 条结果 (用时 0.58 秒)

## Google 学术 : Hepatocytes isolation from resected benign tissues: Results of a 5-year experience

... of the liver: the need for complete **resection** - **Läuffer** - 被引用次数 : 109

... for survival in surgically **resected** non-small-cell lung ... - **Kubuschok** - 被引用次数 : 182

... carcinoma: its role in **tumor** progression and apoptosis - **Huynh** - 被引用次数 : 257

### Isolation of Primary Human Hepatocytes from Normal and Diseased ...

[www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者 : **RH Bhogal** - 2011 - 被引用次数 : 71 - [相关文章](#)

2011年3月29日 - Here we report our **experience isolating** human **hepatocytes** from ... [5]–[6] or from non-diseased **tissue** removed at surgical **resection** for liver ... protocols have been described [3] the overall **results** are so variable in ... For the purposes of the study, liver **tissue** obtained from **resections** for **benign** lesions was ...

### [PDF] Hepatocytes isolated from neoplastic liver-immunomagnetic purging ...

[www.wjgnet.com/esps/DownloadFile.aspx?Type=Digital...1...](http://www.wjgnet.com/esps/DownloadFile.aspx?Type=Digital...1...) ▾ [翻译此页](#)

作者 : **A Gunasegaram** - 2008 - 被引用次数 : 6 - [相关文章](#)

2008年8月28日 - **resection** for neoplasia could provide a novel source of healthy ... **tumor** cell in 106 **hepatocytes**, representing a maximum **tumor** purging ... **Experience** with HT in .... **Benign** liver lesions. 2 ... 5 min at 4°C to separate viable from dead **hepatocytes**. ..... transplantation on a national basis: **results** of a 2-year.

### Factors affecting hepatocyte isolation, engraftment, and replication in ...

[onlinelibrary.wiley.com/doi/10.1002/lt.22099/pdf](http://onlinelibrary.wiley.com/doi/10.1002/lt.22099/pdf) ▾ [翻译此页](#)

作者 : **T Kawahara** - 2010 - 被引用次数 : 31 - [相关文章](#)

**Hepatocytes isolated** from older donors (160 **years**) had lower viability and ... neic cells; the **result** is a transgenic mouse model ... 5). Of these samples, 5 liver segment **tissues** were recovered from the