

全部 图片 新闻 视频 购物 更多 ▼ 搜索工具

找到约 780 条结果 (用时 0.86 秒)

Extramural venous invasion is a potential imaging predictive ... - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▼ 翻译此页

作者: M Chand - 2014 - 被引用次数: 31 - 相关文章

2013年12月3日 - Extramural venous invasion (EMVI) is a poor prognostic factor in rectal cancer and ... All patients had undergone neoadjuvant CRT and curative surgery. ... As an imaging biomarker in rectal cancer, mrEMVI can be used. ... therapy (CRT) by causing vessel fibrosis which can be detected on MRI (Figure 3).

MRI-detected extramural vascular invasion is an independent ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/25500963 ▼ 翻译此页

作者: B Sohn - 2015 - 被引用次数: 11 - 相关文章

2014年12月13日 - MRI-detected extramural vascular invasion is an independent prognostic factor for synchronous metastasis in patients with rectal cancer. ... could predict synchronous distant metastases in rectal cancer. ... EMVI in a large vessel (≥ 3 mm) had a higher risk for metastasis ... Research Support, Non-U.S. Gov't ...

The prognostic significance of postchemoradiotherapy high-resolution ...

www.ncbi.nlm.nih.gov/pubmed/25243543 ▼ 翻译此页

作者: M Chand - 2015 - 被引用次数: 21 - 相关文章

2015 Mar;261(3):473-9. doi: 10.1097/SLA.0000000000000848. The prognostic significance of postchemoradiotherapy high-resolution MRI and histopathology detected extramural venous invasion in rectal cancer. ... BACKGROUND: EMVI is a prognostic factor in rectal cancer but whether this remains so after CRT ...

^[PDF] (NCCN Guidelines®) Rectal Cancer

<https://www.tri-kobe.org/nccn/guideline/colorectal/english/rectal.pdf> ▼

2014年12月9日 - The NCCN Guidelines® and this illustration may not be reproduced in any form without ... circumstances to determine any patient's care or treatment. ... High-risk features include positive margins, lymphovascular invasion, poorly ... defined, locally advanced rectal cancer: Grupo cancer de recto 3 study.



Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 28620

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Combination of three-gene immunohistochemical panel and magnetic resonance imaging-detected extramural vascular invasion to assess prognosis in non-advanced rectal cancer patients

Xiao-Fu Li, Zheng Jiang, Ying Gao, Chun-Xiang Li, Bao-Zhong Shen

Abstract

AIM

To identify a small, clinically applicable immunohistochemistry (IHC) panel that could be combined with magnetic resonance imaging (MRI)-detected extramural vascular invasion (EMVI) for assessment of prognosis concerning

Match Overview

1	Crossref 41 words Shang, Li, Hui-Juan Liu, Jia-Jie Hao, Yan-Yi Jiang, Feng Shi, Yu Zhang, Yan Cai, Xin Xu, Xue-Mei Jia, Qi-Min Zha	1%
2	Internet 38 words crawled on 20-Jun-2016 jco.ascopubs.org	1%
3	Crossref 17 words Lee, Kyu Sang, Yoonjin Kwak, Kyung Han Nam, Duck-Woo Kim, Sung-Bum Kang, Gheeyoung Choe, Woo Ho Ki	1%
4	Internet 15 words crawled on 15-Mar-2016 connection.ebscohost.com	1%
5	Crossref 15 words Y Nancy You. "Is the Increasing Rate of Local Excision ... r Stage I Rectal Cancer in the United States Justified?",	1%
6	Crossref 15 words Jiang, Z.. "Down-regulated GAS1 expression correlates... with recurrence in stage II and III colorectal cancer", Hu	1%
7	Crossref 14 words How, Peter, and Gina Brown. "Magnetic Resonance Im... with Recurrence of Rectal Cancer: A Meta-Analysis of Total Mesorectal	<1%

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 1,010 条结果 (用时 0.85 秒)

Prognostic significance of magnetic resonance imaging-detected ...

www.ncbi.nlm.nih.gov/pubmed/17932879 ▾ [翻译此页](#)

作者: NJ Smith - 2008 - 被引用次数: 141 - [相关文章](#)

Prognostic significance of magnetic resonance imaging-detected extramural vascular invasion in rectal cancer. ... BACKGROUND: Extramural vascular invasion (EMVI) is a poor prognostic feature in colorectal cancer. ... METHODS: Preoperative magnetic resonance images from patients diagnosed with rectal and sigmoid ...

MRI-detected extramural vascular invasion is an independent ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/25500963 ▾ [翻译此页](#)

作者: B Sohn - 2015 - 被引用次数: 12 - [相关文章](#)

2014年12月13日 - MRI-detected extramural vascular invasion is an independent ... factor for synchronous metastasis in patients with rectal cancer. ... OBJECTIVES: To determine whether magnetic resonance imaging ... EMVI in a large vessel (≥ 3 mm) had a higher risk for metastasis than ... Research Support, Non-U.S. Gov't ...

Optimal Imaging Strategies for Rectal Cancer Staging and Ongoing ...

www.ncbi.nlm.nih.gov > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者: S Balyasnikova - 2016 - [相关文章](#)

2016年6月2日 - Imaging determines the optimal treatment for rectal cancer patients. ... MRI enables accurate staging of both early and advanced rectal cancer, ... Combination MRI and CT is essential for distant spread assessment and ... resection margin (CRM) [3] and MR presence of extramural vascular invasion (EMVI) ...

全部 新闻 图片 视频 购物 更多 ▾ 搜索工具

找到约 1,020 条结果 (用时 0.75 秒)

Prognostic significance of magnetic resonance imaging-detected ...

www.ncbi.nlm.nih.gov/pubmed/17932879 ▾ 翻译此页

作者: NJ Smith - 2008 - 被引用次数: 141 - 相关文章

Prognostic significance of magnetic resonance imaging-detected extramural vascular invasion in rectal cancer. ... BACKGROUND: Extramural vascular invasion (EMVI) is a poor prognostic feature in colorectal cancer. ... METHODS: Preoperative magnetic resonance images from patients diagnosed with rectal and sigmoid ...

MRI-detected extramural vascular invasion is an independent ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/25500963 ▾ 翻译此页

作者: B Sohn - 2013 - 被引用次数: 12 - 相关文章

2014年12月13日 - MRI-detected extramural vascular invasion is an independent ... factor for synchronous metastasis in patients with rectal cancer. ... OBJECTIVES: To determine whether magnetic resonance imaging ... EMVI in a large vessel (≥ 3 mm) had a higher risk for metastasis than ... Research Support, Non-U.S. Gov't ...

Optimal Imaging Strategies for Rectal Cancer Staging and Ongoing ...

www.ncbi.nlm.nih.gov ▸ NCBI ▸ Literature ▸ PubMed Central (PMC) - 翻译此页

作者: S Balyasnikova - 2016 - 相关文章

2016年6月2日 - Imaging determines the optimal treatment for rectal cancer patients. ... MRI enables accurate staging of both early and advanced rectal cancer, ... Combination MRI and CT is essential for distant spread assessment and ... resection margin (CRM) [3] and MR presence of extramural vascular invasion (EMVI) ...

[PDF] (NCCN Guidelines®) Rectal Cancer

<https://www.tri-kobe.org/nccn/guideline/colorectal/english/rectal.pdf> ▾

2014年12月9日 - The NCCN Guidelines® and this illustration may not be reproduced in any form without the ... circumstances to determine any patient's care or treatment. concomitant chemoradiotherapy and surgery in magnetic resonance imaging- defined, locally advanced rectal cancer: Grupo cancer de recto 3 study.