

Progression from Low-grade dysplasia to malignancy in patients with barrett's es

全部

图片

视频

新闻

购物

地图

图书

找到约 519,000 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[\[PDF\] Low-Grade Dysplasia in Patients with Barrett's Esophagus ... - ASGE](#)

www.asge.org/WorkArea/DownloadAsset.aspx?id=13294

dysplasia. Pathology slides demonstrating LGD were confirmed by expert external review. At a follow-up office visit, the **patient** is informed about the **diagnosis** of LGD. ... likelihood that **most** will advance to **cancer** is far less than the chance that a ... **Low grade dysplasia in Barrett's esophagus** has a high risk of **progression**.

[The diagnosis of low-grade dysplasia in Barrett's esophagus and its ...](#)

www.ncbi.nlm.nih.gov/pubmed/11151865

OBJECTIVE: The reported risk of **progression from low-grade dysplasia** (LGD) to high-grade **dysplasia** (HGD) or **carcinoma** (CA) in **Barrett's esophagus** varies. biopsy specimens from 43 **patients with Barrett's esophagus diagnosed** and ... RESULTS: Agreement between **two GI pathologists** for a **diagnosis** of LGD was fair ...

[Barrett's esophagus: A review of diagnostic criteria, clinical ... - NCBI](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC3418534/

Clinical follow-up of **patients** with high grade **dysplasia** should be tailored to the ... risk of **malignant progression**, with a reported annual risk of **esophageal** ... are taken along every 2 cm of the BE type mucosa and submitted to **pathology** in ... **Low grade dysplasia** – The **most** important feature of **low grade dysplasia** is ...

[Barrett's esophagus: Incidence, etiology, pathophysiology - NCBI](#)



28688-Review

BY HARSHA MOOLE

Quotes Excluded
Bibliography Excluded8%
SIMILAR

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 28688

Manuscript Type: META-ANALYSIS

Progression from Low-grade dysplasia to malignancy in patients with barrett's esophagus diagnosed by two or more pathologists

Harsha Moole, Jaymon Patel, Zohair Ahmed, Abhiram Duvvuri, Sreekar Vennelaganti, Vishnu Moole, Sowmya Dharmapuri, Raghuveer Boddireddy, Pratyusha Yedama, Naveen Bondalapati, Achuta Uppu, Prashanth Vennelaganti, Srinivas Puli

Match Overview

1	Internet 96 words crawled on 18-Nov-2015 www.ncbi.nlm.nih.gov	2%
2	Crossref 80 words Moole, Harsha, Matthew Bechtold, and Srinivas R. Puli. "Efficacy of preoperative biliary drainage in malignant o...	1%
3	Crossref 51 words Sharma, P.. "Low-grade dysplasia in Barrett's esophagus", <i>Gastroenterology</i> , 200410	1%
4	Internet 38 words crawled on 08-Jun-2012 blue.regence.com	1%
5	Crossref 33 words K K Basu. "Audit of a Barrett's epithelium surveillance t... tabase", <i>European Journal of Gastroenterology & Hepat</i>	1%
6	Crossref 27 words Picardo, S. L., M. P. O'Brien, R. Feighery, D. O'Toole, N. Ravi, N. J. O'Farrell, J. N. O'Sullivan, and J. V. Reynolds. "A	<1%

Progression from Low-grade dysplasia to malignancy in patients with barrett's es

全部

图片

视频

新闻

购物

地图

图书

找到约 519,000 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[\[PDF\] Low-Grade Dysplasia in Patients with Barrett's Esophagus ... - ASGE](#)

www.asge.org/WorkArea/DownloadAsset.aspx?id=13294

dysplasia. Pathology slides demonstrating LGD were confirmed by expert external review. At a follow-up office visit, the **patient** is informed about the **diagnosis** of LGD. ... likelihood that **most** will advance to **cancer** is far less than the chance that a ... **Low grade dysplasia in Barrett's esophagus** has a high risk of **progression**.

[The diagnosis of low-grade dysplasia in Barrett's esophagus and its ...](#)

www.ncbi.nlm.nih.gov/pubmed/11151865

OBJECTIVE: The reported risk of **progression from low-grade dysplasia** (LGD) to high-grade **dysplasia** (HGD) or **carcinoma** (CA) in **Barrett's esophagus** varies. biopsy specimens from 43 **patients with Barrett's esophagus diagnosed** and ... RESULTS: Agreement between **two GI pathologists** for a **diagnosis** of LGD was fair ...

[Barrett's esophagus: A review of diagnostic criteria, clinical ... - NCBI](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC3418534/

Clinical follow-up of **patients** with high grade **dysplasia** should be tailored to the ... risk of **malignant progression**, with a reported annual risk of **esophageal** ... are taken along every 2 cm of the BE type mucosa and submitted to **pathology** in ... **Low grade dysplasia** – The **most** important feature of **low grade dysplasia** is ...

[Barrett's esophagus: Incidence, etiology, pathophysiology - NCBI](#)

[全部](#)[视频](#)[新闻](#)[图片](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 446,000 条结果 (用时 0.91 秒)

您是不是要找：[Progression from Low-grade dysplasia *and* malignancy in patients with barrett's esophagus diagnosed by two or more pathologists](#)

[\[PDF\] Low-Grade Dysplasia in Patients with Barrett's Esophagus ... - ASGE](#)

www.asge.org/WorkArea/DownloadAsset.aspx?id=13294 ▾ [翻译此页](#)

[dysplasia](#), [Pathology](#) slides demonstrating LGD were confirmed by expert external review. At a follow-up office visit, the [patient](#) is informed about the [diagnosis](#) of LGD. ..., likelihood that [most](#) will advance to [cancer](#) is far less than the chance that a ... [Low grade dysplasia in Barrett's esophagus](#) has a high risk of [progression](#).

[The diagnosis of low-grade dysplasia in Barrett's esophagus and its ...](#)

www.ncbi.nlm.nih.gov/pubmed/11151865 ▾ [翻译此页](#)

作者：M Skacel - 2000 - 被引用次数：351 - 相关文章

OBJECTIVE: The reported risk of [progression from low-grade dysplasia](#) (LGD) to high-grade [dysplasia](#) (HGD) or [carcinoma](#) (CA) in [Barrett's esophagus](#) varies. ... biopsy specimens from 43 [patients with Barrett's esophagus diagnosed](#) and ... RESULTS: Agreement between [two](#) GI [pathologists](#) for a [diagnosis](#) of LGD was fair ...

[Barrett's esophagus: A review of diagnostic criteria, clinical ... - NCBI](#)

www.ncbi.nlm.nih.gov ▸ [NCBI](#) ▸ [Literature](#) ▸ [PubMed Central \(PMC\)](#) ▾ [翻译此页](#)

作者：CL Booth - 2012 - 被引用次数：15 - 相关文章

Clinical follow-up of [patients](#) with high grade [dysplasia](#) should be tailored to the ... risk of [malignant progression](#), with a reported annual risk of [esophageal](#) are taken along every 2 cm of the BE type mucosa and submitted to [pathology](#) in ... [Low grade dysplasia](#) – The [most](#) important feature of [low grade dysplasia](#) is ...

[Barrett's esophagus: Incidence, etiology, pathophysiology, prevention ...](#)