



Hepatic non-parenchymal cells; master regulators of alcoholic liver dise



网页

图片

新闻

视频

购物

更多

搜索工具

找到约 87,100 条结果 (用时 0.81 秒)

Google 学术: Hepatic non-parenchymal cells; master regulators of alcoholic liver disease

Lipid peroxidation, stellate cell activation and hepatic ... - George - 被引用次数: 177

... -related parameters in non-alcoholic fatty liver disease ... - Malaguamera - 被引用次数: 102

... of non-alcoholic steatohepatitis (NASH) liver to ... - Serviddio - 被引用次数: 107

Hypoxia and Hypoxia Inducible Factors: Diverse Roles in Liver

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) 翻译此页

作者: B Nath - 2012 - 被引用次数: 66 - 相关文章

In alcoholic liver disease (ALD) and non-alcoholic steatohepatitis (NASH), activation of ... regulation through the inflammatory master regulator group of proteins NFκB. (11)

Exposure of the hepatic stellate cell line LX-2 to hypoxia stimulated in liver parenchymal and nonparenchymal cells; implications for immunity.

Pathogenesis of alcoholic liver disease: Role of oxidative ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) 翻译此页

作者: E Ceni - 2014 - 被引用次数: 6 - 相关文章

2014年12月21日 - Alcoholic liver disease (ALD) is one of the major cause of morbidity and mortality HIFs are the master regulators of oxygen homeostasis and regulate the

... This process is associated with an upheaval of hepatic architecture, stimulate parenchymal and non-parenchymal cells to produce IL-8, ...

Hepatic actions of vitamin D receptor ligands: a sunshine ...

search.proquest.com/openview/.../1?pq-origsite=gscholar 翻译此页

... as a master regulator of skeletal health but also suggest potential therapeutic effects

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 20946

Manuscript Type: TOPIC HIGHLIGHT

2015 Advances in *Alcoholic liver disease*

Hepatic non-parenchymal cells; master regulators of alcoholic liver disease?

Seo W *et al.* Non-parenchymal cells in alcoholic liver disease

Wonhyo Seo, Won-Il Jeong

Abstract

Chronic alcohol consumption is one of the most common causes of the progression of alcoholic liver disease (ALD). In the past, alcohol-mediated hepatocyte injury was assumed to be a significantly major cause of ALD. However, a huge number of

Match Overview

| | | |
|---|--|-----|
| 1 | Internet 221 words crawled on 10-Nov-2011 www.ncbi.nlm.nih.gov | 3% |
| 2 | CrossCheck 31 words G. Sarphie, "Dose- and Time-Dependent Effects of Ethanol on Functional and Structural Aspects of the Liver Si ... | <1% |
| 3 | Internet 29 words crawled on 09-Apr-2015 www.wjgnet.com | <1% |
| 4 | CrossCheck 24 words Fu, Ling, "Interplay of neuropilin-1 and semaphorin 3A ... ter partial hepatectomy in rats", <i>World Journal of Gastroe</i> | <1% |
| 5 | Internet 23 words crawled on 04-Jul-2010 web.kaist.ac.kr | <1% |
| 6 | Internet 19 words crawled on 30-Sep-2015 www.intechopen.com | <1% |
| 7 | CrossCheck 18 words Bertola, Adeline, Stephanie Mathews, Sung Hwan Ki, Hu a-Wang, and Bin Cao, "Mouse model of chronic and hi | <1% |

[网页](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 185,000 条结果 (用时 0.57 秒)

Google 学术: Hepatic non-parenchymal cells; master regulators of alcoholic liver disease?

Lipid peroxidation, stellate cell activation and hepatic ... - George - 被引用次数: 178

... -related parameters in non-alcoholic fatty liver disease ... - Malaguamera - 被引用次数: 102

... of non-alcoholic steatohepatitis (NASH) liver to ... - Serviddio - 被引用次数: 115

An essential role for MCP-1 in alcoholic liver injury ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: P Mandrekar - 2011 - 被引用次数: 72 - [相关文章](#)

Alcoholic liver disease (ALD) is a major health concern and about ninety percent ... In general, MCP-1 seems to play an important role in hepatic inflammatory ... in non-parenchymal cells of alcohol fed WT and MCP-1KO mice (Suppl Fig S3). ... as a master regulator of monocyte/macrophage function has been proposed in ...

Hypoxia and Hypoxia Inducible Factors: Diverse Roles in ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者: B Nath - 2012 - 被引用次数: 81 - [相关文章](#)

We describe regulation of the hypoxia inducible factors and review ... factors that act as master regulators of a homeostatic transcriptional response to hypoxia in virtually ... effects on the function of hepatic parenchymal and nonparenchymal cells. ... In alcoholic liver disease (ALD) and non-alcoholic steatohepatitis (NASH), ...

[全部](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 198,000 条结果 (用时 0.56 秒)

Google 学术 : Hepatic non-parenchymal cells; master regulators of alcoholic liver disease?

Lipid peroxidation, stellate cell activation and hepatic ... - George - 被引用次数 : 178

... -related parameters in non-alcoholic fatty liver disease ... - Malaguarnera - 被引用次数 : 103

... of non-alcoholic steatohepatitis (NASH) liver to ... - Serviddio - 被引用次数 : 114

An essential role for MCP-1 in alcoholic liver injury ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : P Mandrekar - 2011 - 被引用次数 : 72 - [相关文章](#)

Alcoholic liver disease (ALD) is a major health concern and about ninety percent ... In general, MCP-1 seems to play an important role in hepatic inflammatory ... in non-parenchymal cells of alcohol fed WT and MCP-1KO mice (Suppl Fig S3). as a master regulator of monocyte/macrophage function has been proposed in ...

NF-κB in the liver—linking injury, fibrosis and hepatocellular ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : T Luedde - 2011 - 被引用次数 : 259 - [相关文章](#)

Hepatic cirrhosis and hepatocellular carcinoma (HCC) are the most common causes ... factor nuclear factor-κB (NF-κB), a master regulator of inflammation and cell death, ... part of the hepatic wound-healing response to injury induced by alcohol, hepatitis ... Activation of NF-κB in non-parenchymal cells generally promotes ...

Hypoxia and Hypoxia Inducible Factors: Diverse Roles in ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : B Nath - 2012 - 被引用次数 : 82 - [相关文章](#)