

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 21,800 条结果 (用时 0.84 秒)

Google 学术: Establishment of various biliary tract carcinoma cell lines and xenograft models for appropriate preclinical studies

Targeting the Hedgehog pathway in cancer - Rubin - 被引用次数: 600

... of pancreatic adenocarcinoma derived from cell lines ... - Loukopoulos - 被引用次数: 93

... model of ductal adenocarcinoma of the pancreas in ... - Alves - 被引用次数: 45

Establishment of six new human biliary tract carcinoma cell lines and ...

onlinelibrary.wiley.com/doi/10.1111/j.1349-7006.2009.01462.x/pdf - 翻译此页

作者: H Ojima - 2010 - 被引用次数: 12 - 相关文章

The aim of this study was to establish new biliary tract carcinoma (BTC) cell lines ... study, we successfully prepared 19 xenograft models from surgically resected BTC ... BTC cell lines in response to various doses of gemcitabine are shown in Figure 2. ... on the cell lines for both basic and preclinical research. To add.

A novel cell line and xenograft model of ampulla of Vater ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/14677066 - 翻译此页

作者: C Sorio - 2004 - 被引用次数: 8 - 相关文章

2003年12月16日 - A novel cell line and xenograft model of ampulla of Vater adenocarcinoma. ... studies and clarify the relationship between AVC and pancreatic and biliary cancers. ... We report the establishment of a novel AVC cell line (AVC1) derived from a ... mice, making it a suitable model for pre-

Name of journal: World Journal of Gastroenterology

ESPS manuscript NO: 28755

Manuscript Type: LETTER TO THE EDITOR

**Establishment of various biliary tract carcinoma cell lines and xenograft models
for appropriate preclinical studies**

Hidenori Ojima, Seri Yamagishi, Kazuaki Shimada, Tatsuhiro Shibata

Match Overview

1 **Crossref** 29 words
Hidenori Ojima. "Establishment of six new human biliary tract carcinoma cell lines and identification of MAGEH1 as ..."

4%

[全部](#) [图片](#) [视频](#) [新闻](#) [购物](#) [地图](#) [图书](#)

找到约 22,400 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Establishment of various biliary tract carcinoma cell lines and xenograft models for appropriate preclinical studies](#)

[Targeting the Hedgehog pathway in cancer](#) - Rubin - 被引用次数: 605

[... of pancreatic adenocarcinoma derived from cell lines ...](#) - Loukopoulos - 被引用次数: 94

[... model of ductal adenocarcinoma of the pancreas in ...](#) - Alves - 被引用次数: 47

[Establishment and characterization of a human intrahepatic ...](#) - NCBI

www.ncbi.nlm.nih.gov/pmc/articles/PMC4844644/

20 Oct 2015 ... **Cancer cell lines** are affordable **models**, reflecting the ... derived from cholangiocytes in **different** tracts of **biliary tree**. ... In this work, we describe a human Italian ICC **cell line** obtained from a patient-derived **xenograft** (PDX) [33]. This **cell line** could provide a new suitable **model** for **preclinical studies** of ...

[Establishment of a patient-derived intrahepatic cholangiocarcinoma ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4750214/

11 Feb 2016 ... **Cell lines** and animal **models**, even partially reflecting **tumor** characteristics, ... Cholangiocarcinoma (CCA) is the most common **biliary tract** ... to provide a more suitable **model** for **preclinical studies** or to test drug efficacy. Go to: Methods. **Establishment** and characterization of patient derived **xenograft** (PDX).

[PDF] [Establishment of six new human biliary tract carcinoma cell lines ...](#)



[全部](#) [图片](#) [新闻](#) [购物](#) [地图](#) [更多 ▾](#) [搜索工具](#)

找到约 22,000 条结果 (用时 0.83 秒)

Google 学术 : Establishment of various biliary tract carcinoma cell lines and xenograft models for appropriate preclinical studies

Targeting the Hedgehog pathway in cancer - Rubin - 被引用次数 : 605

... of pancreatic adenocarcinoma derived from cell lines ... - Loukopoulos - 被引用次数 : 95

... model of ductal adenocarcinoma of the pancreas in ... - Alves - 被引用次数 : 47

Establishment and characterization of a human intrahepatic ... - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者 : G Cavalloni - 2016 - 被引用次数 : 1 - [相关文章](#)

2015年10月20日 - Cancer cell lines are affordable models, reflecting the ... derived from cholangiocytes in different tracts of biliary tree. ... In this work, we describe a human Italian ICC cell line obtained from a patient-derived xenograft (PDX) [33]. This cell line could provide a new suitable model for preclinical studies of ...

Establishment of a patient-derived intrahepatic cholangiocarcinoma ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者 : G Cavalloni - 2016 - [相关文章](#)

2016年2月11日 - Cell lines and animal models, even partially reflecting tumor characteristics, ... Cholangiocarcinoma (CCA) is the most common biliary tract ... to provide a more suitable model for preclinical studies or to test drug efficacy. Go to: Methods. Establishment and characterization of patient derived xenograft (PDX).

Establishment of six new human biliary tract carcinoma cell lines and ...

onlinelibrary.wiley.com/doi/10.1111/j.1349-7006.2009.01462.x/pdf - [翻译此页](#)

作者 : H Ojima - 2010 - 被引用次数 : 12 - [相关文章](#)

The aim of this study was to establish new biliary tract carcinoma. (BTC) cell lines ... study, we successfully prepared 19 xenograft models from surgically resected BTC BTC cell lines in