

[全部](#)[图片](#)[新闻](#)[视频](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 13,200 条结果 (用时 0.55 秒)

Google 学术: Th17 involvement in NAFLD progression to NASH.

Interleukin - 17 exacerbates hepatic steatosis and ... - Tang - 被引用次数: 104

IL - 17 signaling accelerates the progression of ... - Harley - 被引用次数: 39

... alleviates steatosis and the progression of NASH ... - Liu - 被引用次数: 10

Interleukin-17 exacerbates hepatic steatosis and inflammation ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/21985374 ▾ 翻译此页

作者: Y Tang - 2011 - 被引用次数: 104 - 相关文章

Regulatory T cells (T(regs)) play a critical role in regulating inflammatory ... this study focused on characterizing the role of Th17 cells using a NAFLD mouse model. ... of preventing progression to NASH and advanced liver diseases in NAFLD ...

Interleukin-17 exacerbates hepatic steatosis and inflammation ... - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: Y Tang - 2011 - 被引用次数: 104 - 相关文章

Th17 cells and IL-17 were associated with hepatic steatosis and ... as a means of preventing progression to NASH and advanced liver diseases in NAFLD patients. ... Th17 cells played a key role in NAFLD-associated inflammatory processes.

IL-17 Axis Driven Inflammation in Non-Alcoholic Fatty Liver Disease ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: DA Giles - 2015 - 被引用次数: 5 - 相关文章

2016年7月1日 - NASH prevalence is lower, affecting up to 30% of adults and children with Thus, the

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 27822

Manuscript Type: REVIEW

Th17 involvement in nonalcoholic fatty liver disease progression to non-alcoholic steatohepatitis

Chackelevicius CM *et al.* Th17 cells

Carla Melisa Chackelevicius, Sabrina Eliana Gambaro, Claudio Tiribelli, Natalia Rosso

Abstract

The nonalcoholic fatty liver disease (NAFLD) is the hepatic manifestation of the metabolic syndrome. NAFLD encompasses a wide histological spectrum ranging from benign simple steatosis to non-alcoholic steatohepatitis (NASH). Sustained

Match Overview

1	Internet 125 words crawled on 05-Jul-2016 www.ncbi.nlm.nih.gov	4%
2	Internet 33 words crawled on 04-Jul-2016 www.wjgnet.com	1%
3	Internet 29 words crawled on 12-May-2016 journals.plos.org	1%
4	Crossref 26 words Zhang, Shuai, Di Huang, Jiefeng Weng, Yu Huang, Shuo Liu, Qiang Zhang, Ning Li, Minjie Wen, Guanghui Zhu, F	1%
5	Internet 19 words crawled on 01-Sep-2016 www.nature.com	1%
6	Crossref 17 words Gaffen, S.L. "An overview of IL-17 function and signaling", Cytokine, 200809	1%
7	Crossref 16 words Seo, Nam Kyoung, Hyung suk Koo, Ji-hee Haam, Hyung	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 18,400 条结果 (用时 0.78 秒)

Google 学术: Th17 involvement in nonalcoholic fatty liver disease progression to non-alcoholic steatohepatitis

... steatosis and inflammation in non - alcoholic fatty liver ... - Tang - 被引用次数: 104

... against alcoholic and nonalcoholic fatty liver disease - Wan - 被引用次数: 103

Role of IL-17 and Th17 cells in liver diseases - Hammerich - 被引用次数: 137

Interleukin-17 exacerbates hepatic steatosis and inflammation in non ...

www.ncbi.nlm.nih.gov/pubmed/21985374 ▾ 翻译此页

作者: Y Tang - 2011 - 被引用次数: 104 - 相关文章

... the progression of simple steatosis to non-alcoholic fatty liver disease (NAFLD) ... study focused on characterizing the role of Th17 cells using a NAFLD mouse ...

Interleukin-17 exacerbates hepatic steatosis and inflammation in non ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者: Y Tang - 2011 - 被引用次数: 104 - 相关文章

Th17 cells and IL-17 were associated with hepatic steatosis and ... of preventing progression to NASH and advanced liver diseases in NAFLD ... Th17 cells played a key role in NAFLD-associated inflammatory processes. Dietary factors alter hepatic innate immune system in mice with nonalcoholic fatty liver disease.

Progression from Nonalcoholic Fatty Liver to Nonalcoholic ...

www.jimmunol.org/content/196/1/97.abstract ▾ 翻译此页

作者: M Rau - 2016 - 被引用次数: 2 - 相关文章

全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 18,600 条结果 (用时 0.78 秒)

Google 学术 : Th17 involvement in nonalcoholic fatty liver disease progression to non-alcoholic steatohepatitis

... steatosis and inflammation in non-alcoholic fatty liver ... - Tang - 被引用次数 : 104

... against alcoholic and nonalcoholic fatty liver disease - Wan - 被引用次数 : 104

Role of IL-17 and Th17 cells in liver diseases - Hammerich - 被引用次数 : 138

Interleukin-17 exacerbates hepatic steatosis and inflammation in non ...

www.ncbi.nlm.nih.gov/pubmed/21985374 ▾ 翻译此页

作者 : Y Tang - 2011 - 被引用次数 : 104 - 相关文章

... the progression of simple steatosis to non-alcoholic fatty liver disease (NAFLD) ... study focused on characterizing the role of Th17 cells using a NAFLD mouse ...

Interleukin-17 exacerbates hepatic steatosis and inflammation in non ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ 翻译此页

作者 : Y Tang - 2011 - 被引用次数 : 104 - 相关文章

Th17 cells and IL-17 were associated with hepatic steatosis and ... of preventing progression to NASH and advanced liver diseases in NAFLD ... Th17 cells played a key role in NAFLD-associated inflammatory processes. Dietary factors alter hepatic innate immune system in mice with nonalcoholic fatty liver disease.

Progression from Nonalcoholic Fatty Liver to Nonalcoholic ...

www.jimmunol.org/content/196/1/97.abstract ▾ 翻译此页

作者 : M Rau - 2016 - 被引用次数 : 2 - 相关文章

2016年1月1日 - Nonalcoholic fatty liver disease is increasing in prevalence. It can be subdivided into nonalcoholic fatty liver (NAFL) and nonalcoholic steatohepatitis (NASH). ... Increased hepatic Th17 cells and IL-17 expression were observed in NASH mice ... NAFLD: nonalcoholic fatty liver disease; NASH: nonalcoholic ...

Non-Alcoholic Steatohepatitis: From Pathophysiology to ... - Karger

<https://www.karger.com/Article/Pdf/444547> - 翻译此页