

Reoperation in an adult female with 'right-sided' Hirschsprung's disease com

全部

图片

视频

新闻

购物

地图

图书

找到约 1,270 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Case Report: Surgical management of pheochromocytoma in a 13 ...](#)

www.ncbi.nlm.nih.gov/pmc/articles/PMC4039882/

21 May 2014 ... In this **case report** PCC was detected in the second trimester, and managed successfully. ... clinic because of **hypertension refractory** to medical therapy and **right** adrenal mass. A brief **literature review** showed that patients with PCC during ... Endocrine **disorders** in pregnancy: pheochromocytoma and ...

[\[PDF\] Supplementum 203 - Swiss Medical Weekly](#)

www.smw.ch/fileadmin/smw/pdf/Supplementa.../SMW_Suppl-203.pdf

26 May 2014 ... Official journal of the Swiss Society of Infectious **Diseases**, and 252 **adults** (163 **female**, Mage = 23.20, SDage = 2.68) from the However, data on the natural **history** of **cough** only approximately 40 **cases** are reported in the English **literature**. and she is slowly regaining **right** ankle function.

[Minoxidil experience in Australia 1974-1980. - Abstract - Europe PMC](#)

europepmc.org/abstract/med/7254111

Search worldwide, life-sciences **literature**. **Search**. Advanced ... The most common **side** effects were fluid retention and hypertrichosis. In 71% of patients the ...

[Gandhi Medical College - Contributions - ResearchGate](#)

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript No.: 28127

Manuscript Type: CASE REPORT

Reoperation in an adult female with "right-sided" Hirschsprung's disease complicated by refractory hypertension and cough

Zhi-Jian Wei, Lei Huang, A-Man Xu

Abstract

Hirschsprung's disease (HD) is an intestinal malformation caused by the innate

Match Overview

1	Internet 44 words crawled on 21-Oct-2013 www.science.gov	1%
2	Internet 19 words crawled on 15-Mar-2016 www.wjgnet.com	1%
3	Internet 14 words crawled on 15-Apr-2016 www.surgery.uthscsa.edu	<1%

Reoperation in an adult female with "right-sided" Hirschsprung's disease complic

全部

图片

视频

新闻

购物

地图

图书

找到约 749 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Saudi license exam \(SLE\) 3rd edt. UQU - Scribd](#)

<https://www.scribd.com/doc/.../Saudi-license-exam-SLE-3rd-edt-UQU>

评分: 4.5 - 7 票

Most common cause of secondary hypertension in female adolescent is: a) Cushing In most adults with symptomatic, severe aortic stenosis, aortic valve Patient in ER: dyspnea, right sided chest pain, engorged neck veins and disease but presents the risk of the development of Vancomycin-resistant enterococcus.

[\[PDF\] Abstracts - American Federation for Medical Research](#)

afmr.org/files/Southern/2015/Southern_Regional_Meeting_Abstracts.pdf

26 Feb 2015 ... cortisone to urinary cortisol ratio in resistant hypertension patients Caucasian female in the Adult ICU with a fulminate illness following a ... also reported new onset headache and progressive right-sided ... rhinitis presented with cough, chest pain, and shortness of breath for the need of reoperation.

[Full text of "TAIF UQU 4th Edition \(FINAL REVISION\)"](#)

[https://archive.org/.../TAIF%20____UQU%204th%20edition%20\(FINAL%20REVISION\)_djvu.txt](https://archive.org/.../TAIF%20____UQU%204th%20edition%20(FINAL%20REVISION)_djvu.txt)

Most common cause of secondary hypertension in female adolescent is: a) Cushing The most common cause of cough in adults is a) Asthma b) GERD c) Patient in ER: dyspnea, right sided chest pain, engorged neck veins and weak diagnosis is: a) Midgut volvulus b) Meconium ileus c) Hirschsprung disease 217.

[全部](#)[新闻](#)[图片](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 750 条结果 (用时 1.02 秒)

Saudi license exam (SLE) 3rd edt. UQU - Scribd

<https://www.scribd.com/doc/.../Saudi-license-exam-SLE-3rd-edt-UQU> ▾ [翻译此页](#)

★★★★★ 评分：4.5 - 7 票

Most common cause of secondary **hypertension** in **female** adolescent is: a) Cushing The most common cause of **cough** in **adults** is a) Asthma b) GERD c) Patient in ER: dyspnea, **right sided** chest pain, engorged neck veins and weak diagnosis is: a) Midgut volvulus b) Meconium ileus c) **Hirschsprung disease** 195.

[PDF] Abstracts - American Federation for Medical Research

afmr.org/files/Southern/2015/Southern_Regional_Meeting_Abstracts.pdf ▾

2015年2月26日 - cortisone to urinary cortisol ratio in **resistant hypertension** patients Caucasian **female** in the **Adult** ICU with a fulminate illness following a ... also reported new onset headache and progressive **right-sided** ... rhinitis presented with **cough**, chest pain, and shortness of breath for the need of **reoperation**.

Full text of "TAIF UQU 4th Edition (FINAL REVISION)" - Internet Archive

[https://archive.org/.../TAIF%20__UQU%204th%20edition%20\(FINAL%20REVISIO...](https://archive.org/.../TAIF%20__UQU%204th%20edition%20(FINAL%20REVISIO...) ▾

Most common cause of secondary **hypertension** in **female** adolescent is: a) Cushing The most common cause of **cough** in **adults** is a) Asthma b) GERD c) Patient in ER: dyspnea, **right sided** chest pain, engorged neck veins and weak diagnosis is: a) Midgut volvulus b) Meconium ileus c) **Hirschsprung disease** 217.

Full text of "The Mont Reid Surgical Handbook" - Internet Archive

https://archive.org/stream/.../The%20Mont%20Reid%20Surgical%20Handbook_djvu.txt

Chronic illnesses — diabetes mellitus, **hypertension**, coronary artery **disease**, chronic Palpate bulges with the patient standing relaxed, then **coughing**/ straining, feeling for ... Blood at the male penile meatus or **female** urethra, gross hematuria b. For patients with low cardiac output, **right-sided** cardiomegaly, unusual ...