

Low phase angle is associated with the development of hepatic encephalopathy i

全部 图片 视频 新闻 地图 图书

找到约 32,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Nutritional assessment in cirrhotic patients with hepatic ... - NCBI](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4691697/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4691697/>

28 Dec 2015 ... Keywords: **Hepatic encephalopathy**, Liver **cirrhosis**, Malnutrition, In **cirrhosis**, obesity can be **associated** with loss of muscle mass, ... in the assessment of nutritional status in **cirrhotic patients** who **developed** ... in **cirrhotic patients** with HE because **phase angle** and BCM convey prognostic information[72].

[Role of Nutrition in the Management of Hepatic Encephalopathy in ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3017957/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3017957/>

22 Dec 2010 ... Role of Nutrition in the Management of **Hepatic Encephalopathy** in End-Stage Liver Failure ... energy metabolism may **shift** from carbohydrate to fat oxidation [7] while ... **Cirrhotic patients** may unintentionally consume a **low** energy diet, ... Malnourished **patients with cirrhosis** are prone to the **development** of ...

[\[PDF\] Phase angle as a nutritional evaluation tool in all stages of chronic ...](https://scielo.isciii.es/pdf/nh/v27n6/37original26.pdf)

scielo.isciii.es/pdf/nh/v27n6/37original26.pdf

stages of CLD, including chronic hepatitis, liver **cirrhosis** ... **patients** without **hepatic encephalopathy** ... PhA was strongly **associated** with survival and ... tion. 13 **Phase angle** (PhA) is obtained from the direct ... be consistent with **low** Xc or with cell death or a break- ... intake and malnutrition may facilitate the **development** ...

[Nutritional assessment in patients with cirrhosis - SciELO](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0004...)

www.scielo.br/scielo.php?script=sci_arttext&pid=S0004...

CONTEXT: Malnutrition in **cirrhotic patients** with end-stage disease is ... RESULTS: Through **phase angle** of bioelectrical impedance analysis (BIA) method, ... and more than 60% of these **patients** with severe **hepatic** dysfunction(21, 22). The assessment of the **association** between the reference values of HGS and BIA ...

[Hepatic Encephalopathy: Definition, Pathogenesis, Clinical Features ...](#)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 29063

Manuscript Type: ORIGINAL ARTICLE

Prospective Study

Low phase angle is associated with the development of hepatic encephalopathy
in patients with cirrhosis

Astrid Ruiz-Margáin, Ricardo Ulises Macías-Rodríguez, Javier Ampuero, Francisco
Javier Cubero, Luis Chi-Cervera, Silvia L Ríos Torres, Andrés Duarte-Rojo,
Ángeles Espinosa-Cuevas, Manuel Romero-Gómez, Aldo Torre

Abstract

AIM

Evaluate the association between phase angle and the development of hepatic

Match Overview

- | Match Number | Crossref | Words | Match Text | Similarity |
|--------------|----------|----------|--|------------|
| 1 | Crossref | 17 words | Luis Compañy. "Extrapyramidal signs predict the developm
ent of overt hepatic encephalopathy in patients with liver c.... | 1% |
| 2 | Crossref | 16 words | "Poster Session 1: Encephalopathy and Other Complicatio
ns", Hepatology, 2014. | 1% |

Low phase angle is associated with the development of hepatic encephalopathy i

全部 图片 视频 新闻 地图 图书

找到约 32,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Nutritional assessment in cirrhotic patients with hepatic ... - NCBI](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4691697/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4691697/>

28 Dec 2015 ... Keywords: **Hepatic encephalopathy**, Liver **cirrhosis**, Malnutrition, In **cirrhosis**, obesity can be **associated** with loss of muscle mass, ... in the assessment of nutritional status in **cirrhotic patients** who **developed** ... in **cirrhotic patients** with HE because **phase angle** and BCM convey prognostic information[72].

[Role of Nutrition in the Management of Hepatic Encephalopathy in ...](https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3017957/)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3017957/>

22 Dec 2010 ... Role of Nutrition in the Management of **Hepatic Encephalopathy** in End-Stage Liver Failure ... energy metabolism may **shift** from carbohydrate to fat oxidation [7] while ... **Cirrhotic patients** may unintentionally consume a **low** energy diet, ... Malnourished **patients with cirrhosis** are prone to the **development** of ...

[\[PDF\] Phase angle as a nutritional evaluation tool in all stages of chronic ...](https://scielo.isciii.es/pdf/nh/v27n6/37original26.pdf)

scielo.isciii.es/pdf/nh/v27n6/37original26.pdf

stages of CLD, including chronic hepatitis, liver **cirrhosis** ... **patients** without **hepatic encephalopathy** ... PhA was strongly **associated** with survival and ... tion. 13 **Phase angle** (PhA) is obtained from the direct ... be consistent with **low** Xc or with cell death or a break- ... intake and malnutrition may facilitate the **development** ...

[Nutritional assessment in patients with cirrhosis - SciELO](http://www.scielo.br/scielo.php?script=sci_arttext&pid=S0004...)

www.scielo.br/scielo.php?script=sci_arttext&pid=S0004...

CONTEXT: Malnutrition in **cirrhotic patients** with end-stage disease is ... RESULTS: Through **phase angle** of bioelectrical impedance analysis (BIA) method, ... and more than 60% of these **patients** with severe **hepatic** dysfunction(21, 22). The assessment of the **association** between the reference values of HGS and BIA ...

[Hepatic Encephalopathy: Definition, Pathogenesis, Clinical Features ...](#)

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 21,100 条结果 (用时 0.82 秒)

Nutritional assessment in cirrhotic patients with hepatic encephalopathy

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者：FG Romeiro - 2015 - 被引用次数：3 - [相关文章](#)

2015年12月28日 - Keywords: **Hepatic encephalopathy**, Liver **cirrhosis**, Malnutrition, Anthropometry, However, glutamine synthesis and ammonia uptake are **low** during rest[19]. ... and oxidative stress has been **associated** to HE **development**[17]. in **cirrhotic patients** with HE because **phase angle** and BCM convey ...

Role of Nutrition in the Management of Hepatic Encephalopathy in ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者：C Bémour - 2010 - 被引用次数：46 - [相关文章](#)

2010年12月22日 - **Cirrhotic patients** may unintentionally consume a **low** energy diet, an observation ... Malnourished **patients with cirrhosis** are prone to the **development** of ... biased by the presence of edema or ascites **associated** with **liver** failure. during the anhepatic **phase** and to induce an exacerbated proinflammatory ...

Hepatic Encephalopathy: Definition, Pathogenesis, Clinical Features ...

emedicine.medscape.com/article/186101-overview ▾ [翻译此页](#)

2015年12月8日 - **Hepatic encephalopathy** is a syndrome observed in **patients with cirrhosis**. ... The **development of hepatic encephalopathy** negatively impacts patient survival. ... is **associated** with a survival probability of 42% at 1 year of follow-up and ... In contrast, brain edema is rarely reported in **patients with cirrhosis**.

缺少字词：angle

Sleepwake abnormalities in patients with cirrhosis - Wiley Online Library

onlinelibrary.wiley.com/doi/10.1002/hep.26555/pdf ▾ [翻译此页](#)

作者：S Montagnese - 2014 - 被引用次数：33 - [相关文章](#)

A considerable proportion of **patients with cirrhosis** exhibit insomnia, delayed sleep habits ... nation of high/**low** sleep propensity in relation to dark/light cues. ... However, while melatonin abnormalities are