

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多 ▾](#)[搜索工具](#)

找到约 31,100 条结果 (用时 0.44 秒)

Google 学术: Sirtuins and Nonalcoholic Fatty Liver Disease

... with the severity of nonalcoholic fatty liver disease - Min - 被引用次数: 155

... target for treatment of nonalcoholic fatty liver disease - Colak - 被引用次数: 54

... accumulation in non-alcoholic fatty liver disease - Kawano - 被引用次数: 145

Direct evidence of sirtuin downregulation in the liver of non-alcoholic ...

www.ncbi.nlm.nih.gov/pubmed/25361925 ▾ 翻译此页

作者: T Wu - 2014 - 被引用次数: 5 - 相关文章

Ann Clin Lab Sci. 2014 Fall;44(4):410-8. Direct evidence of sirtuin downregulation in the liver of non-alcoholic fatty liver disease patients. Wu T(1), Liu YH(1), ...

SIRT1 as a potential therapeutic target for treatment of nonalcoholic ...

www.ncbi.nlm.nih.gov/pubmed/21525818 ▾ 翻译此页

作者: Y Colak - 2011 - 被引用次数: 54 - 相关文章

There are 7 different sirtuins in mammals (SIRT1-7), of which SIRT1 is the best ... Nonalcoholic fatty liver disease (NAFLD) is currently the most common liver ...

A potential treatment of non-alcoholic fatty liver disease with SIRT1 ...

www.ncbi.nlm.nih.gov/pubmed/25267960 ▾ 翻译此页

作者: Y Colak - 2014 - 被引用次数: 14 - 相关文章

A potential treatment of non-alcoholic fatty liver disease with SIRT1 activators. ... Sirtuins (SIRTs) are members of the silent information regulator-2 family and act ...

The expression of SIRT1 in nonalcoholic fatty liver disease induced by ...

www.ncbi.nlm.nih.gov/pubmed/17498258 ▾ 翻译此页

作者: XQ Deng - 2007 - 被引用次数: 97 - 相关文章

The expression of SIRT1 in nonalcoholic fatty liver disease induced by high-fat diet ...
Liver/ultrastructure; Male; Rats; Rats, Wistar; Sirtuin 1; Sirtuins/metabolism* ...

[PDF] Role of Mitochondria in Nonalcoholic Fatty Liver Disease - MDPI.com

www.mdpi.com/1422-0067/15/5/8713/pdf ▾ 翻译此页

作者: E Nasciaro - 2014 - 被引用次数: 29 - 相关文章

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 29484

Manuscript Type: REVIEW

Sirtuins and nonalcoholic fatty liver disease

Nassir F *et al.* Sirtuins as therapeutic target for NAFLD

Fatiha Nassir, Jamal A Ibdah

Fatiha Nassir, Jamal A Ibdah, Research Service, Harry S Truman Memorial VA Hospital, University of Missouri, Columbia, MO 65201, United States

Fatiha Nassir, Departments of Medicine, Division of Gastroenterology and Hepatology, Nutrition and Exercise Physiology, University of Missouri, Columbia, MO 65212, United States

Jamal A Ibdah, Departments of Medicine, Division of Gastroenterology and

Match Overview

1	Internet 202 words crawled on 30-Oct-2015 www.ncbi.nlm.nih.gov	5%
2	Internet 168 words crawled on 15-Mar-2016 www.wjgnet.com	5%
3	Crossref 86 words Haas, Joel T., Sven Francque, and Bart Staels. "Pathophy... ology and Mechanisms of Nonalcoholic Fatty Liver Disease"	2%
4	Crossref 23 words Hebert, Alexander S., Kristin E. Dittenhafer-Reed, Wei Yu, Derek J. Bailey, Ebru Selin Selen, Melissa D. Boersma, Jos	1%
5	Internet 23 words crawled on 08-Jul-2016 mdpi.com	1%
6	Internet 20 words crawled on 10-Apr-2016 tpp.uah.edu	1%
7	Internet 20 words crawled on 06-Jul-2016 www.linkedin.com	1%

学术搜索

找到约 2,700 条结果 (用时0.08秒)

文章

我的图书馆

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

创建快讯

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

[HTML] Increased hepatic synthesis and dysregulation of cholesterol metabolism is associated with the severity of nonalcoholic fatty liver disease

[HTML] sciencedirect.com

HK Min, A Kapoor, M Fuchs, F Mirshahi, H Zhou... - Cell metabolism, 2012 - Elsevier
... The clinical-histologic phenotype of NAFLD extends from a nonalcoholic fatty liver (NAFL) to nonalcoholic steatohepatitis (NASH ... It inhibits sirtuin-1 which is known to dephosphorylate AMP kinase a known regulator of HMGCR phosphorylation (Beg et al., 1984, Chaudhary ...
被引用次数: 163 相关文章 所有 18 个版本 引用 保存

SIRT1 as a potential therapeutic target for treatment of nonalcoholic fatty liver disease

[PDF] researchgate.net

Y Colak, O Ozturk, E Senates, I Tuncer... - Medical Science ..., 2011 - medscimonit.com
... Currently, the best-studied sirtuin homolog, SIRT1, is expressed in metabolic tissues such as ... T, Heltweg B et al: Substrate-specific activa- tion of sirtuins by resveratrol. ... Effects of calorie restriction on SIRT1 ex- pression in liver of nonalcoholic fatty liver disease: experiment with ...
被引用次数: 56 相关文章 所有 9 个版本 引用 保存

Mechanisms of hepatic triglyceride accumulation in non-alcoholic fatty liver disease

[HTML] springer.com

Y Kawano, DE Cohen - Journal of gastroenterology, 2013 - Springer
... Accumulating evidence also indicates that members of the sirtuin gene family ... Sirtuins control target proteins by deacetylating histones and other transcriptional regulators. ... Fabbrini E, Sullivan S, Klein S. Obesity and nonalcoholic fatty liver disease: biochemical, metabolic, and ...
被引用次数: 151 相关文章 所有 18 个版本 引用 保存

[HTML] Mitochondrial adaptations and dysfunctions in nonalcoholic fatty liver disease

[HTML] wiley.com

K Begriche, J Massart, MA Robin, F Bonnet... - ..., 2013 - Wiley Online Library
... 14,15 Sirtuins also interact with PPARα coactivator 1α (PGC1α), thus favoring mitochondrial ... study, 13 C-KICA decarboxylation was unaffected in patients with nonalcoholic fatty liver, while it ... of MRC and OXPHOS seem to be more consistently observed in fatty liver induced by a ...
被引用次数: 126 相关文章 所有 7 个版本 引用 保存

Resveratrol improves non - alcoholic fatty liver disease by activating AMP -



全部 图片 新闻 视频 购物 更多 ▾ 搜索工具

找到约 80,100 条结果 (用时 0.67 秒)

Google 学术 : Sirtuins and nonalcoholic fatty liver disease

... with the severity of nonalcoholic fatty liver disease - Min - 被引用次数 : 164

... target for treatment of nonalcoholic fatty liver disease - Colak - 被引用次数 : 56

... accumulation in non-alcoholic fatty liver disease - Kawano - 被引用次数 : 154

Direct evidence of sirtuin downregulation in the liver of non-alcoholic ...

<https://www.ncbi.nlm.nih.gov/pubmed/25361925> - 翻译此页

作者 : T Wu - 2014 - 被引用次数 : 5 - 相关文章

Ann Clin Lab Sci. 2014 Fall;44(4):410-8. Direct evidence of sirtuin downregulation in the liver of non-alcoholic fatty liver disease patients. Wu T(1), Liu YH(1), ...

SIRT1 as a potential therapeutic target for treatment of nonalcoholic ...

<https://www.ncbi.nlm.nih.gov/pubmed/21525818> - 翻译此页

作者 : Y Colak - 2011 - 被引用次数 : 56 - 相关文章

There are 7 different sirtuins in mammals (SIRT1-7), of which SIRT1 is the best ... Nonalcoholic fatty liver disease (NAFLD) is currently the most common liver ...

The expression of SIRT1 in nonalcoholic fatty liver disease induced by ...

www.ncbi.nlm.nih.gov/pubmed/17498258 ▾ 翻译此页

作者 : XQ Deng - 2007 - 被引用次数 : 101 - 相关文章

The expression of SIRT1 in nonalcoholic fatty liver disease induced by high-fat diet ... Liver/ultrastructure; Male; Rats; Rats, Wistar; Sirtuin 1; Sirtuins/metabolism* ...

[PDF] Role of Mitochondria in Nonalcoholic Fatty Liver Disease - MDPI.com

www.mdpi.com/1422-0067/15/5/8713/pdf ▾ 翻译此页

作者 : F Nassir - 2014 - 被引用次数 : 36 - 相关文章

2014年5月15日 - sirtuin-1 (SIRT1); sirtuin-3 (SIRT3). 1. Introduction. Nonalcoholic fatty liver disease