

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 27216

Manuscript Type: ORIGINAL ARTICLE

Case Control Study

**Effect of switching from treatment with nucleos(t)ide analogues to
pegylated interferon α -2a on virological and serological response in chronic
hepatitis B patients**

He LT *et al.* Switching to PegIFN α -2a in NA-treated CHB patients

Li-Ting He, Xiao-Guang Ye, Xiao-Yuan Zhou

Li-Ting He, Xiao-Guang Ye, Xiao-Yuan Zhou, Department of Infectious Dis-
eases, The Second Affiliated Hospital of Guangzhou Medical University,
Guangzhou 510260, Guangdong Province, China

Match Overview

1

Internet 218 words
crawled on 13-May-2016
www.wjgnet.com

5%

学术搜索

找到约 249 条结果 (用时0.14秒)

文章

我的图书馆

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2016以来

2015以来

2012以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☐ 创建快讯

[HTML] Peginterferon alpha - 2b plus adefovir induce strong cccDNA decline and HBsAg reduction in patients with chronic hepatitis B

[HTML] wiley.com

..., T Volz, P Buggisch, B Zollner, T Longerich... - ..., 2006 - Wiley Online Library

... As HBV polymerase inhibitors do not have a direct effect on cccDNA formation, the decreased intrahepatic ... Because liver biopsies were taken only at the end of therapy, the impact of combination therapy ... Table 2, Summary of Effects of Peg-IFN and ADV in CH-B Patients After 48 ...

被引用次数: 402 相关文章 所有 10 个版本 引用 保存

[HTML] Peginterferon Alfa-2a, lamivudine, and the combination for HBeAg-positive chronic hepatitis B

[HTML] nejm.org

GKK Lau, T Piratvisuth, KX Luo... - ... England Journal of ..., 2005 - Mass Medical Soc

... alfa-2a, like conventional interferon alfa, has a dual immunomodulatory and antiviral mode of action. ... similar to rates in the overall study population (Table 3Table 3 Effect of Baseline ... with HBeAg-negative chronic hepatitis B, and there were no unexpected adverse effects. 29 The ...

被引用次数: 1534 相关文章 所有 15 个版本 引用 保存

[HTML] Long-term therapy with adefovir dipivoxil for HBeAg-negative chronic hepatitis B

[HTML] nejm.org

SJ Hadziyannis, NC Tassopoulos... - ... England Journal of ..., 2005 - Mass Medical Soc

... improvement in patients with hepatitis B e antigen (HBeAg)-negative chronic hepatitis B. We evaluated the effect of continued ... (2014) Nucleoside/nucleotide analog inhibitors of hepatitis B virus polymerase: mechanism of action and resistance. ... (2013) Effects of ketoconazole ...

被引用次数: 756 相关文章 所有 18 个版本 引用 保存

[HTML] Peginterferon alfa-2a alone, lamivudine alone, and the two in combination in patients with HBeAg-negative chronic hepatitis B

[HTML] nejm.org

..., R Jin, ZM Lu, T Piratvisuth... - ... England Journal of ..., 2004 - Mass Medical Soc

... interferon alfa, has a dual immunomodulatory and antiviral mode of action but has ... Only if the overall test of the treatment effect was significant were pairwise comparisons performed ... in combination with lamivudine were satisfactory, and there were no unexpected adverse effects, ...

被引用次数: 1156 相关文章 所有 10 个版本 引用 保存

Sustained HBeAg and HBsAg loss after long-term follow-up of HBeAg-positive patients treated with peginterferon α-2b

[HTML] theclinics.com

..., J Trojan, F Tabak, TMK So, SV Feinman, T Mach... - Gastroenterology, 2008 - Elsevier

... α-2b was lowered to 50 µg/wk to prevent early discontinuation of therapy due to side effects. ... A preliminary report on sustainability of HBeAg loss after treatment with PEG-IFN α-2a ... or telbivudine, HBeAg seroconversion was observed in 20%, 21% after 2 years of therapy. 16 and

[全部](#)[图片](#)[新闻](#)[购物](#)[地图](#)[更多 ▾](#)[搜索工具](#)

找到约 6,190 条结果 (用时 0.99 秒)

Combination of Pegylated Interferon and Nucleos(t)ide Therapy - NCBI

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: EJ Cho - 2016

2016年7月15日 - See "Potential Efficacy of Pegylated Interferon- α and a Nucleos(t)ide ... Current treatments for chronic hepatitis B virus (HBV) infection based on nucleos(t)ide analogues ... (HBeAg)-positive chronic hepatitis B patients, combination therapy ... Serological response rates, including HBeAg seroconversion and ...

Sequential Combination Therapy with Pegylated Interferon Leads to ...

www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) ▾ [翻译此页](#)

作者: GJ Li - 2015 - 被引用次数: 7 - [相关文章](#)

2015年5月4日 - Nucleos(t)ide analogues rarely result in a durable off-treatment response in chronic hepatitis B infection, whereas pegylated interferon ... We assessed the effect of sequential combination therapy with Peg-IFN- α 2a and entecavir in hepatitis B e ... More patients in the S-C therapy group achieved HBeAg ...

Predictors of response to pegylated interferon in chronic hepatitis B: a ...

www.nature.com > Scientific Reports > Articles - [翻译此页](#)

作者: YC Wang - 2016

2016年7月12日 - The treatment options for chronic hepatitis B include nucleoside or nucleotide ... with a sustained off-treatment response to PEG-IFN α 2a based treatment using a ... Serological, virological and combined responses Durability of Nucleos(t)ide Analogues Treatment in Patients With Chronic Hepatitis B ...

Pegylated-Interferon Alpha Therapy for Treatment-Experienced ...

journals.plos.org/plosone/article?id=10.1371%2Fjournal.pone.0122259 ▾ [翻译此页](#)

作者: ML Yeh - 2015 - 被引用次数: 6 - [相关文章](#)

2015年4月2日 - All of the patients were treated with Peg-IFN α -2a at 180 μ g weekly ... Studies are limited on pegylated interferon (Peg-IFN) therapy for chronic hepatitis B (CHB) patients ... HBeAg-