

全部 新闻 图片 视频 购物 更多 搜索工具



找到约 203,000 条结果 (用时 0.70 秒)

Google 学术: Visceral hypersensitivity in inflammatory bowel diseases and irritable bowel syndrome: the role of proteases

- ... activity in visceral pain in irritable bowel syndrome - Cenac - 被引用次数: 433
- ... visceral hypersensitivity and improves intestinal ... - Klooker - 被引用次数: 248
- ... sensory neurons in irritable bowel syndrome - Barbara - 被引用次数: 505

Role for protease activity in visceral pain in irritable bowel syndrome

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: N Cenac - 2007 - 被引用次数: 432 - 相关文章

2007年2月15日 - Role for protease activity in visceral pain in irritable bowel syndrome ... IBS can also appear in patients with inflammatory bowel disease (IBD) who IBS tissue supernatants cause visceral hypersensitivity in mice: a protease ...

Protease inhibition as new therapeutic strategy for GI diseases ...

gut.bmj.com > Volume 65, Issue 7 - 翻译此页

作者: N Vergnolle - 2016 - 被引用次数: 3 - 相关文章

2016年4月12日 - The therapeutic potential of protease inhibition in GI diseases is parts of the colon of those patients, where no inflammatory cell infiltration was detected. PAR2 activation causes visceral hypersensitivity, modifies intestinal motility Role for protease activity in visceral pain in irritable bowel syndrome.

Irritable bowel syndrome : Nature Reviews Disease Primers

www.nature.com > Nature Reviews Disease Primers > Primers

2016年3月24日 - Irritable bowel syndrome (IBS) is a functional gastrointestinal disease with a high prevalence. ... hyperpermeability, dysbiosis, inflammation, visceral hypersensitivity, ... QOL and the putative roles of inflammation, genetics, the intestinal Increased levels of histamine and proteases in tissue supernatants

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 30250

Manuscript Type: REVIEW

Visceral hypersensitivity in inflammatory bowel diseases and irritable bowel syndrome: the role of proteases

Hannah Ceuleers, Hanne Van Spaendonk, Nikita Hanning, Jelena Heirbaut, Anne-Marie Lambeir, Jurgen Joossens, Koen Augustyns, Joris G De Man, Ingrid De Meester, Benedicte Y De Winter

Abstract

Proteases, enzymes catalyzing the hydrolysis of peptide bonds, are present at high concentrations in the gastrointestinal tract. Besides their well-known role in the digestive process, they also function as signaling molecules through the activation of protease-

Match Overview

1	Internet 77 words crawled on 20-Nov-2016 www.wjgnet.com	2%
2	Crossref 30 words Nathalie Vergnolle, "Modulation of visceral pain and inflammation by protease-activated receptors", <i>British Journal of Gastroenterology</i> , 2004; 48(10): 1101-1108	1%
3	Internet 23 words crawled on 15-Oct-2013 www.europeanreview.org	<1%
4	Crossref 18 words Fang, Y. "Label-free receptor assays", <i>Drug Discovery Today: Technologies</i> , 2010; 2(1): 21-28	<1%
5	Crossref 17 words N. Cenac, "Proteases and Protease-Activated Receptors (PARs): Novel Signals for Pain", <i>Current Topics in Medicinal Chemistry</i> , 2009; 9(1): 1-10	<1%
6	Crossref 17 words T. Li, "Induction of Interleukin-6 Release from Monocytes by Serine Proteinases and its Potential Mechanisms", <i>Journal of Cellular Biochemistry</i> , 2004; 92(1): 1-10	<1%
7	Publications 17 words Kolfschoten, Lauren. "The pathophysiology, diagnosis, and treatment of IBS: a thorough history and examination of the literature", <i>Journal of Clinical Gastroenterology</i> , 2004; 38(1): 1-10	<1%
8	Crossref 15 words Roka, R. "A Pilot Study of Fecal Serine-Protease Activity: A Pathophysiologic Factor in Diarrhea-Predominant Irritable Bowel Syndrome", <i>Journal of Clinical Gastroenterology</i> , 2004; 38(1): 1-10	<1%

Visceral hypersensitivity in inflammatory bowel diseases and irritable bowel synd

全部 图片 视频 新闻 地图 图书

找到约 203,000 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Visceral hypersensitivity in inflammatory bowel diseases and irritable bowel syndrome: the role of proteases](#)

[... activity in visceral pain in irritable bowel syndrome](#) - Cenac - 被引用次数: 433

[... visceral hypersensitivity and improves intestinal ...](#) - Klooker - 被引用次数: 248

[... sensory neurons in irritable bowel syndrome](#) - Barbara - 被引用次数: 505

[Protease inhibition as new therapeutic strategy for GI diseases ...](#)
[gut.bmj.com/content/65/7/1215.full](#)

12 Apr 2016 ... The therapeutic potential of **protease** inhibition in GI **diseases** is parts of the **colon** of those patients, where no **inflammatory** cell infiltration was detected. PAR2 activation causes **visceral hypersensitivity**, modifies **intestinal motility** Role for **protease** activity in visceral pain in **irritable bowel syndrome**.

[Role for protease activity in visceral pain in irritable bowel syndrome](#)
<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1794118/>

15 Feb 2007 ... **Role** for **protease** activity in visceral pain in **irritable bowel syndrome** ... IBS can also appear in patients with **inflammatory bowel disease** (IBD) who IBS tissue supernatants cause **visceral hypersensitivity** in mice: a **protease**

[Irritable bowel syndrome : Nature Reviews Disease Primers](#)
[www.nature.com/articles/nrdp201614](#)

24 Mar 2016 ... **Irritable bowel syndrome** (IBS) is a functional gastrointestinal **disease** with a high prevalence. ... hyperpermeability, dysbiosis, **inflammation**, **visceral hypersensitivity**, ... QOL and the putative **roles** of **inflammation**, genetics, the **intestinal** Increased levels of histamine and **proteases** in biopsy supernatants[§] .

[The Immune System in Irritable Bowel Syndrome](#)
[www.jnmjournal.org/view.html?uid=217&vmd=Full](#) ▼



全部

新闻

图片

视频

购物

更多 ▼

搜索工具

找到约 211,000 条结果 (用时 0.74 秒)

Google 学术 : Visceral hypersensitivity in inflammatory bowel diseases and irritable bowel syndrome: The role of proteases

... activity in visceral pain in irritable bowel syndrome - Cenac - 被引用次数 : 433

... visceral hypersensitivity and improves intestinal ... - Klooker - 被引用次数 : 248

... sensory neurons in irritable bowel syndrome - Barbara - 被引用次数 : 505

Role for protease activity in visceral pain in irritable bowel syndrome

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : N Cenac - 2007 - 被引用次数 : 433 - 相关文章

2007年2月15日 - Role for protease activity in visceral pain in irritable bowel syndrome ... IBS can also appear in patients with inflammatory bowel disease (IBD) who IBS tissue supernatants cause visceral hypersensitivity in mice: a protease ...

Irritable Bowel Syndrome: The Role of Food in Pathogenesis and ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : PA Hayes - 2014 - 被引用次数 : 35 - 相关文章

Keywords: Irritable bowel syndrome, diet, FODMAPs, gut hormones, gluten, food ... from gut dysmotility, visceral hypersensitivity, low-grade mucosal inflammation, Some of the symptoms that commonly occur in patients with celiac disease, amounts of proteases, which have been shown to stimulate sensory afferents in ...

Protease inhibition as new therapeutic strategy for GI diseases ...

gut.bmj.com > Volume 65, Issue 7 - 翻译此页

作者 : N Vergnolle - 2016 - 被引用次数 : 3 - 相关文章

2016年4月12日 - The therapeutic potential of protease inhibition in GI diseases is parts of the colon of those patients, where no inflammatory cell infiltration was detected. PAR2 activation causes visceral hypersensitivity, modifies intestinal motility Role for protease activity in visceral pain in irritable bowel syndrome.