

全部 图片 新闻 视频 购物 更多▼ 搜索工具

找到约 16,800 条结果 (用时 0.66 秒)

Mo1960 Dysphagia After Sleeve Gastrectomy: Evaluation of Risk ...

www.gastrojournal.org/article/S0016-5085(16)32802-5/abstract - 翻译此页作者: A Nath - 2016

Mo1960 Dysphagia After Sleeve Gastrectomy: Evaluation of Risk Factors and Assessment of Intervention ... Roux-en-Y Gastric Bypass and Vertical Sleeve Gastrectomy: Divergent Pathways to Improved ... Mo1961 Endoscopic Septotomy: An Effective Strategy for Drainage of Sleeve Gastrectomy-Associated Collections.

[PDF] The role of endoscopy in the bariatric surgery patient - ASGE

www.asge.org/uploadedFiles/.../**Endoscopy**_bariatric_surgery_patient.pdf ▼ 翻译此页作者: JA Evans - 2015 - 被引用次数: 12 - 相关文章

sleeve gastrectomy (SG) has supplanted LAGB in this re- gard. Vertical ... upper GI endoscopic evaluation before bariatric surgery. 18. Multiple The endoscopist should assess for pouch dilation or Obesity itself is a risk factor for GERD, which is present in or dysphagia, usually within the first year after surgery. Ste-.

Endoscopic Evaluation of Symptomatic Patients following Bariatric ...

www.hindawi.com/journals/dte/2012/753472/▼翻译此页

作者: M Subhani - 2012 - 被引用次数: 6 - 相关文章

2012年3月11日 - Gastric bypass surgery is currently the only effective treatment option available for

Risk factors for anastomotic leaks after sleeve gastrectomy include poor suture line healing ...

Dysphagia is a troubling complication in the bariatric surgery patient. Upper endoscopy is a useful tool to assess this complaint.

29835-Review-check.docx

Quotes Included Bibliography Included

1% SIMILAR

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 29835

Manuscript Type: ORIGINAL ARTICLE

Retrospective Cohort Study

Dysphagia after vertical sleeve gastrectomy: Evaluation of risk factors and assessment of endoscopic intervention

Nath A. et al. Dysphagia after sleeve gastrectomy

Anand Nath, Sayali Yewale, Tung Tran, John S Brebbia, Timothy R Shope, Timothy R Koch

Abstract

AIM

Vertical sleeve gastrectomy is being performed more frequently worldwide as a treatment for medically-complicated obesity, and dysphagia is common post-















全部 图片 新闻 视频 购物 更多▼ 搜索工具

找到约 10,600 条结果 (用时 0.74 秒)

[PDF] The role of endoscopy in the bariatric surgery patient - ASGE www.asge:org/uploadedFiles/.../Endoscopy_bariatric_surgery_patient.pdf ▼ 翻译此页 作者: JA Evans - 2015 - 被引用次数: 14 - 相关文章 sleeve gastrectomy (SG) has supplanted LAGB in this re- gard. Vertical ... upper GI endoscopic evaluation before bariatric surgery.18. Multiple The endoscopist should assess for pouch dilation or ... Obesity itself is a risk factor for GERD, which is present in or dysphagia, usually within the first year after surgery. Ste-.

Endoscopic Evaluation of Symptomatic Patients following Bariatric ... https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页 作者: M Subhani - 2012 - 被引用次数: 7 - 相关文章 2012年5月15日 - Then came about laparoscopic vertical banded gastroplasty (LVBG), followed by Risk factors for anastomotic leaks after sleeve gastrectomy include poor suture line healing ... Dysphagia is a troubling complication in the bariatric surgery patient. Upper endoscopy is a useful tool to assess this complaint.

Endoscopic management of post-bariatric surgery complications - NCBI www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页 2016年9月16日 - In the immediate 48 h after LRYGB, hemorrhage is reported to occur with an The most common endoscopic interventions performed for the ... The most important factor in the immediate control of bleeding is likely Sleeve gastrectomy and the risk of leak; a systematic analysis of 4,888 patients.

Mo1962 Systematic Treatment of Sleeve Gastrectomy Stenosis With ... 14_9/abstract_翻译此而



全部 图片 新闻 视频 购物 更多▼ 搜索工具

找到约 10,700 条结果 (用时 0.80 秒)

IPDF] The role of endoscopy in the bariatric surgery patient - ASGE www.asge.org/uploadedFiles/.../Endoscopy_bariatric_surgery_patient.pdf ▼ 翻译此页作者: JA Evans - 2015 - 被引用次数: 15 - 相关文章 sleeve gastrectomy (SG) has supplanted LAGB in this re- gard. Vertical ... upper Gl endoscopic evaluation before bariatric surgery.18. Multiple The endoscopist should assess for pouch dilation or ... Obesity itself is a risk factor for GERD, which is present in or dysphagia, usually within the first year after surgery. Ste-.

Gastrointestinal Complications After Bariatric Surgery - NCBI

https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页作者: IT Ma - 2015 - 被引用次数: 3 - 相关文章

Laparoscopie adjustable gastric banding (A) and vertical sleeve gastrectomy (B). which can delay diagnosis and treatment.34 Identified risk factors for patients who after bariatric surgery ranges from 22% to 71%.6,52,53 In assessing the ... with endoscopic evaluation and treatment being the primary approaches.

Gastric leaks post sleeve gastrectomy: Review of its prevention and ... https://www.ncbi.nlm.nih.gov > NCBI > Literature > PubMed Central (PMC) - 翻译此页作者: AA Rached - 2014 - 被引用次数: 29 - 相关文章 2014年10月14日 - Gastric sleeve gastrectomy has become a frequent bariatric procedure. ... After diagnosis, the management of leak depends mainly on the clinical ... The management options include also endoscopic interventions with closure ... Fever and tachycardia are the two most important clinical factors in the ...

Assessment of Complications Following Primary Bariatric Surgery ...
www.academia.edu/.../Assessment_of_Complications_Following_Primary_Bariatric_Sur...
4 OS2.03 Endoscopic Internal Drainage for Treatment of Leaks Following Sleeve Gastrectomy 7
OS2.12 Residual Gastric Volume After Sleeve Gastrectomy 15 OS6.05 Preparing Patients for
Bariatric Surgery: Evaluating Methodology Roux-en-Y Gastric Bypass and Gastric Banding):
Incidence and Risk Factors at ...