

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 20752

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Interleukin-22 contributes to liver regeneration in concanavalin A-mediated murine hepatitis model after hepatectomy

Zhang YM *et al.* IL-22 promote liver regeneration of CP

Ya-Min Zhang, Zi-Rong Liu, Zi-Lin Cui, Chao Yang, Long Yang, Yang Li,
Zhong-Yang Shen

Yan-Min Zhang, Zi-Lin Cui, Long Yang, Yang Li, Zhong-Yang Shen,
Department of Hepatobiliary Surgery, First Central Hospital, Tianjin 300192,
China

Match Overview

1	CrossCheck 109 words Han, Ke-Qi. "Inflammatory microenvironment and expression of chemokines in hepatocellular carcinoma", <i>World Journal of Gastroenterology</i> , 11/11/2007	2%
2	CrossCheck 90 words Yu, Hong-Jing. "Probiotic BIFIDCO cocktail ameliorates Helicobacter pylori induced gastritis", <i>World Journal of Gastroenterology</i> , 11/11/2007	2%
3	CrossCheck 51 words Svetlana Radaeva. "Interleukin 22 (IL-22) plays a protective role in T cell-mediated murine hepatitis: IL-22 is a survival factor", <i>Journal of Interpersonal Medicine</i> , 01/01/2007	1%
4	CrossCheck 29 words X. Ren. "IL-22 is involved in liver regeneration after hepatectomy", <i>AJP Gastrointestinal and Liver Physiology</i> , 01/01/2007	1%
5	CrossCheck 27 words Brandoni, Anabel. "Expression of renal Oat5 and NaDC1 transporters in rats with acute biliary obstruction", <i>World Journal of Gastroenterology</i> , 11/11/2007	1%
6	CrossCheck 21 words Xiaoling Jin. "Interleukin-6 inhibits oxidative injury and necrosis after extreme liver resection", <i>Hepatology</i> , 09/2007	<1%

学术搜索

获得 5 条结果 (用时0.08秒)

文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☒ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Myeloid STAT3 inhibits T cell-mediated hepatitis by regulating T helper 1 cytokine and interleukin-17 production

F Lafdil, H Wang, O Park, W Zhang, Y Moritoki, S Yin... - Gastroenterology, 2009 - Elsevier
... Activation of STAT3 in T Cells Contributes to the Th17 Cytokine IL-17, But Not Th1 ... 33 we hypothesized that higher levels of IFN-γ and IL-17 may contribute to enhanced ... of CD4 + T cells produces inflammatory cytokines and controls CD8 + T-cell cytotoxicity, contributing to the ...
被引用次数: 66 相关文章 所有 9 个版本 引用 保存

nih.gov 中的 [HTML]

[HTML] Therapeutic opportunities of the IL-22-IL-22R1 system

R Sabat, W Quyang, K Wolk - Nature reviews Drug discovery, 2014 - nature.com
... to produce chemokines that attract neutrophilic granulocytes, and it thereby contributes to the ... cytokine receptor system that has high potential clinical relevance is the interleukin-22 (IL-22 ... might have a beneficial impact in inflammatory bowel disease, alcoholic liver and pancreas ...
被引用次数: 76 相关文章 所有 7 个版本 引用 保存

nature.com 中的 [HTML]

Opposite effects of high and low doses of interleukin-2 on T cell-mediated hepatitis in mice (interleukin-2 on hepatitis)

X Zhang, HX Wei, S Rui, H Wei, Z Tian - Hepatology international, 2010 - Springer
... X. Zhang and H.-X. Wei contributed equally to this work. ... Therefore, these molecules and cytokines may contribute to apoptosis of NKT cells. ... Radaeva S, Sun R, Pan HN, Hong F, Gao B. Interleukin 22 (IL-22) plays a protective role in T cell-mediated murine hepatitis: IL-22 is a ...
被引用次数: 7 相关文章 所有 8 个版本 引用 保存

nih.gov 中的 [HTML]

Stromal TIMP3 regulates liver lymphocyte populations and provides protection against Th1 T cell-driven autoimmune hepatitis

A Murthy, YW Shao, V Defamie... - The Journal of ..., 2012 - Am Assoc Immunol
... 1 AM and YWS contributed equally to ... Alternatively, circulating dendritic cells can contribute to protective proinflammatory responses during infection by recruiting lymphocytes to the ... oxygen species is a feature of TNF-mediated stress signaling and potentially contributes to the ...
被引用次数: 15 相关文章 所有 4 个版本 引用 保存

Rapamycin prevents Concanavalin A - induced liver injury by inhibiting lymphocyte activation

Y Zhang, X Xiao, X Li, H Wei - Journal of gastroenterology and ..., 2009 - Wiley Online Library
... Interleukin 22 (IL-22) plays a protective role in T cell-mediated murine hepatitis: IL-22 is a survival factor for hepatocytes via STAT3 activation. Hepatology 2004; 39: 1332-42. ... Critical contribution of liver natural killer T cells to a murine model of hepatitis. ...
被引用次数: 2 相关文章 所有 4 个版本 引用 保存

[网页](#) [图片](#) [新闻](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 14,700 条结果 (用时 0.94 秒)

Google 学术 : Interleukin-22 contributes to liver regeneration in concanavalin A-mediated murine hepatitis model after hepatectomy

... STAT3 inhibits T cell-mediated hepatitis by regulating T ... - Lafdil - 被引用次数 : 67

Therapeutic opportunities of the IL-22-IL-22R1 system - Sabat - 被引用次数 : 82

... -mediated hepatitis in mice (interleukin-2 on hepatitis) - Zhang - 被引用次数 : 8

In vivo consequences of liver-specific interleukin-22 ...

www.ncbi.nlm.nih.gov > ... > PubMed Central (PMC) ▾ [翻译此页](#)

作者 : Q Park - 2011 - 被引用次数 : 101 - [相关文章](#)

Diethylnitrosamine (DEN)-induced hepatocellular carcinoma model ... and AST in this T cell hepatitis model induced by different doses of Concanavalin A, ... IL-22TG mice have accelerated liver regeneration after partial hepatectomy (PHx) Gao B. Interleukin 22 (IL-22) plays a protective role in T cell-mediated murine ...

Hepatoprotective and antifibrotic functions of interleukin22 ...

onlinelibrary.wiley.com/doi/10.1111/jgh.12032/pdf - [翻译此页](#)

作者 : X Kong - 2013 - 被引用次数 : 19 - [相关文章](#)

treatment with IL-22 in vitro promotes proliferation of liver progenitor cells via a signal transducer and ... murine liver injury induced by Concanavalin A (Con A), carbon ... inhibited STAT3 activation.²³ In a liver regeneration model, the levels of ... sion were significantly increased after 70% partial hepatectomy. Blockage of ...

Th17 cells and their associated cytokines in liver ... - Nature

www.nature.com > Journal home > Archive > Reviews - [翻译此页](#)

作者 : F Lafdil - 2010 - 被引用次数 : 55 - [相关文章](#)