

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 20432

Manuscript Type: Basic Study

Choledochojejunostomy with an innovative magnetic compressive anastomosis: How to determine optimal pressure?

Xue F *et al.* Optimal magnetic pressure in choledochojejunostomy magnamosis

Fei Xue, Hong-Chang Guo, Jian-Peng Li, Jian-Wen Lu, Hao-Hua Wang, Feng Ma, Ya-Xiong Liu, Yi Lv

Fei Xue, Jian-Peng Li, Jian-Wen Lu, ⁵Yi Lv, Department of Hepatobiliary Surgery, First Affiliated Hospital, Xi'an Jiaotong University, Xi'an 710061, Shaanxi Province, China

Fei Xue, Jian-Peng Li, Jian-Wen Lu, Hao-Hua Wang, Feng Ma, Yi Lv, XJTU Research Institute of Advanced Surgical Technology and Engineering, Xi'an

Match Overview

1	CrossCheck 109 words Pan, Qin. "Linked PNPLA3 polymorphisms confer susceptibility to nonalcoholic steatohepatitis and decreased viral lo ..."	2%
2	CrossCheck 33 words Pang, Qing. "Platelet to lymphocyte ratio as a novel prognostic tool for gallbladder carcinoma", <i>World Journal of Gastro</i>	1%
3	CrossCheck 29 words Lambe, T., M. G. O. Riordain, R. A. Cahill, and P. Cantillon-Murphy. "Magnetic Compression in Gastrointestinal and E ..."	<1%
4	CrossCheck 23 words Sipos, Ferenc. "Association of hepatocyte-derived growth factor receptor/caudal type homeobox 2 co-expression with ..."	<1%
5	CrossCheck 21 words Yan, Xiao-Peng. "Magnamosis combined with endoscopy ... A new endoscopic technique in digestive surgery", <i>World C</i>	<1%



学术搜索

找到约 471 条结果 (用时0.04秒)

文章

我的图书馆

您是不是要找: Prognostic value of **perioperative** leukocyte count in resectable gastric cancer

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

时间不限

2015以来

2014以来

2011以来

自定义范围...

Prospective randomized trial of early postoperative intraperitoneal chemotherapy as an adjuvant to **resectable gastric cancer**.

nih.gov 中的 [PDF]

W Yu, I Whang, I Suh, A Averbach, D Chang... - Annals of ..., 1998 - ncbi.nlm.nih.gov
 ... Eligible patients had a **white blood cell count** of at least 4000/mcl or more, a platelet **count** 150,000/mcl or more, a blood urea nitrogen **level** of less than 30 mg/dl, and a creatinine ... Mean of follow-up in months (**range**) ... Surgery Surgery + EPIC Only Category (%) (% p value ...
 被引用次数: 243 相关文章 所有 5 个版本 引用 保存

按相关性排序

按日期排序

Neo-adjuvant chemotherapy for operable **gastric cancer**: long term results of the Dutch randomised FAMTX trial

HH Hartgrink, CJH Van de Velde, H Putter... - European Journal of ..., 2004 - Elsevier
 ... With a two-sided significance **level** of 0.05 and a power of 0.90, requiring 225 patients in each arm; the ... n, Median, 5 year, p-Value. ... For the group without response the prolonged time interval between randomisation and operation may have led to a strongly decreased **prognosis**.
 被引用次数: 157 相关文章 所有 7 个版本 引用 保存

搜索所有网页

中文网页

简体中文网页

... III trial of preoperative chemotherapy with hyperfractionation radiotherapy plus surgery versus surgery alone for **resectable esophageal squamous cell carcinoma**

oxfordjournals.org 中的 [HTML]

JL Lee, SI Park, SB Kim, HY Jung, GH Lee... - Annals of ..., 2004 - Eur Soc Med Oncology
 ... creatinine clearance of >50 ml/min; normal liver function with serum bilirubin **level** of <1.5 ... grade 2 or worse renal toxicity, cisplatin was withheld until the creatinine **value** recovered to ... Multivariate analysis showed that only weight loss was an independent **prognostic** factor for OS ...
 被引用次数: 241 相关文章 所有 5 个版本 引用 保存

包括专利

包含引用

创建快讯

[PDF] Pretreatment serum albumin as a predictor of **cancer** survival: a systematic review of the epidemiological literature

biomedcentral.com 中的 [PDF]

D Gupta, CG Lis - Nutr J, 2010 - biomedcentral.com
 ... analysis, serum albumin **level** and the pathologic T, N statuses were independent factors correlated with **prognosis** [63]. A study was done to assess the **value** of combination of hypoalbuminemia and an elevated circulating concentra- tion of CRP as a **prognostic** score in ...
 被引用次数: 147 相关文章 所有 18 个版本 引用 保存 更多

Surgical treatment of pancreatic **adenocarcinoma**: actual survival and **prognostic** factors in 343 patients

researchgate.net 中的 [PDF]

KFD Kuhlmann, SMM De Castro, JG Wesseling... - ... Journal of Cancer, 2004 - Elsevier
 ... In the operating room, the surgeon marked the resection margins at the resection specimen at



[全部](#) [图片](#) [视频](#) [购物](#) [新闻](#) [更多 ▾](#) [搜索工具](#)

找到约 7,540 条结果 (用时 0.62 秒)

您是不是要找：[Choledochojejunostomy with an innovative magnetic **compression** anastomosis: How to determine optimal pressure?](#)

[compressive anastomosis Publications and Abstracts](#)

[www.pubfacts.com/search/compressive+anastomosis](#) ▾ [翻译此页](#)

Surgical stapling device-tissue interactions: what surgeons need to know to ... limitations in regards to tissue interactions that, if known, allow for optimal use. ... To investigate the advantages of magnetic compressive anastomosis (MCA) ... and varying compressive pressures when applied to create ex vivo intestinal seals.

[Extrahepatic portacaval shunt via a magnetic compression ...](#)

[www.ncbi.nlm.nih.gov](#) > ... > Literature > PubMed Central (PMC) - [翻译此页](#)

作者：XP Yan - 2015

2015年7月14日 - Core tip: The portacaval shunt using magnetic compression technique ... Previously, we applied Roux-en-Y choledochojunostomy ... the portacaval anastomosis using the magnetic compression technique. The average diameters of the IVC determined from CT image Innovations and breakthroughs.

缺少字词：pressure

[Portacaval Shunt Established in Six Dogs Using Magnetic ...](#)

[journals.plos.org/plosone/article?id=10.1371/journal.pone...](#) ▾ [翻译此页](#)

作者：X Yan - 2013 - 被引用次数：8 - 相关文章

2013年9月30日 - The daughter magnet was moved to the anastomosis position on the ... choledochojunostomy procedures using magnetic compression in a canine ... We found that 5 to 7 days after the first surgery was the best time to perform the ...