



Glycosyltransferases and non-alcoholic fatty liver disease



学术搜索

找到约 203 条结果 (用时0.06秒)

我的著作引用情况



文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

Diagnostic utility of laboratory tests in alcoholic liver disease.

AS Rosman, CS Lieber - Clinical chemistry, 1994 - Am Assoc Clin Chem

... III Non- Alcoholic Fatty Cdm Uver I- *4 I- * -U-I I * ... Values for serum Fab-P- ill-P >45 pgIL (the upper limit of those with steatosis) detected 55% of patients with PVF, 62% of those with septal fibrosis, and 90% of the cirrhotic subjects (24). ... Non- Slapis Alcoholic Fatty Control Liver ...

被引用次数: 73 相关文章 所有 5 个版本 引用 保存

clinchem.org 中的 [PDF]

Carbohydrate deficient transferrin in alcoholic liver disease: mechanisms and clinical implications

CS Lieber - Alcohol, 1999 - Elsevier

... is that ethanol intake decreases the activity of glycoprotein glycosyltransferase enzymes, namely ... stem from the decreased activities of liver glycoprotein glycosyltransferases involved in ... liver disease (both abstinent alcoholics and patients with non-alcoholic liver disease), the IEF ...

被引用次数: 30 相关文章 所有 7 个版本 引用 保存

High fat diet induced hepatic steatosis and insulin resistance: Role of dysregulated ceramide metabolism

L Longato, M Tong, JR Wands... - Hepatology ..., 2012 - Wiley Online Library

... SPTLC), and (f) uridine diphosphate (UDP) glucose ceramide glycosyltransferase (UGCG). ... Figure 4. Chronic high fat diet (HFD) causes hepatic oxidative stress. ... NON-ALCOHOLIC FATTY LIVER disease and NASH, like T2DM and metabolic syndrome, are insulin resistance ...

被引用次数: 37 相关文章 所有 8 个版本 引用 保存

nih.gov 中的 [HTML]

Altered UDP-glucuronosyltransferase and sulfotransferase expression and function during progressive stages of human nonalcoholic fatty liver disease

aspetjournals.org 中的 [HTML]



19521-Review

BY YU-TAO ZHAN

Quotes Excluded
Bibliography Excluded14%
SIMILAR

Name of journal: World Journal of Gastroenterology

ESPS Manuscript NO: 19521

Manuscript Type: Review

Glycosyltransferases and non-alcoholic fatty liver disease

Yu-Tao Zhan, Hai-Ying Su, Wei An

Match Overview

1	Internet 210 words crawled on 06-Jan-2015 www.ncbi.nlm.nih.gov	2%
2	Internet 156 words crawled on 24-Apr-2010 glycob.oxfordjournals.org	2%
3	CrossCheck 108 words Jianxin Fu. "Endothelial cell O-glycan deficiency causes blood/lymphatic misconnections and consequent fatty ..."	1%
4	Internet 81 words crawled on 15-Mar-2014 dmd.aspetjournals.org	1%
5	Internet 76 words crawled on 23-Apr-2014 www.bayer.com	1%
6	Internet 70 words crawled on 06-Jun-2012 jwk.ucsf.edu	1%



Glycosyltransferases and non-alcoholic fatty liver disease



学术搜索

找到约 203 条结果 (用时0.06秒)

我的著作引用情况



文章

我的图书馆

时间不限

2015以来

2014以来

2011以来

自定义范围...

按相关性排序

按日期排序

搜索所有网页

中文网页

简体中文网页

☒ 包括专利☐ 包含引用

小提示：只搜索中文(简体)结果，可在 学术搜索设置 指定搜索语言

Diagnostic utility of laboratory tests in alcoholic liver disease.

AS Rosman, CS Lieber - Clinical chemistry, 1994 - Am Assoc Clin Chem

... III Non- Alcoholic Fatty Cdmi Uver I- *4 I- * -U-I I * ... Values for serum Fab-P- ill-P >45 pgIL (the upper limit of those with steatosis) detected 55% of patients with PVF, 62% of those with septal fibrosis, and 90% of the cirrhotic subjects (24). ... Non- Slapis Alcoholic Fatty Control Liver ...

被引用次数: 73 相关文章 所有 5 个版本 引用 保存

clinchem.org 中的 [PDF]

Carbohydrate deficient transferrin in alcoholic liver disease: mechanisms and clinical implications

CS Lieber - Alcohol, 1999 - Elsevier

... is that ethanol intake decreases the activity of glycoprotein glycosyltransferase enzymes, namely ... stem from the decreased activities of liver glycoprotein glycosyltransferases involved in ... liver disease (both abstinent alcoholics and patients with non-alcoholic liver disease), the IEF ...

被引用次数: 30 相关文章 所有 7 个版本 引用 保存

High fat diet induced hepatic steatosis and insulin resistance: Role of dysregulated ceramide metabolism

L Longato, M Tong, JR Wands... - Hepatology ..., 2012 - Wiley Online Library

... SPTLC), and (f) uridine diphosphate (UDP) glucose ceramide glycosyltransferase (UGCG). ... Figure 4. Chronic high fat diet (HFD) causes hepatic oxidative stress. ... NON-ALCOHOLIC FATTY LIVER disease and NASH, like T2DM and metabolic syndrome, are insulin resistance ...

被引用次数: 37 相关文章 所有 8 个版本 引用 保存

nih.gov 中的 [HTML]

Altered UDP-glucuronosyltransferase and sulfotransferase expression and function during progressive stages of human nonalcoholic fatty liver disease

aspetjournals.org 中的 [HTML]

[网页](#) [新闻](#) [图片](#) [视频](#) [购物](#) [更多 ▾](#) [搜索工具](#)

找到约 108,000 条结果 (用时 0.63 秒)

Google 学术 : Glycosyltransferases and non-alcoholic fatty liver disease

... utility of laboratory tests in alcoholic liver disease: - Rosman - 被引用次数 : 73

... deficient transferrin in alcoholic liver disease: ... - Lieber - 被引用次数 : 30

... fat diet induced hepatic steatosis and insulin resistance ... - Longato - 被引用次数 : 37

Lipids in Health and Disease | Full text | Mechanism of the ...

www.lipidworld.com/content/14/1/43 ▾ [翻译此页](#)作者 : Y Zhan - 2015 - [相关文章](#)

2015年5月8日 - Recent studies have shown that some glycosyltransferases are involved in the development of nonalcoholic fatty liver disease (NAFLD).

[PDF] Mechanism of the effect of glycosyltransferase GLT8D2 o...

www.lipidworld.com/content/pdf/s12944-015-0040-3.pdf ▾ [翻译此页](#)作者 : Y Zhan - 2015 - [相关文章](#)

GLT8D2 expression increased in steatosis HepG2 cells compared with that in normal ... Non-alcoholic fatty liver disease (NAFLD) is one of the most common ...

Mechanism of the effect of glycosyltransferase GLT8D2 on ...

www.ncbi.nlm.nih.gov > ... > Literature > PubMed Central (PMC) - [翻译此页](#)作者 : Y Zhan - 2015 - [相关文章](#)

2015年5月8日 - Recent studies have shown that some glycosyltransferases are involved in the development of nonalcoholic fatty liver disease (NAFLD).

Mechanism of the effect of glycosyltransferase GLT8D2 on ...