

Effect of treatment failure on the CagA EPIYA motif in Helicobacter pylori strains f

全部 图片 视频 新闻 地图 图书

找到约 547 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Effect of treatment failure on the CagA EPIYA motif in Helicobacter pylori strains from Colombian subjects](#)

[... of cagA in Helicobacter pylori strains from Colombian ...](#) - Loh - 被引用次数: 32

[... of Helicobacter pylori infection and histologic gastritis ...](#) - Dooley - 被引用次数: 932

[Helicobacter pylori infection](#) - Suerbaum - 被引用次数: 2580

[Analysis of cagA in Helicobacter pylori strains from Colombian ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3189306/>

22 Aug 2011 ... CagA proteins with a greater number of EPIYA motifs undergo more ... Several strains were excluded from analysis due to contamination or failure to grow in liquid broth culture. ... Expression of CagA in Colombian H. pylori strains ... Gastric mucosa biopsy samples from the 36 subjects were obtained from ...

[CagA C-terminal variations in Helicobacter pylori strains from ... - NCBI](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2837774/>

15 May 2009 ... The C-terminus of the Helicobacter pylori CagA protein is polymorphic, ... lesions was examined in H. pylori-infected Colombian patients from areas of high and low risk for GC. ... Strains with one EPIYA-C motif were associated with less severe ... In most infected subjects, this bacterium survives for decades ...

[Analysis of Helicobacter pylori cagA Promoter Elements Required ...](#)

iai.asm.org/content/80/9/3094.full

Mutagenesis experiments confirmed that two copies of the TAATGA motif are required for ... 14), but these studies failed to detect any effect of salt on CagA expression. ... We then examine salt-regulated CagA expression in H. pylori strains cagA promoter sequences from Colombian strains were introduced into H. pylori ...

[Variations in Helicobacter pylori Cytotoxin-Associated Genes and ...](#)

journals.plos.org/plosone/article?id=10.1371/journal.pone

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 31008

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Effect of treatment failure on the CagA EPIYA motif in *Helicobacter pylori* strains from Colombian subjects

Javier Andres Bustamante-Rengifo, Andres Jenuer Matta, Alvaro Jairo Pazos,
Luis Eduardo Bravo

Abstract

AIM

To evaluate effect of treatment failure on *cagA* and *vacA* genotypes in

Match Overview

1	Internet 140 words crawled on 27-Jun-2016 www.wjgnet.com	3%
2	Internet 83 words crawled on 03-Feb-2009 jcm.asm.org	2%
3	Internet 35 words crawled on 11-Jul-2010 www.actagastro.org	1%
4	Internet 21 words crawled on 12-Dec-2014 onlinelibrary.wiley.com	<1%
5	Crossref 14 words Hasan Umit. "The Relationship Between Virulence Factor... of <i>Helicobacter pylori</i> and Severity of Gastritis in Infected Pa	<1%

Effect of treatment failure on the CagA EPIYA motif in Helicobacter pylori isolates

全部 图片 视频 新闻 购物 地图 图书

找到约 1,560 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Google 学术: Effect of treatment failure on the CagA EPIYA motif in Helicobacter pylori isolates](#)

[... as an intracellular target of Helicobacter pylori CagA ...](#) - Higashi - 被引用次数: 967

[Pathogenesis of Helicobacter pylori infection](#) - Kusters - 被引用次数: 1372

[c-Src/Lyn kinases activate Helicobacter pylori CagA ...](#) - Stein - 被引用次数: 451

[In vitro effect of amoxicillin and clarithromycin on the 3' region of ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC3785626/](#) ▼

28 Sep 2013 ... Resistant isolates at baseline and in amoxicillin and clarithromycin ... The gain and loss of EPIYA motifs resulted in a diversity of H. pylori ... process of Helicobacter pylori in cases with treatment failure. ... Hence, determination of the number of EPIYA motifs within the cagA variable region in clinical H. pylori ...

[Role of Helicobacter pylori cagA EPIYA motif and vacA genotypes for ...](#)

[https://bmcinfectdis.biomedcentral.com/.../10.../1471-2334-12-223](#) ▼

Helicobacter pylori cagA vacA EPIYA Gastric cancer Peptic ulcer disease Southeast Asian ... Greater than 90% of H. pylori strains isolated from East Asian populations The effects of cagA status, EPIYA motif genotype, and vacA genotypes on the risk of pylori cagA and vacA status for the efficacy of antibiotic treatment.

[High Prevalence of Clarithromycin-Resistant Helicobacter pylori ...](#)

[jcm.asm.org/content/48/10/3703.full](#) ▼

We studied 118 H. pylori strains isolated from the same number of patients. ... Many factors have been implicated as causes of treatment failure, including Clarithromycin resistance relationship with cagA gene and vacA mosaicism distribution. of repeating EPIYA phosphorylation motifs in the carboxyl terminus of CagA ...

[JCI - Multistep activation of the Helicobacter pylori effector CagA](#)

[https://www.jci.org/articles/view/61578](#)

Effect of treatment failure on the CagA EPIYA motif in Helicobacter pylori isolates

全部 图片 视频 新闻 购物 地图 图书

找到约 1,790 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

Google 学术: [Effect of treatment failure on the CagA EPIYA motif in Helicobacter pylori isolates](#)

[... as an intracellular target of Helicobacter pylori CagA ...](#) - Higashi - 被引用次数: 975

[Pathogenesis of Helicobacter pylori infection](#) - Kusters - 被引用次数: 1409

[c-Src/Lyn kinases activate Helicobacter pylori CagA ...](#) - Stein - 被引用次数: 460

[In vitro effect of amoxicillin and clarithromycin on the 3' region of ...](#)

[www.ncbi.nlm.nih.gov/pmc/articles/PMC3785626/](#) ▼

28 Sep 2013 ... The gain and loss of EPIYA motifs resulted in a diversity of H. pylori subclones after ... process of Helicobacter pylori in cases with treatment failure. ... CagA- positive clinical isolates with increased number of EPIYA motifs ...

[Helicobacter pylori virulence factor CagA promotes tumorigenesis of ...](#)

[https://biosignaling.biomedcentral.com/.../10.../s12964-015-0111-0](#) ▼

11 Jul 2015 ... H. pylori also affects the prognosis of gastric cancer. ... in determining the clinical outcomes, and clinically isolated H. pylori ... several repeat Glu-Pro-Ile-Tyr-Ala (EPIYA) motifs, while other CagA Although this process was found to be independent of CagA, H. pylori lacking a functional T4SS failed to ...

[In vitro effect of amoxicillin and clarithromycin on the 3 ' region of ...](#)

[https://www.researchgate.net/.../257535535_In_vitro_effect_of_amoxicillin_and_clarithromycin_on_the_3_'_region_of_cagA_gene_in_Helicobacter_...](#)

In vitro effect of amoxicillin and clarithromycin on the 3 ' region of cagA gene in ... The gain and loss of EPIYA motifs resulted in a diversity of H. pylori subclones region in H. pylori isolates from patients with treatment failure and deciphering.

[High Prevalence of Clarithromycin-Resistant Helicobacter pylori ...](#)

[jcm.asm.org/content/48/10/3703.full](#) ▼