

Endosonographic Surveillance of 1-3 cm Gastric Submucosal Tumors Originating

全部 图片 视频 新闻 购物 地图 图书

找到约 1,990 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

[Endosonographic surveillance of small gastrointestinal tumors ...](#)

[www.ncbi.nlm.nih.gov/pubmed/19565047](http://www.ncbi.nlm.nih.gov/pubmed/19565047)

Endosonographic surveillance of small gastrointestinal tumors originating ... INTRODUCTION: Submucosal tumors (SMT) are not uncommonly found during upper endoscopy. Management for small SMT originating from muscularis propria (MP) is ... Patients with large tumors (maximal diameter > 3 cm), heterogeneous echo ...

[\[PDF\] download Full Article \(PDF file\)](#)

[www.jgld.ro/2009/2/7.pdf](http://www.jgld.ro/2009/2/7.pdf)

Abstract. Introduction. Submucosal tumors (SMT) are not ... for small SMT originating from muscularis propria (MP) is ... 3 cm), heterogeneous echopattern, presence of cystic spaces or irregular ... Submucosal tumor (SMT) refers to a bulge underneath the mucosa ... tumors were most commonly located in the gastric body (n.

[Small EUS-suspected gastrointestinal stromal tumors of the stomach ...](#)

[www.eusjournal.com/article.asp?issn=2303-9027;year=2016;...](http://www.eusjournal.com/article.asp?issn=2303-9027;year=2016;...)

They represent a distinct group of GI tumors originating from the interstitial cells of Cajal and are ... with the wide availability of endoscopic/endosonographic examination. ... small hypoechoic muscularis propria (MP) lesion, surgery, surveillance .... gastric submucosal tumors having a median diameter of 1.8 (0.5- 4.0) cm (the ...

[Tunneling endoscopic muscularis dissection for subepithelial tumors ...](#)

[link.springer.com/article/10.1007/s00464-013-3023-3](http://link.springer.com/article/10.1007/s00464-013-3023-3)

14 Jun 2013 ... Twelve patients with SETs originating from the MP of the esophagus (n = 7) or ... when the SETs originate from the muscularis propria (MP) layer [1–3]. Endoscopic surveillance of these patients is a common approach, but the ... 70 W ) approximately 5 cm proximal to the lesion, then a submucosal tunnel was ...

[Upper gastrointestinal submucosal lesions-Clinical and ...](#)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 30963

Manuscript Type: ORIGINAL ARTICLE

*Retrospective Cohort Study*

Endosonographic Surveillance of 1-3 cm <sup>1</sup> Gastric Submucosal Tumors

Originating from Muscularis Propria

Ming-Luen Hu, Keng-Liang Wu, Chi-Sin Changchien, Seng-Kee Chuah,

Yi-Chun Chiu

Abstract

### Match Overview

1	Crossref 148 words "Poster - Endoscopy : Poster - Endoscopy", Journal of Gastroenterology and Hepatology, 2015.	6%
2	Crossref 59 words Fang, Y.J.. "Suggested cutoff tumor size for management of small EUS-suspected gastric gastrointestinal strom...	3%
3	Internet 23 words crawled on 04-Jan-2017 <a href="http://www.omicsonline.org">www.omicsonline.org</a>	1%
4	Internet 18 words crawled on 25-Dec-2016 <a href="http://office.wjgnet.com">office.wjgnet.com</a>	1%
5	Crossref 15 words Susan C. Abraham. "Distinguishing Gastrointestinal S... omal Tumors From Their Mimics: An Update", Advances	1%
6	Internet 13 words crawled on 15-Nov-2016 <a href="http://www.wjgnet.com">www.wjgnet.com</a>	1%
7	Internet 13 words crawled on 30-Apr-2012 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	1%
8	Internet 13 words crawled on 30-Aug-2010 <a href="http://www.clinicaladvances.com">www.clinicaladvances.com</a>	1%



Endosonographic Surveillance of 1-3 cm Gastric Submucosal Tumors Originating

全部 图片 视频 新闻 地图 图书

找到约 3,220 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

[Endosonographic surveillance of small gastrointestinal tumors ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/19565047>

INTRODUCTION: **Submucosal tumors** (SMT) are not uncommonly found during ... Management for small SMT **originating from muscularis propria** (MP) is controversial. ... Patients with large tumors (**maximal diameter** > 3 **cm**), heterogeneous echo ... **Gastric** Mucosa/ultrasonography; Gastrointestinal Stromal Tumors/surgery ...

[\[PDF\] download Full Article \(PDF file\)](#)

[www.jgld.ro/2009/2/7.pdf](http://www.jgld.ro/2009/2/7.pdf)

Abstract. Introduction. **Submucosal tumors** (SMT) are not ... for small SMT **originating from muscularis propria** (MP) is ... 3 **cm**), heterogeneous echopattern, presence of cystic spaces or irregular ... **Submucosal tumor** (SMT) refers to a bulge underneath the mucosa ... tumors were most commonly located in the **gastric** body (n.

[EUS features - GastroHep](#)

[www.gastrohep.com/ebooks/ebook.asp?book=1405120819&id=4](http://www.gastrohep.com/ebooks/ebook.asp?book=1405120819&id=4)

The majority of **submucosal tumors** are asymptomatic and often are found ... recently, GISTs have been reclassified as mesenchymal tumors theorized to **originate** from ... EUS features associated with malignancy include large tumor size (> 4 **cm**), ... Lesions arising from the **muscularis propria** (fourth **endosonographic** layer) ...

[Endoscopic submucosal dissection of gastric fundus subepithelial ...](#)

<https://www.spandidos-publications.com/etm/6/2/391>

25 Jun 2013 ... Endoscopic resection of **gastric** subepithelial **tumors** (SETs) carries a high risk of ... for **gastric** fundus subepithelial **tumors** **originating** from the **muscularis propria**. ... However, the detection of a small SET (diameter, ≤3 **cm**) presents ..... Alternative **endosonographic surveillance** may delay the diagnosis of ...

[# 2037 Endosonographic surveillance of 1–3 cm gastric submucosal ...](#)

找到约 3,220 条结果 (用时 0.75 秒)

## Endosonographic surveillance of small gastrointestinal tumors ... - NCBI

<https://www.ncbi.nlm.nih.gov/pubmed/19565047> - 翻译此页

作者: KH Lok - 2009 - 被引用次数: 50 - 相关文章

INTRODUCTION: **Submucosal tumors** (SMT) are not uncommonly found during ... Management for small SMT **originating from muscularis propria** (MP) is controversial. ... Patients with large tumors (maximal diameter > 3 **cm**), heterogeneous echo ... **Gastric** Mucosa/ultrasonography; Gastrointestinal Stromal Tumors/surgery ...

## Endoscopic approach to subepithelial lesions - NCBI - NIH

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者: L Menon - 2014 - 被引用次数: 19 - 相关文章

They most commonly occur within the **stomach**, but are also regularly noted in the .... As most of these lesions arise from the **muscularis propria** layer of the GI tract wall, ... For small lesions of <1–2 **cm** in size, **surveillance** EGD and/or EUS may be ... (b) EUS view of a hypoechoic carcinoid **tumor originating** from the mucosal ...

## [PDF] Endosonographic Surveillance of Small Gastrointestinal Tumors ...

[www.jgld.ro/2009/2/7.pdf](http://www.jgld.ro/2009/2/7.pdf) ▼ 翻译此页

作者: KH Lok - 2009 - 被引用次数: 50 - 相关文章

Abstract. Introduction. **Submucosal tumors** (SMT) are not ... for small SMT **originating from muscularis propria** (MP) is ... 3 **cm**), heterogeneous echopattern, presence of cystic spaces or irregular ... **Submucosal tumor** (SMT) refers to a bulge underneath the mucosa ... tumors were most commonly located in the **gastric** body (n.

## EUS features - GastroHep

[www.gastrohep.com/ebooks/ebook.asp?book=1405120819&id=4](http://www.gastrohep.com/ebooks/ebook.asp?book=1405120819&id=4) ▼ 翻译此页

The majority of **submucosal tumors** are asymptomatic and often are found ... recently, GISTs have