

Comparing acid steatocrit and faecal elastase estimations for use in M-ANNHE

全部 图片 视频 新闻 地图 图书

找到约 110 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Fecal elastase1 and acid steatocrit estimation in chronic pancreatitis](#)

www.ncbi.nlm.nih.gov/pubmed/20177867

23 Feb 2010 ... Fecal elastase1 and acid steatocrit estimation in chronic pancreatitis. ... measured in 101 patients using polyclonal ELISA and acid steatocrit was measured ... Pancreatic Elastase/analysis*, Pancreatitis, Chronic/complications ...

[PDF] [Faecal elastase 1: not helpful in diagnosing chronic pancreatitis](#) ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1727065/.../v042p00551.pdf>

moderate exocrine pancreatic insufficiency ... estimation of faecal elastase 1 is a valuable ... evaluated by comparing it with faecal chymotrypsin ... faecal chymotrypsin (Boehringer-Mannheim, ... group, they used our staging system with the

[PDF] [Fecal elastase1 and acid steatocrit estimation in chronic pancreatitis](#)

medind.nic.in/ica/t09/i6/icat09i6p201.pdf

using polyclonal ELISA and acid steatocrit was measured in ... estimation of fecal elastase 1 Fecal elastase1 testing does ... into another test tube and 100 µL of 5 M perchloric acid ... as compared to CP patients with normal pancreatic ela-

[PDF] [FESCC, EFLM Postgraduate Course 12 \(2012\) - Handbook.pdf](#)

www.eflm.eu/.../FESCC,

[%20EFLM%20Postgraduate%20Course%202012%20\(2012\)%20-...](#)

Gastroesophageal reflux disease (GERD) is one of the major acid peptic diseases. ... Marino having the highest incidence (more than 8-10-fold higher compared to ... cancer, cystic pancreatic neoplasms and other non endocrine pancreatic tumors. ... Mindemark M, Larsson A. Ruling out IBD: Estimation of the possible ...

[Fecal Elastase 1 Determination in Chronic Pancreatitis - Documents](#)

docslide.us/.../fecal-elastase-1-determination-in-chronic-pancreatitis.html

6 Aug 2016 ... Fifty-three healthy subjects, 44 patients with chronic pancreatitis (22 severe, ... Fecal elastase 1 has a high sensitivity, superior to that of fecal chymotrypsin, in the ... Boehringer Mannheim GmbH Diagnostica, Mannheim, Germany). ... Fecal elastase1 and acid steatocrit estimation in chronic pancreatitis

Match Overview



There are no matching sources for this report

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 31822

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Comparing acid steatocrit and faecal elastase estimations for use in M-ANNHEIM staging for pancreatitis

M Ganesh Kamath, C Ganesh Pai, Asha Kamath, Annamma Kurien

Abstract

AIM

Comparing acid steatocrit and faecal elastase estimations for use in M-ANNHE

全部 图片 视频 新闻 地图 图书

找到约 194 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[\[PDF\] Fecal elastase1 and acid steatocrit estimation in chronic pancreatitis](#)

[medind.nic.in/ica/t09/i6/icat09i6p201.pdf](#)

using polyclonal ELISA and acid steatocrit was measured in ... estimation of fecal elastase.1 Fecal elastase1 testing does ... into another test tube and 100 µL of 5 M perchloric acid ... as compared to CP patients with normal pancreatic ela-

[Fecal elastase-1 determination: 'gold standard' of indirect pancreatic ...](#)

<https://www.ncbi.nlm.nih.gov/pubmed/11589385>

... University Hospital of Heidelberg at Mannheim, Germany. BACKGROUND: Tubeless pancreatic function tests measuring the content of elastase-1 and the activity of chymotrypsin in stool are used with different cut-off levels and with ... insufficiency and is of significantly higher sensitivity than fecal chymotrypsin estimation.

[comparison of fecal elastase 1 for exocrine pancreatic ... - SciELO](#)

[www.scielo.br/scielo.php?script=sci_arttext&pid=S0004...](#)

Fecal elastase is a noninvasive test for pancreatic insufficiency diagnosis. ... The levels of fecal elastase 1 were measured using a commercial kit. ... The diagnosis of pancreatic exocrine insufficiency at an early stage is challenging, ... Balakrishnan V. Fecal elastase1 and acid steatocrit estimation in chronic pancreatitis.

[Belgian consensus on chronic pancreatitis in adults and children ...](#)

[https://www.researchgate.net/.../261879903_Belgian_consensus_on_chronic_p](#)
[ancreatitis_in_adults_and_children_Statements_on_diagnosis_and...](#)

Pain should be assessed using the Izbicki score and preferably treated using the ... Screening for pancreatic cancer is not recommended in CP patients, except in ... 48 M. Delhaye et al. ... for CP diagnosis, a consensus-based scoring system has ... elastase tests ... erides breath, acid steatocrit, and quantitative fecal fat).

[Chronic pancreatitis - Resources - References - Best Practice ...](#)

[https://bestpractice.bmj.com/best-practice/.../67/.../references.html?locale...](#)

11 Nov 2016 ... Pregabalin for decreasing pancreatic pain in chronic pancreatitis. ... of chronic pancreatitis; comparison of outcomes at 16-year follow-up. J Am Coll Surg. 2014; 219:208-

[全部](#)[图片](#)[新闻](#)[购物](#)[视频](#)[更多](#)[设置](#)[工具](#)

找到约 206 条结果 (用时 0.80 秒)

Fecal elastase-1 determination: 'gold standard' of indirect pancreatic ...<https://www.ncbi.nlm.nih.gov/pubmed/11589385> - 翻译此页

作者：S Lüth - 2001 - 被引用次数：68 - 相关文章

... University Hospital of Heidelberg at Mannheim, Germany. BACKGROUND: Tubeless pancreatic function tests measuring the content of elastase-1 and the activity of chymotrypsin in stool are used with different cut-off levels and with ... insufficiency and is of significantly higher sensitivity than fecal chymotrypsin estimation.

缺少字词：comparing acid steatocrit staging

Using faecal elastase-1 to screen for chronic pancreatitis in patients ...<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者：RC Turner - 2006 - 被引用次数：11 - 相关文章

Thus a 2x2 table comparing [FE-1] and clinical diagnosis was constructed. ... Keywords: Faecal elastase-1, diagnostic tests, chronic pancreatitis, acute ... including faecal chymotrypsin estimation, the para-aminobenzoic acid test ... Severity of the acute episode was graded according to Ranson's predictive scoring system.

缺少字词：steatocrit annheim

[PDF] Fecal elastase1 and acid steatocrit estimation in chronic pancreatitismedind.nic.in/ica/t09/i6/icat09i6p201.pdf ▼ 翻译此页

作者：BN Girish - 2009 - 被引用次数：15 - 相关文章

using polyclonal ELISA and acid steatocrit was measured in ... estimation of fecal elastase.1 Fecal elastase1 testing does ... into another test tube and 100 µL of 5 M perchloric acid ... as compared to CP patients with normal pancreatic ela-.

缺少字词：annheim staging