





全部 图片 新闻 视频 购物 更多▼ 搜索工具

获得 9 条结果 (用时 0.76 秒)

您是不是要找: CUL4A associated with epithelial to mesenchymal transition and poor prognosis in perihilar cholangiocarcinoma

Google 学术: CUL4A associates with epithelial to mesenchymal transition and poor prognosis in perihilar cholangiocarcinoma

FBXW7 suppresses epithelial-mesenchymal transition, ... - Yang - 被引用次数: 12

High Infiltration of Tumor-Associated Macrophages Influences Poor ...

www.ncbi.nlm.nih.gov/pubmed/26871785 - 翻译此页

作者: J Zhang - 2016 - 被引用次数: 5 - 相关文章

Tumor-associated macrophages (TAMs) are associated with poor prognosis in numerous ... Epithelialmesenchymal transition (EMT) contributes to invasion and ...

缺少字词: cul4a perihilar cholangiocarcinoma

Extra-Hepatic Bile duct cancer (cholangiocarcinoma) | CancerIndex

www.cancerindex.org/clinks5u.htm ▼ 翻译此页

METHODS: In vitro cultured cholangiocarcinoma QBC939 cell line was ducts is higher than the percentage of the neoplasms with perihilar localization. hepatocyte dedifferentiation, enhanced epithelial-mesenchymal transition ... BACKGROUND: Intrahepatic cholangiocarcinoma (ICC) is associated with poor survival.

Overexpression of epithelial-mesenchymal transition-related markers ...

Quotes Excluded Bibliography Excluded

15% SIMILAR

Name of Journal: World Journal of Gastroenterology

ESPS Manuscript NO: 31422

Manuscript Type: ORIGINAL ARTICLE

Basic Study

iThenticate[®]

Cullin 4A is associated with the epithelial to mesenchymal transition and poor prognosis in perihilar cholangiocarcinoma


Zhang TJ et al. CUL4A in PHCC EMT and prognosis

Tong-Jun Zhang, Dong Xue, Cheng-De Zhang, Ze-Dong Zhang, Qing-Ran Liu, Jian-Qiang Wang

Abstract

AIM

To explore the functional role of Cullin 4A (CUL4A), a core subunit of E3



















图片

全部

购物

更多

工具

找到约 190 条结果 (用时 0.81 秒)

Extra-Hepatic Bile duct cancer (cholangiocarcinoma) | CancerIndex

www.cancerindex.org/clinks5u.htm ▼ 翻译此页

新闻

Information for Health Professionals / Researchers (4 links) bile ducts is higher than the percentage of the neoplasms with perihilar localization. hepatocyte dedifferentiation, enhanced epithelialmesenchymal transition and cell ... BACKGROUND: Intrahepatic cholangiocarcinoma (ICC) is associated with poor survival.

PubPDF - Find Full Text Journal Articles about Cholangiocarcinoma

www.pubpdf.com/search/Cholangiocarcinoma/18 ▼ 翻译此页

视频

Epithelial-to-Mesenchymal Transition and Cancer Invasiveness: What Can We Intrahepatic cholangiocarcinoma (ICC) is associated with poor survival. ... Outcome of curative resection for perihilar. cholangiocarcinoma in Northeast Thailand. ... target genes, including cullin 4A (CUL4A), insulin receptor substrate 2 (IRS2), ...

PubPDF - Find Full Text Journal Articles about Bile Duct Tumors

www.pubpdf.com/search/Bile+Duct+Tumors/7 ▼ 翻译此页

Perihilar cholangiocarcinoma (pCCA) has a dismal prognosis. inflammatory response is associated with a poor long-term prognosis after Interleukin - 6 induces epithelial - mesenchymal transition in human intrahepatic biliary epithelial cells. target genes, including cullin 4A (CUL4A), insulin receptor substrate 2 (IRS2) ...

The Cullin 4A/B-DDB1-Cerebion E3 Ubiquitin Ligase Complex ... - NCBI

https://www.ncbi.nlm.nih.gov/pubmed/26021757 - 翻译此页

作者: YA Chen - 2015 - 被引用次数: 9 - 相关文章



Cullin 4A is associated with the epithelial to mesenchymal transition and poor p



全部

图片

新闻

视频

购物

更多

设置

工具

找到约65条结果(用时0.86秒)

Extra-Hepatic Bile duct cancer (cholangiocarcinoma) | CancerIndex www.cancerindex.org/clinks5u.htm ▼ 翻译此页

METHODS: In vitro cultured cholangiocarcinoma QBC939 cell line was exposed to inflammatory response is associated with a poor long-term prognosis after ... hepatocyte dedifferentiation, enhanced epithelial-mesenchymal transition and target genes, including cullin 4A (CUL4A), insulin receptor substrate 2 (IRS2), ...

PubPDF - Find Full Text Journal Articles about Cholangiocarcinoma www.pubpdf.com/search/Cholangiocarcinoma/18 ▼ 翻译此页

Epithelial-to-Mesenchymal Transition and Cancer Invasiveness: What Can We Intrahepatic cholangiocarcinoma (ICC) is associated with poor survival. ... Outcome of curative resection for perihilar cholangiocarcinoma in Northeast Thailand. ... target genes, including cullin 4A (CUL4A), insulin receptor substrate 2 (IRS2), ...

PubPDF - Find Full Text Journal Articles about Bile Duct Tumors

www.pubpdf.com/search/Bile+Duct+Tumors/7 ▼ 翻译此页
Perihilar cholangiocarcinoma (pCCA) has a dismal prognosis. inflammatory response is
associated with a poor long-term prognosis after Interleukin-6 induces epithelial-mesenchymal
transition in human intrahepatic biliary epithelial cells. target genes, including cullin 4A (CUL4A),
insulin receptor substrate 2 (IRS2)

The cullin 4B-based UV-damaged DNA-binding protein ligase binds to ...

https://www.ncbi.nlm.nih.gov/pubmed/018593899 - 翻译此页

作者: J Guerrero-Santoro - 2008 - 被引用次数: 93 - 相关文章

2008年7月1日 - UV-DDB consists of DDB1 and DDB2, and it is a component of the cullin 4A (CUL4A) -based ubiquitin ligase, DDB1-CUL4A(DDB2).

缺少字词: epithelial mesenchymal transition poor prognosis perihilar cholangiocarcinoma