

Notch mediated by TGF- β /Smad pathway in Concanavalin A-induced liver fibrosi

全部 图片 视频 新闻 购物 地图 图书

找到约 5,940 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Preventive Effect of Halofuginone on Concanavalin A-Induced Liver ...](#)

journals.plos.org/plosone/article?id=10.1371/journal.pone...

16 Dec 2013 ... Since ConA-induced liver fibrosis is caused by the repeated activation of T ... The direct targets in TGF- β 1 pathway, Smads (Smad2, and ... This inhibitory effect was probably mediated by blocking TGF- β induced Smad3 activation [33]. ... Concanavalin A (ConA)-induced liver injury is well accepted as a rat ...

[Preventive Effect of Halofuginone on Concanavalin A-Induced Liver ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3864948/>

16 Dec 2013 ... Since ConA-induced liver fibrosis is caused by the repeated activation of T ... The direct targets in TGF- β 1 pathway, Smads (Smad2, and ... This inhibitory effect was probably mediated by blocking TGF- β induced Smad3 activation [33]. ... Concanavalin A (ConA)-induced liver injury is well accepted as a rat ...

[Hepatoprotective and Anti-fibrotic Agents: It's Time to Take the Next ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4703795/>

7 Jan 2016 ... Hepatic fibrosis and cirrhosis cause strong human suffering and Consequently , catalase, an enzymatic scavenger of H₂O₂, abrogated TGF- β -mediated type I In dimethylnitrosamine (DMN)-induced liver fibrosis in rats, ... and inhibition of cytokine signaling pathways (TGF- β and PDGF; Li et al., 2015a).

[Strategies to prevent and reverse liver fibrosis in humans and ...](#)

https://www.researchgate.net/.../276209969_Strategies_to_prevent_and_reverse_liver_fibrosis_in_humans_and_laboratory_animals

mediated by various signaling pathways such as the JNK, and p38 rat models of liver fibrosis via unknown mechanisms. (Paulsen et al. 2013). Smad3/4 complex formation and TGF- β -induced Smad ... The administration of concanavalin A (ConA). induced Notch shRNA transferred to CCl₄-treated rat (Zheng et al.

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 32048

Manuscript Type: ORIGINAL ARTICLE

Basic Study

Notch signaling mediated by TGF- β /Smad pathway in
concanavalin-A-induced liver fibrosis in rats

Yi Wang, Ruo-Wu Shen, Bing Han, Zhen Li, Le Xiong, Feng-Yu Zhang,
Bei-Bei Cong, Bei Zhang

Abstract

Match Overview

1	Crossref 72 words Xiao, Zhicheng, Jing Zhang, Xiaogang Peng, Yanjun Dong, Lixin Jia, Huihua Li, and Jie Du. "The Notch γ -secretase"	3%
2	Crossref 46 words Clara Dees. "Inhibition of Notch signaling prevents experimental fibrosis and induces regression of established f"	2%
3	Internet 25 words crawled on 25-Jun-2016 www.wjgnet.com	1%
4	Internet 22 words crawled on 09-Nov-2009 www.ncbi.nlm.nih.gov	1%
5	Publications 20 words "Scientists at University of Massachusetts report research in hematology". <i>Blood Weekly</i> , Oct 23 2008 Issue	1%
6	Crossref 19 words Lan, Tian, Tatiana Kisseleva, and David A. Brenner. "Deficiency of NOX1 or NOX4 Prevents Liver Inflammation a ..."	1%
7	Internet 17 words crawled on 13-Oct-2016 www.spandidos-publications.com	1%
8	Crossref 15 words Y Inagaki. "Emerging insights into Transforming growth factor β 1 (TGF- β 1) as a downstream Smad signal in hepatic fibrogenesis", <i>Gut</i> , 2/1/20	1%

Notch signaling mediated by TGF- β /Smad pathway in concanavalin-A-induced liv

全部 图片 视频 新闻 地图 图书

找到约 5,460 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

Google 学术: [Notch signaling mediated by TGF- \$\beta\$ /Smad pathway in concanavalin-A-induced liver fibrosis in rats](#)

[... to prevent and reverse liver fibrosis in humans and ...](#) - Chen - 被引用次数: 9

[Novel investigations of flavonoids as chemopreventive ...](#) - Liao - 被引用次数: 5

[Hepatoprotective and Anti-fibrotic Agents: It's Time to Take the Next ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4703795/>

7 Jan 2016 ... In dimethylnitrosamine (DMN)-induced liver fibrosis in rats, ... and inhibition of cytokine signaling pathways (TGF- β and PDGF; Li et al., Transforming growth factor (TGF- β) and its intracellular signaling mediators, the Smad proteins, transduce signals via the classical JAK-STAT pathway and mediate ...

[Effects of 18 \$\alpha\$ -glycyrrhizin on TGF- \$\beta\$ 1/Smad signaling pathway in ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4396252/>

1 Feb 2015 ... Conclusion: 18 α -GL is able to attenuate CCl₄ induced liver fibrosis in rat. ... improved liver fibrosis by acting on the TGF- β /Smad signal pathway, suggesting that Effects of 18 α -GL on the collagen I and III expression in Rat Liver TGF- β mediates the activation of Smad2 primarily in undifferentiated cells ...

[Preventive Effect of Halofuginone on Concanavalin A-Induced Liver ...](#)

journals.plos.org/plosone/article?id=10.1371/journal.pone... ▼

16 Dec 2013 ... Since ConA-induced liver fibrosis is caused by the repeated activation of T ... by suppressing the synthesis of collagen I and inflammation-mediated liver injury. ... Concanavalin A (ConA)-induced liver injury is well accepted as a rat model of and Smad-independent pathways in TGF- β family signalling.

[PubPDF - Find Full Text Journal Articles Authored By Cheng-hai Liu](#)

www.pubpdf.com/search/author/Cheng-hai+Liu ▼

However, whether HSP affects the Notch signaling pathway is unclear. Effect of Euzhang

[全部](#)[图片](#)[视频](#)[新闻](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 5,480 条结果 (用时 0.91 秒)

Google 学术 : Notch signaling mediated by TGF- β /Smad pathway in concanavalin-A-induced liver fibrosis in rats

... to prevent and reverse liver fibrosis in humans and ... - Chen - 被引用次数 : 9

Novel investigations of flavonoids as chemopreventive ... - Liao - 被引用次数 : 5

Hepatoprotective and Anti-fibrotic Agents: It's Time to Take the Next Step<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC4703795/> - 翻译此页

作者 : R Weiskirchen - 2015 - 被引用次数 : 2 - 相关文章

2016年1月7日 - In dimethylnitrosamine (DMN)-induced liver fibrosis in rats, ... and inhibition of cytokine signaling pathways (TGF- β and PDGF; Li et al., Transforming growth factor (TGF- β) and its intracellular signaling mediators, the Smad proteins, transduce signals via the classical JAK-STAT pathway and mediate ...

Effects of 18 α -glycyrrhizin on TGF- β 1/Smad signaling pathway in rats ...<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : Y Qu - 2015 - 被引用次数 : 7 - 相关文章

2015年2月1日 - Conclusion: 18 α -GL is able to attenuate CCl₄ induced liver fibrosis in rat. ... 18 α -GL improved liver fibrosis by acting on the TGF- β /Smad signal pathway, TGF- β mediates the activation of Smad2 primarily in undifferentiated ...

缺少字词 : notch concanavalin

Preventive Effect of Halofuginone on Concanavalin A-Induced Liver ...journals.plos.org/plosone/article?id=10.1371/journal.pone.0082232 - 翻译此页

作者 : J Liang - 2013 - 被引用次数 : 12 - 相关文章

2013年12月16日 - Since ConA-induced liver fibrosis is caused by the repeated activation of T ... by suppressing the synthesis of collagen I and inflammation-mediated liver injury. ... Concanavalin A