

Clinical Significance of Hypoechoic Submandibular Gland Lesions in Type 1 Autc



全部 图片 视频 新闻 购物 地图 图书

找到约 9,800 条结果

时间不限

过去 1 小时内
过去 24 小时内
过去 1 周内
过去 1 个月内
过去 1 年内

所有结果

精确匹配

[Hypoechoic Lesions in Submandibular Glands Are Diagnostic ...](#)

www.ncbi.nlm.nih.gov/pubmed/26658049

1. Pancreas. 2016 Jan;45(1):e8-9. doi: 10.1097/MPA.0000000000000442. ... Glands Are Diagnostic Markers of Type 1 Autoimmune Pancreatitis. Takano S(1) ...

[Hypoechoic Lesions in Submandibular Glands Are Diagnostic Ma ...](#)

journals.lww.com/pancreasjournal/.../Hypoechoic_Lesions_in_Submandibular_Glands_Are.29.aspx ▼

Hypoechoic Lesions in Submandibular Glands Are Diagnostic Markers of Type 1 Autoimmune Pancreatitis ... according to the International Consensus Diagnostic Criteria (ICDC) for AIP or the 2011 Japanese clinical diagnostic criteria for AIP.

[Pancreas - Wolters Kluwer Health](#)

journals.lww.com/pancreasjournal/.../Hypoechoic_Lesions_in_Submandibular_Glands_Are.29.aspx

Hypoechoic Lesions in Submandibular Glands Are Diagnostic Markers of Type 1 Autoimmune Pancreatitis The amendment of the Clinical Diagnostic Criteria in Japan JPS2011 in response to the proposal of the International Consensus of ...

[\[PDF\] Localized or Diffuse Lesions of the Submandibular Glands in ...](#)

www.jultrasoundmed.org/content/32/5/731.full.pdf ▼

salivary glands, pancreas, bile duct, lymph nodes, and kidneys.4–9 ... nal organs, such as autoimmune pancreatitis and sclerosing cholangitis. ... of IgG4-related disease and their clinical implications in ... (Table 1). Findings of the localized tumor-forming type were a hypoechoic coarse tumor bulging from the normal.

[IgG4-related Sclerosing Disease: Autoimmune Pancreatitis and ...](#)

pubs.rsna.org/doi/full/10.1148/rg.315105735 ▼

21 Mar 2011. Autoimmune pancreatitis is a distinct type of chronic pancreatitis and is

Name of journal: *World Journal of Gastroenterology*

Manuscript NO: 32412

Manuscript Type: ORIGINAL ARTICLE

Retrospective Study

Clinical significance of hypoechoic submandibular gland lesions in Type 1 autoimmune pancreatitis

Shinichi Takano, Mitsuharu Fukasawa, Makoto Kadokura, Hiroko Shindo, Ei Takahashi, Sumio Hirose, Yoshimitsu Fukasawa, Satoshi Kawakami, Tadashi Sato, Nobuyuki Enomoto

Abstract

Match Overview

1	Crossref 30 words Kuruma, Sawako, Terumi Kamisawa, Taku Tabata, Seiichi Hara, Takashi Fujiwara, Go Kuwata, Hideto Egarashira, Koi	1%
2	Crossref 22 words Ikeura, Tsukasa, Sönke Dellefsen, Giuseppe Zamboni, Riccardo Manfredi, Riccardo Negrelli, Antonio Amodio, France	1%
3	Crossref 14 words ERCP and EUS, 2015.	1%
4	Crossref 13 words Terumi Kamisawa. "Extraintestinal Manifestations of Inflammatory Bowel Disease: Autoimmune Pancreatitis and Ot...	1%

Clinical significance of hypoechoic submandibular gland lesions in Type 1 autoim

全部

图片

视频

新闻

地图

图书

找到约 65,100 条结果

时间不限

过去 1 小时内

过去 24 小时内

过去 1 周内

过去 1 个月内

过去 1 年内

所有结果

精确匹配

[Google 学术: Clinical significance of hypoechoic submandibular gland lesions in Type 1 autoimmune pancreatitis](#)

[Autoimmune pancreatitis](#) - Finkelberg - 被引用次数: 504

[... clinical guidelines for autoimmune pancreatitis](#) - Okazaki - 被引用次数: 164

[Recent advances in autoimmune pancreatitis](#) - Park - 被引用次数: 211

[Hypoechoic Lesions in Submandibular Glands Are ... - NCBI](#)

www.ncbi.nlm.nih.gov/pubmed/26658049

[Hypoechoic Lesions in Submandibular Glands Are ... - ResearchGate](#)

https://www.researchgate.net/.../286762491_Hypoechoic_Lesions_in_Submandibular_Glands_Are_Diagnostic_Markers_of_Type_1_Autoimmun...

Hypoechoic Lesions in Submandibular Glands Are Diagnostic Markers of Type 1 Autoimmune Pancreatitis on ResearchGate, the professional network for ...

[Hypoechoic Lesions in Submandibular Glands Are Diagnostic Ma ...](#)

journals.lww.com/pancreasjournal/.../Hypoechoic_Lesions_in_Submandibular_Glands_Are.29.aspx

Hypoechoic Lesions in Submandibular Glands Are Diagnostic Markers of Type 1 Autoimmune Pancreatitis. Takano, Shinichi MD, PhD; Sato, Tadashi MD, PhD; ...

[IgG4-related Sclerosing Disease: Autoimmune Pancreatitis and ...](#)

pubs.rsna.org/doi/full/10.1148/rg.315105735

21 Mar 2011 ... Autoimmune pancreatitis is a distinct type of chronic pancreatitis and is characterized by ... Kamisawa et al (1) proposed the concept of IgG4-related sclerosing disease and ... Salivary gland involvement in a 46-year-old man. ... Clinical significance of extrapancreatic lesions in autoimmune pancreatitis.

找到约 70,400 条结果 (用时 0.37 秒)

Google 学术 : Clinical significance of hypoechoic submandibular gland lesions in Type 1 autoimmune pancreatitis

Autoimmune pancreatitis - Finkelberg - 被引用次数 : 504

... clinical guidelines for autoimmune pancreatitis - Okazaki - 被引用次数 : 164

Recent advances in autoimmune pancreatitis - Park - 被引用次数 : 211

Hypoechoic Lesions in Submandibular Glands Are Diagnostic ... - NCBI

www.ncbi.nlm.nih.gov/pubmed/26658049 - 翻译此页

Hypoechoic Lesions in Submandibular Glands Are Diagnostic ...

https://www.researchgate.net/.../286762491_Hypoechoic_Lesions_in_Subma... - 翻译此页

Hypoechoic Lesions in Submandibular Glands Are Diagnostic Markers of Type 1 Autoimmune Pancreatitis on ResearchGate, the professional network for ...

Hypoechoic Lesions in Submandibular Glands Are Diagnostic Ma ...

journals.lww.com › Home › January 2016 - Volume 45 - Issue 1 - 翻译此页

作者 : S Takano - 2016 - 相关文章

Hypoechoic Lesions in Submandibular Glands Are Diagnostic Markers of Type 1 Autoimmune Pancreatitis. Takano, Shinichi MD, PhD; Sato, Tadashi MD, PhD; ...

IgG4-related Sclerosing Disease: Autoimmune Pancreatitis and ...

pubs.rsna.org/doi/full/10.1148/rg.315105735 ▼ 翻译此页

作者 : PA Vlachou - 2011 - 被引用次数 : 124 - 相关文章

2011年3月21日 - Autoimmune pancreatitis is a distinct type of chronic pancreatitis and is characterized by ... Kamisawa et al (1) proposed the concept of IgG4-related sclerosing disease and Salivary gland involvement in a 46-year-old man. Clinical significance of extrapancreatic lesions in autoimmune pancreatitis.

Localized or Diffuse Lesions of the Submandibular Glands in ...