

学术搜索

找到约 3,060 条结果 (用时0.04秒)

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☐ 包括专利☐ 包含引用

创建快讯

小提示: 只搜索**中文(简体)**结果, 可在 [学术搜索设置](#) 指定搜索语言**Safety of an ofloxacin - based antitubercular regimen for the treatment of tuberculosis in patients with underlying chronic liver disease: A preliminary report**

S Saigal, SR Agarwal, HP Nandeesh... - Journal of ..., 2001 - Wiley Online Library

... value of P less than 0.05 was considered to represent a statistically significant difference. ... Lack of relationship between hepatic toxicity and acetylator phenotypes in three thousand South Indian ... incidence of isoniazid hepatitis in rapid acetylators: possible relations to hydrazine ...

被引用次数: 62 相关文章 所有 5 个版本 引用 保存

Clinicopathological analysis of alcoholic liver disease complicating chronic type C hepatitis

O Noguchi, K Yamaoka, T Ikeda, S Tozuka... - Liver ..., 1991 - Wiley Online Library

... Fig. 2. Relationship between histological scores and Chiron C100 antibody. ... liver disease complicating NANB chronic hepatitis (Group B). There was no significant difference between Groups A ... In relation to the third possibility, chronic alcoholics are also known to be positive for ...

被引用次数: 20 相关文章 所有 5 个版本 引用 保存

Treatment with interferon-alpha (IFN [alpha]) of hepatitis C patients induces lower serum dipeptidyl peptidase IV activity, which is related to IFN [alpha]-induced ...

M Maes, S Bonaccorso, V Marino, A Puzella... - Molecular ..., 2001 - search.proquest.com

... serum DPP IV activity in relation to induction of the ... means of Pearsons product moment correlation coefficients. ... Eur Neuropsychopharmacol 2000; 10: 129132.24 Taylor MW, Feng G. Relationship between interferon-gamma, indolamine 2,3-dioxygenase, and tryptophan ...

被引用次数: 97 相关文章 所有 13 个版本 引用 保存

[引用] Serum neopterin levels in children with hepatitis-B-related chronic liver disease and its relationship to disease severity

EM Gulcan, I Tirit, A Anil, E Adal, G Ozbay - World J Gastroenterol, 2008

被引用次数: 13 相关文章 所有 12 个版本 引用 保存

[HTML] A review on liver function test

S Gowda, P Desai, V Hull, A Math... - Pan African ..., 2009 - panafrican-med-journal.com

... relationship between histological findings

[\[PDF\] researchgate.net](#)[\[PDF\] researchgate.net](#)[\[HTML\] panafrican-med-journal.com](#)

Name of journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 33386

Columns: ORIGINAL ARTICLE

Retrospective Study

Relationship between serum adenosine deaminase levels and liver histology in autoimmune hepatitis

Torgutalp M *et al.* Relationship between ADA and liver biopsy in AIH

Murat Torgutalp, Cumali Efe, Hakan Babaoglu, Taylan Kav

Murat Torgutalp, Cumali Efe, Hakan Babaoglu, Taylan Kav, Department of Gastroenterology, Hacettepe University Faculty of Medicine, Ankara, Turkey

Author contributions: Torgutalp M designed and performed the research and

Match Overview

1	Crossref 165 words Efe, Cumali, Mustafa Cengiz, Evrim Kahramanoğlu-Aksoy, Bülent Yılmaz, Burak Özşeker, Yavuz Beyazıt, Alpaslan Tan	5%
2	Internet 136 words crawled on 22-Jan-2016 www.wjgnet.com	4%

学术搜索

找到约 3,170 条结果 (用时0.13秒)

文章

我的图书馆

时间不限

2017以来

2016以来

2013以来

自定义范围...

按相关性排序

按日期排序

不限语言

中文网页

简体中文网页

☒ 包括专利

☒ 包含引用

☒ 创建快讯

小提示: 只搜索中文(简体)结果, 可在 学术搜索设置 指定搜索语言

Safety of an ofloxacin - based antitubercular regimen for the treatment of tuberculosis in patients with underlying chronic liver disease: A preliminary report

S Saigal, SR Agarwal, HP Nandeesh... - Journal of ..., 2001 - Wiley Online Library

... value of P less than 0.05 was considered to represent a statistically significant difference. ... Lack of relationship between hepatic toxicity and acetylator phenotypes in three thousand South Indian ... incidence of isoniazid hepatitis in rapid acetylators: possible relations to hydrazine ...

被引用次数: 62 相关文章 所有 5 个版本 引用 保存

Clinicopathological analysis of alcoholic liver disease complicating chronic type C hepatitis

O Noguchi, K Yamaoka, T Ikeda, S Tozuka... - Liver ..., 1991 - Wiley Online Library

... Fig. 2. Relationship between histological scores and Chiron C100 antibody. ... liver disease complicating NANB chronic hepatitis (Group B). There was no significant difference between Groups A ... In relation to the third possibility, chronic alcoholics are also known to be positive for ...

被引用次数: 20 相关文章 所有 5 个版本 引用 保存

Treatment with interferon-alpha (IFN [alpha]) of hepatitis C patients induces lower serum dipeptidyl peptidase IV activity, which is related to IFN [alpha]-induced ...

M Maes, S Bonaccorso, V Marino, A Puzella... - Molecular ..., 2001 - search.proquest.com

... serum DPP IV activity in relation to induction of the ... means of Pearsons product moment correlation coefficients. ... Eur Neuropsychopharmacol 2000; 10: 129132.24 Taylor MW, Feng G. Relationship between interferon-gamma, indolamine 2,3-dioxygenase, and tryptophan ...

被引用次数: 98 相关文章 所有 13 个版本 引用 保存

[引用] Serum neopterin levels in children with hepatitis-B-related chronic liver disease and its relationship to disease severity

EM Gulcan, I Tirit, A Anil, E Adal, G Ozbay - World J Gastroenterol, 2008

被引用次数: 13 相关文章 所有 12 个版本 引用 保存

Summary: A review on liver function test

[PDF] researchgate.net

[PDF] researchgate.net

Summary: nonafrican med journal.com

找到约 3,420,000 条结果 (用时 0.89 秒)

Google 学术 : Relationship between serum adenosine deaminase levels and liver histology in autoimmune hepatitis

... with underlying chronic liver disease: A preliminary ... - Saigal - 被引用次数 : 62

... liver disease complicating chronic type C hepatitis - Noguchi - 被引用次数 : 20

... IFN [alpha] of hepatitis C patients induces lower serum ... - Maes - 被引用次数 : 98

[Adenosine deaminase activity--an indicator of inflammation in liver ...

<https://www.ncbi.nlm.nih.gov/pubmed/3436285> - 翻译此页

作者 : R Nilius - 1987 - 被引用次数 : 7 - 相关文章

Inflammatory liver diseases display higher levels in serum ADA activity ... hepatitis, in active liver cirrhoses, extremely high levels in some liver tumours. Due to ... This can be validated by assessment of histological criteria. ... This interpretation is evidenced by correlative relations between the activity of ADA in liver biopsy ...

缺少字词 : autoimmune

Liver Involvement in Patients with Systemic Autoimmune Diseases

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : M COJOCARU - 2013 - 被引用次数 : 2 - 相关文章

Liver histology described in SAD is mostly based on needle biopsy. ... Differentiating features for autoimmune hepatitis (AIH) from SLE-related liver disease are ... the rise in serum alkaline phosphatase in view of the possible association between ... Abnormal ALT/AST levels developed in 14-35% of patients with RA initiating ...

缺少字词 : adenosine deaminase

[PDF] Adenosine deaminase activity in serum of patients with hepatitis - JMII

www.ejmii.com/article_pdf.php?code=PDT4b0e52bd97b2a - 翻译此页