

Bcl-2 degradation is an additional pro-apoptotic effect of polo-like kinase inhibit

全部 图片 视频 新闻 地图 图书

找到约 16,800 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

[Google 学术: Bcl-2 degradation is an additional pro-apoptotic effect of polo-like kinase inhibition in cholangiocarcinoma cells](#)

[Overview of major classes of plant-derived anticancer ...](#) - Amin - 被引用次数: 72  
[... activation in NSCLC H460 cells that involves RIP1 and ...](#) - Azijli - 被引用次数: 29  
[... of calmodulin in cell proliferation, programmed cell ...](#) - Berchtold - 被引用次数: 58

[Polo-Like Kinase 2 Is a Mediator of Hedgehog Survival Signaling in ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3811036/>

6 Aug 2013 ... **Polo-Like Kinase 2** Is a Mediator of Hedgehog Survival Signaling in ...  
**Cholangiocarcinoma (CCA) cells** paradoxically express the death ligand tumor ... Hh survival  
signaling by **inhibition** of Mcl-1 proteasomal **degradation**, ..... **Proapoptotic Effects** of PLK2  
**Inhibition** Are Mediated by Mcl-1 **Degradation**.

[Mcl-1 antibody | antibody review based on formal publications](#)

<https://www.labome.com/review/gene/.../Mcl-1-antibody.html>

In order to study promotion of survival of breast cancer **cells** in response to .... blot in order to  
investigate the regulatory **effect** of deamidation on **Bcl-xL degradation**. ... anti-Mcl-1 antibody  
was used in western blot to study dendritic **cell apoptosis** in the ..... "Pharmacological  
**inhibition** of **Polo-like kinase 1 (PLK1)** by BI-2536 ...

[Cooperative integration between HEDGEHOG-GLI signalling and ...](#)

<https://www.cambridge.org/core/product/.../core-reader>

9 Feb 2015 ... Recently, the **pro-apoptotic** protein Fem1b has been found to interact with  
mammalian Gli1 ... The serine/threonine unc-51-like **kinase 3 (ULK3)** enhances GLI1 (and  
GLI2) ... Modes of **action** of HH-GLI signalling in cancer ..... in vitro and in vivo through  
**inhibition** of the survival molecules AKT and **Bcl-2** (Ref.

[Normal Cells, but Not Cancer Cells, Survive Severe Plk1 Depletion](#)

[mcb.asm.org/content/26/6/2093.full](http://mcb.asm.org/content/26/6/2093.full)

We previously reported the phenotype of depletion of **polo-like kinase 1 (Plk1)** using RNA ... In  
HeLa **cells**, two weak hypomorphs showed only slight G2/M arrest, ... A **more** detailed analysis  
of the Plk1/p53 interaction reported recently showed ... Because **inhibition** of **cell proliferation**

Name of Journal: World Journal of Gastroenterology

Manuscript NO: 32539

Manuscript Type: ORIGINAL ARTICLE

*Basic Study*

**Bcl-2 degradation is an additional pro-apoptotic effect of polo-like kinase inhibition in cholangiocarcinoma cells**

Svenja Sydor, Sami Jafoui, Lena Wingerter, Sandra Swoboda, Joachim C Mertens,  
Guido Gerken, Ali Canbay, Andreas Paul, Christian D Fingas

**Abstract**

**AIM**

To examine the influence on apoptotic mechanisms following inhibition of polo-like

## Match Overview

1	<b>Internet</b> 104 words crawled on 09-Dec-2016 <a href="https://pdfs.semanticscholar.org">pdfs.semanticscholar.org</a>	3%
2	<b>Crossref</b> 69 words Sydor, Svenja, Yanli Gu, Martin Schlattjan, Lars P Bechmann, Ursula Rauen, Jan Best, Andreas Paul, Hideo A Ba	2%
3	<b>Crossref</b> 62 words Olympia Anastasiou, Svenja Sydor, Jan-Peter Sowa, Paul Manka, Antonios Katsounas, Wing-Kin Syn, Dagmar F	2%
4	<b>Crossref</b> 49 words Fingas, Christian D, Joachim C Mertens, Nataliya Razumilava, Svenja Sydor, Steven F Bronk, John D Christense	2%
5	<b>Internet</b> 32 words crawled on 14-Nov-2014 <a href="http://www.ncbi.nlm.nih.gov">www.ncbi.nlm.nih.gov</a>	1%
6	<b>Crossref</b> 23 words Boris R.A. Blechacz. "Sorafenib inhibits STAT3 signalin ... in cholangiocarcinoma cells by activating the phosphata	1%
7	<b>Crossref</b> 20 words Yang, Ming-Ming, Wei Huang, and Dian-Ming Jiang. "Tetramethylpyrazine protects Schwann cells from ischemia-l	1%
8	<b>Internet</b> 18 words crawled on 27-May-2010 <a href="http://www.biomedcentral.com">www.biomedcentral.com</a>	1%
9	<b>Internet</b> 17 words crawled on 18-Apr-2010	1%



Bcl-2 degradation is an additional pro-apoptotic effect of polo-like kinase inhibit

全部 图片 视频 新闻 地图 图书

找到约 22,400 条结果

时间不限

过去 1 小时内  
过去 24 小时内  
过去 1 周内  
过去 1 个月内  
过去 1 年内

所有结果

精确匹配

[Google 学术: Bcl-2 degradation is an additional pro-apoptotic effect of polo-like kinase inhibition in cholangiocarcinoma cells](#)

[Overview of major classes of plant-derived anticancer ...](#) - Amin - 被引用次数: 81

[... activation in NSCLC H460 cells that involves RIP1 and ...](#) - Azijli - 被引用次数: 29

[... anticancer properties through inhibition of kinase ...](#) - Lamoral-Theys - 被引用次数: 92

[Polo-Like Kinase 2 Is a Mediator of Hedgehog Survival Signaling in ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3811036/>

6 Aug 2013 ... **Cholangiocarcinoma** (CCA) cells paradoxically express the death ligand ... for the cell division regulating serine/threonine kinase **polo-like kinase 2** (PLK2). ... Hh survival signaling by inhibition of Mcl-1 proteasomal degradation, ..... **Proapoptotic Effects** of PLK2 Inhibition Are Mediated by Mcl-1 Degradation.

[Autocrine and Paracrine Mechanisms Promoting Chemoresistance ...](#)

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5297782/>

13 Jan 2017 ... **Cholangiocarcinoma** (CCA) is one of the most aggressive and lethal malignancies in humans. .... On the other hand, inactivation of Bax, a **pro-apoptotic** protein member of ... Similar to nucleolin, calmodulin exerts **inhibitory effects** on ..... the expression of the cell division regulator **polo-like kinase** (PLK)-2 , ...

[\[PDF\] Pololike kinase 2 is a mediator of hedgehog survival signaling in ...](#)

[onlinelibrary.wiley.com/doi/10.1002/hep.26484/pdf](https://onlinelibrary.wiley.com/doi/10.1002/hep.26484/pdf)

**Cholangiocarcinoma** (CCA) cells paradoxically express the death ligand tumor necrosis ... sion regulating serine/threonine kinase **polo-like kinase 2** (PLK2). .... **Additional Supporting Information** may be found in the online version of this article. .... **Proapoptotic effects** of PLK2 inhibition are mediated by Mcl-1 degradation.

[BCL2L1 | Cancer Genetics Web - CancerIndex](#)

[www.cancerindex.org/geneweb/BCL2L1.htm](http://www.cancerindex.org/geneweb/BCL2L1.htm)

20 Aug 2015 ... Gene: BCL2L1; **BCL2-like 1** ... **BCL-2** family members form hetero- or homodimers and act as anti- or **pro-apoptotic** regulators that are involved in a wide variety of cellular ... **Inhibitor** of Apoptosis Proteins; **Cell Proliferation**; Breast Cancer ... MATERIALS


[全部](#)
[图片](#)
[新闻](#)
[视频](#)
[购物](#)
[更多](#)
[设置](#)
[工具](#)

找到约 17,300 条结果 (用时 0.66 秒)

## Google 学术 : Bcl-2 degradation is an additional pro-apoptotic effect of polo-like kinase inhibition in cholangiocarcinoma cells

Overview of major classes of plant-derived anticancer ... - Amin - 被引用次数 : 81

... activation in NSCLC H460 cells that involves RIP1 and ... - Azijli - 被引用次数 : 29

... anticancer properties through inhibition of kinase ... - Lamoral-Theys - 被引用次数 : 92

## Polo-Like Kinase 2 Is a Mediator of Hedgehog Survival Signaling in ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - 翻译此页

作者 : CD Fingas - 2013 - 被引用次数 : 16 - 相关文章

2013年8月6日 - Cholangiocarcinoma (CCA) cells paradoxically express the death ligand ... for the cell division regulating serine/threonine kinase polo-like kinase 2 (PLK2). ... Hh survival signaling by inhibition of Mcl-1 proteasomal degradation, ..... Proapoptotic Effects of PLK2 Inhibition Are Mediated by Mcl-1 Degradation.

## Autocrine and Paracrine Mechanisms Promoting Chemoresistance in ...

<https://www.ncbi.nlm.nih.gov/pmc/articles/PMC5297782/> - 翻译此页

作者 : M Cadamuro - 2017

2017年1月13日 - Cholangiocarcinoma (CCA) is one of the most aggressive and lethal malignancies in humans. ... Mechanisms underpinning chemoresistance are multiple and affect drug .... On the other hand, inactivation of Bax, a pro-apoptotic protein ..... the expression of the cell division regulator polo-like kinase (PLK)-2, ...

## Pololike kinase 2 is a mediator of hedgehog survival signaling in ...

[onlinelibrary.wiley.com/doi/10.1002/hep.26484/pdf](https://onlinelibrary.wiley.com/doi/10.1002/hep.26484/pdf) - 翻译此页

作者 : CD Fingas - 2013 - 被引用次数 : 16 - 相关文章

Cholangiocarcinoma (CCA) cells paradoxically express the death ligand tumor necrosis ... sion regulating serine/threonine kinase polo-like kinase 2 (PLK2). .... Additional Supporting Information may be found in the online version of this article. .... Proapoptotic effects of PLK2 inhibition are mediated by Mcl-1 degradation.