

[全部](#)[图片](#)[新闻](#)[视频](#)[更多 ▾](#)[搜索工具](#)

找到约 4,230 条结果 (用时 0.63 秒)

[Google 学术: VARIABILITY OF ANTI-HUMAN TRANSGLUTAMINASE TESTING ACROSS MEDITERRANEAN COUNTRIES](#)

The prevalence of celiac disease in Europe: results of a ... - [Mustalahti](#) - 被引用次数: 409

Serologic and genetic markers of celiac disease: a ... - [Bonamico](#) - 被引用次数: 85

... Lymphoma Patients: A Serum Anti-Transglutaminase- ... - [Carroccio](#) - 被引用次数: 18

[Testing for gluten-related disorders in clinical practice: The role of ...](#)

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: [D Armstrong](#) - 2011 - 被引用次数: 30 - [相关文章](#)

Antitissue [transglutaminase \(anti-tTG\) tests](#) followed and, more recently, ... The ability of [human](#) leukocyte antigen (HLA) typing for HLA-DQ2 and HLA-DQB ... The AGA-IgG [test](#) has sensitivity from 42% to 93%, and specificity in the [region](#) of 84% to 97%. ... Serological [tests: Variability](#) in celiac disease [Annu Rev Med](#).

[Performance of Antibodies against Tissue Transglutaminase for the ...](#)

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: [E Zintzaras](#) - 2006 - 被引用次数: 64 - [相关文章](#)

[Human](#) tTG ELISA is sensitive and specific, and it can be used for mass screening. ... OR [anti-tTG](#) OR tTG) AND ([sensitivity and specificity] OR diagnostic [test](#)). ... The [area](#) under the SROC curve (AUC) is related to the common log(OR), and it ... The SROC curve can be fit weighted by the inverse of the [variance](#) of the ...

[Antibodies to human recombinant tissue transglutaminase may detect ...](#)

Name of Journal: *World Journal of Gastroenterology*

ESPS Manuscript NO: 31493

Manuscript Type: ORIGINAL ARTICLE

Observational Study

Variability of anti-human transglutaminase testing across Mediterranean countries

Smarrazzo A *et al.* Transglutaminase testing across Mediterranean countries

Andrea Smarrazzo, Giuseppe Magazzù, Mongi Ben-Hariz, Maria Legarda Tamara, Virtut Velmishi, Eleftheria Roma, Aydan Kansu, Dušanka Mičetić-Turk, Enzo Bravi, Pio Stellato, Carmela Arcidiaco, Luigi Greco

Abstract

AIM

To verify the precision and accuracy of transglutaminase antibodies (TGA) assays across

Match Overview

1	Internet 77 words crawled on 08-Jul-2016 journals.lww.com	2%
2	Internet 28 words crawled on 20-Nov-2014 archive.ahrq.gov	1%
3	Crossref 16 words Bienvenu, Françoise, Cécile Besson Duvanel, Camille Seignover, Paul Rouzaire, Alain Lachaux, and Jacques Bie	<1%

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 9,490 条结果 (用时 0.64 秒)

Google 学术: Variability of anti-human transglutaminase testing across Mediterranean countries

The prevalence of celiac disease in Europe: results of a ... - Mustalahti - 被引用次数: 428

... of four tissue transglutaminase autoantibody tests ... - Martini - 被引用次数: 33

Serologic and genetic markers of celiac disease: a ... - Bonamico - 被引用次数: 86

Serologic testing in celiac disease - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: M Rashid - 2016 - 被引用次数: 11 - [相关文章](#)

Testing for human leukocyte antigens DQ2 and DQ8 can help exclude the diagnosis. ... a positive immunoglobulin A (IgA) tissue transglutaminase (tTG) antibody test leading to potential interobserver variability in interpretation of the test results. ... self-testing kit for CD has been available in pharmacies across the country ...

Testing for Antireticulin Antibodies in Patients with Celiac Disease Is ...

<https://www.ncbi.nlm.nih.gov> > NCBI > Literature > PubMed Central (PMC) - [翻译此页](#)

作者: SL Nandiwada - 2013 - 被引用次数: 9 - [相关文章](#)

EMA is detected by IFA using monkey esophageal or human umbilical cord tissue, ... The anti-DGP antibody test provides a more specific marker for disease ... tTG IgG tests are not dependable due to the variability in their sensitivities. ... BMC Med. Antibodies against human tissue transglutaminase and endomysium in ...

Anti-tissue transglutaminase antibodies in the follow-up of adult ...

[全部](#)[图片](#)[新闻](#)[视频](#)[购物](#)[更多](#)[设置](#)[工具](#)

找到约 10,300 条结果 (用时 0.63 秒)

Google 学术 : Variability of anti-human transglutaminase testing across Mediterranean countries

The prevalence of celiac disease in Europe: results of a ... - [Mustalahti](#) - 被引用次数 : 431

... of four tissue [transglutaminase](#) autoantibody tests ... - [Martini](#) - 被引用次数 : 33

Serologic and genetic markers of celiac disease: a ... - [Bonamico](#) - 被引用次数 : 86

Serologic testing in celiac disease - NCBI - National Institutes of Health

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者 : [M Rashid](#) - 2016 - 被引用次数 : 11 - [相关文章](#)

[Testing](#) for [human](#) leukocyte antigens DQ2 and DQ8 can help exclude the diagnosis. ... a positive immunoglobulin A (IgA) tissue [transglutaminase](#) (tTG) antibody [test](#) leading to potential interobserver [variability](#) in interpretation of the [test](#) results. ... self-[testing](#) kit for CD has been available in pharmacies [across](#) the [country](#) ...

Testing for Antireticulin Antibodies in Patients with Celiac Disease Is ...

<https://www.ncbi.nlm.nih.gov> > [NCBI](#) > [Literature](#) > [PubMed Central \(PMC\)](#) - [翻译此页](#)

作者 : [SL Nandiwada](#) - 2013 - 被引用次数 : 9 - [相关文章](#)

This minireview highlights the advances in CD-specific serologic [testing](#) and ... and more-specific serologic [tests](#) such as the tissue [transglutaminase](#) (tTG), EMA is detected by IFA using monkey esophageal or [human](#) umbilical cord ... The [anti-DGP](#) antibody [test](#) provides a more specific marker for disease [BMC Med.](#)

Anti-tissue transglutaminase antibodies and their role in the ...

acb.sagepub.com/content/43/2/105.full.pdf - [翻译此页](#)

作者 : [PG Hill](#) - 2006 - 被引用次数 : 61 - [相关文章](#)

immunoassays using [human](#) tissue TGA as antigen have high diagnostic sensitivity ... adopting IgA